

SOCIETY OF AMERICAN BACTERIOLOGISTS

*MONOGRAPHS ON SYSTEMATIC
BACTERIOLOGY*

Volume I GENERAL SYSTEMATIC BACTERIOLOGY
 R. E. Buchanan, *Iowa State College*
 597 pages. \$6.00, net postpaid

**GENERAL
SYSTEMATIC BACTERIOLOGY**

MONOGRAPHS ON SYSTEMATIC BACTERIOLOGY

GENERAL SYSTEMATIC BACTERIOLOGY

History, Nomenclature, Groups of Bacteria

BY

R E BUCHANAN, PH.D

*Professor of Bacteriology, Bacteriologist of the Iowa Agricultural Experiment Station
and Dean of the Graduate College, Iowa State College,
Ames, Iowa*



BALTIMORE
WILLIAMS & WILKINS COMPANY
1925

COPYRIGHT 1925
WILLIAMS & WILKINS COMPANY

Made in United States of America

Published January, 1925

ALL RIGHTS RESERVED

COMPOSED AND PRINTED AT THE
WAVERLY PRESS
BY THE WILLIAMS & WILKINS COMPANY
BALTIMORE, MD , U S A

CONTENTS

Preface	9
I Classification of the Genera and Higher Groups of Bacteria Historical, 1773-1922	15
II Codes of Nomenclature and Their Application in Bacteriology	109
The Rules of Bacterial Nomenclature According to Lehmann and Neumann	110
The International Code of Botanical Nomenclature	113
The International Code of Zoological Nomenclature	137
The Type Basis Code Suggested by the Committee on Botanical Nomen- clature of the Botanical Society of America	146
Suggestions and Recommendations of the Society of American Bacteri- ologists	149
III Nomenclatural Status of Names Which Have Been Applied to Groups of Bacteria of Higher Rank than Species Alphabetically Arranged	152
Bibliography	533
Subject Index	561
Author Index	589

PREFACE

The present volume is the first of a series of monographs in the general field of systematic bacteriology. The data presented have been collected, compiled and annotated largely in connection with courses of lectures in systematic bacteriology given to graduate students in bacteriology at the Iowa State College during the past fifteen years. It is an attempt to bring together material which may prove of value to those who are desirous of knowing the probable nomenclatural status of the various names which have been used in bacterial terminology

If science is to be defined as a system of *classified* knowledge, the subject of bacteriology is laboring under a serious handicap in lacking, probably more than any other branch of science, the advantages conferred by a satisfactory system of terminology or of nomenclature. It would seem that systematic bacteriology is deserving of even more recognition at the hands of teachers and investigators than has been accorded to it in the past.

Systematic bacteriology has two principal aims or functions which are of importance to the teacher and to the investigator. The first of these aims has to do with the presentation in graphic form of our present conception of the phylogeny and of the relationships of various groups of bacteria. The second is to give a greater degree of stability to the names used for particular groups of organisms and to prevent unnecessary nomenclatural confusion in literature. These aims are thoroughly scientific, their accomplishment will prove useful, we must therefore accord them careful consideration if our nomenclature is to be stabilized.

It is somewhat difficult to tabulate satisfactorily all of the factors instrumental in bringing about the present state of confusion and uncertainty in bacterial terminology. That there is such confusion is, of course, a self-evident fact. It has been rare in texts or articles (until comparatively recently) to find a consistent use made of generic names, and systems of classification taught in elementary courses in bacteriology have rarely been made use of later. In some quarters there are at the present time certain ill-advised attempts being made

to standardize nomenclature. Certain American publishers for example have sought to fix upon the correct generic names for the designation of certain microorganisms, but apparently with no clear conception of the rules which should govern such standardization.

A study of the catalogs of various colleges and universities in the United States in which bacteriology is being taught at the present time reveals the fact that in very few cases are courses offered in which there is any emphasis placed upon systematic bacteriology and phylogeny. An examination of the courses offered in other biological sciences would show very different conditions. Another factor which has undoubtedly tended to confuse nomenclature in this science is the fact that until recent years most bacteriologists received their training in medical schools. During their medical course they were familiarized with several systems of nomenclature used in other sciences. In anatomy, for example, agreement is rapidly being reached that for each organ and anatomical unit a single international Latin name shall be adopted, and the student becomes familiar with the general rules for the formation of such names. In pathology and medicine diseases are designated in much the same fashion, a descriptive Latin phrase, frequently consisting of several words, is used. It is exceptional for the teacher in bacteriology to emphasize the fact that systems of names used in anatomy and pathology are utterly different and based upon other rules than those used in naming living plants and animals. Many of the polynomials which cumber our bacteriological literature have been applied by workers who do not have such distinctions in mind.

By far the most important factor which has determined the attitude of teachers and investigators toward this problem has been the lack of a really adequate system of group differentiation, of classification. It has taken bacteriological systematists a long time to realize that such group differentiation (the separation of species and genera), upon other than morphological differences, may be wholly valid. The botanical bacteriologist has ignored physiology which is so important. The physiological bacteriologist has, in many cases, not possessed the background of the systematist. The latter worker may point out the characters which should be used, but finds difficulty in their practical application to the specific problems of taxonomy.

The most hopeful sign of importance in this respect probably, has been the work of the committee on taxonomy of bacteria of the Society of American Bacteriologists under the chairmanship of Dr Winslow,

and of the more recent work of a committee on classification of bacteria under the chairmanship of Dr Bergey. The work of Mrs Enlows in the names of bacterial genera is also to be commended. It is to be expected that, as a result of their work, eventually a practical system of nomenclature which will be satisfactory and applicable to all fields of bacteriology will be evolved

But, it may be asked, is a scientific classification of bacteria really important and desirable? Are we not making progress in our science quite as rapidly as we would without the increased emphasis upon taxonomy? The following statement by H. Marshall Ward, published two decades ago is apropos

"The only really valid objection to a purely scientific classification is the old objection of the purely utilitarian practical man, and even there the objection is relative. This leads me to bring out the point that the bacteriologists in the widest sense of the word are really looking at the question of classification from at least two very different points of view. On the one hand, we have the botanists, who direct their attention to the organism, the *Schizomycete* itself, as a biological phenomenon to be examined and reported upon as thoroughly as possible, for them no classification is complete which does not record, or (which amounts to the same thing) imply in its records, all of the life phenomena of the organism, including its pedigree

"On the other hand, we have the pathologists, hygienists, brewers, chemists, etc., who regard the organism simply as an object to be named for convenience in reference, because it brings about certain changes in tissues, waters, and other media which they are more especially concerned with. They do not care, and naturally so, what vagaries the organism exhibits, so long as they can recognize it when they meet with it. As a matter of experience, however, it is just these vagaries that bring about the sources of error which beset them on all hands, and hence they are equally interested with the botanist in having them cleared up and explained. When we come to the conclusion that, whatever may be believed to the contrary, the real interests of bacteriologists of all kinds are identical."

The first chapter of the present volume is devoted to a brief history of the various classifications of bacteria that have been proposed up to the present time. An effort has also been made to give dichotomous keys to the genera and higher groups recognized by the various writers whenever there seemed to be sufficient data for their formulation. Contrary to commonly accepted opinion, it will be found that a large

number of observers have made contributions in this field, but the writings are so widely scattered and, in many cases, the journals and other publications so difficult of access that for the most part they are quite unknown to our modern writers or, at least, unheeded by them. It is believed that a synopsis of these various schemes of classification that have been proposed and a comparison of the various bases upon which these rest must surely be of service in developing a stable bacteriological nomenclature

The second chapter has to do with codes of nomenclature and their relationships to the problems of bacterial terminology. The subject of nomenclature is, in part, distinct from that of classification. The latter deals with methods and criteria of use in the differentiation of groups from each other. The former has for its function the determination of the kind of a name that shall be applied to a group and the validity and suitability of names which have already been applied. This frequent confusion of nomenclature in classification has made us hesitate to give a name to that which we cannot accurately define. It is true, however, that species have not been accurately delimited in the older biological sciences, but this fact has not interfered with at least tentative arrangements of the genera and higher groups nor of application of correct names. It seems to be self-evident that until the bacteriologists can agree upon a code and follow it consistently, there is little hope or remedy for our present chaos. It is believed that the discussion of the International Botanical Code given in this second section will show that it may constitute an adequate basis, which, with some modification, may prove satisfactory as a guide.

The third chapter is an alphabetical list of all of the names which have been used by various authors to designate bacterial subgenera, genera or higher groups so far as a rather comprehensive study of the literature has revealed them. Wherever practicable the original description has been quoted, the descriptions used by various subsequent writers discussed, and sufficient data included to make possible in most cases, at least, the formation of an opinion as to the probable nomenclatural status of each name. It is believed that this list will prove helpful in the determination of the exact meaning of the various names, in the formulation of adequate diagnoses for the determination of synonymy, and in preventing the introduction of new names where there are valid older names.

The section on bibliography has been compiled with particular care, as have also the various indices.

I wish gratefully to acknowledge the assistance of many members of the staff in bacteriology at the Iowa State College, especially Miss Clarissa Clark, Mrs C. H. Werkman, Miss Ruth Confare, Mr. J. C. Weldin and Dr Max Levine. For interest and assistance in bibliographic details acknowledgment is due Miss Margaret Brown and Prof. C. H. Brown of the library of Iowa State College.

CHAPTER I

CLASSIFICATION OF THE GENERA AND HIGHER GROUPS OF BACTERIA

Apparently the first definite named description of any organism included now among the bacteria is probably that of Mueller in his *Vermium terrestrium et fluviatilium* (1773). The following key, abridged from his introduction, gives the main groupings and the genera of the group in which the bacteria fall.

Mueller's Classification of Vermes (1773)

- a. Tentaculi destituta.
 - b. Vagantes *Infusoria*
 - c. Organus externis nullis
 - d. Teretes.
 - 1. Corpus punctiforme *Monas*
 - 2. Corpus sphaericina *Volvox*
 - 3. Corpus cylindraceum *Encheus*
 - 4. Corpus elongatum *Vibrio*
 - 5. Corpus cavum *Bursaria*
 - dd. Complanati (With four genera)
 - cc. Organus externus (With four genera)
 - bb. Serpentes *Helminthica*
(With eight genera)
- aa. Tentaculis instructi *Testacea*
(With twelve genera)

Later in the introduction to the above volume the two genera *Monas* and *Vibrio* (which contain his bacterial species) are noted as found among the *Aquatici*, also among the *Hyalini*, also under *Tentaculis nullis*, *inconspicui*, *teretes*, and still later under *Vacillantes*, *pellucidi*. Finally he places these genera with *Volvox* and *Encheus* in a classification under the headings *Organus externis nullis*, *crassiuscula*.

Three species of *Monas* are described. The generic description is "*Vermis inconspicuus, simplicissimus pellucidus, punctiformus*." The genus *Vibrio* is described as "*Vermis inconspicuus, simplicissimus, teres, elongatus*." Fifteen species are described. Several of them were doubtless bacterial forms.

In the year 1786 there appeared a posthumous volume by Mueller, entitled "*Animalcula Infusoria fluviatilia et marina*." The infusoria

were divided into two groups "*Organis externus nullus*" and "*Organis externus*." The forms now included among the bacteria were all to be found in the ten genera of the first group. These were separated as follows (p. XXVI)

Mueller's Classification of Protozoa, etc. (1786)

Crassiuscula

- 1 *Monas* Punctiforme
- 2 *Proteus*: Mutabile
- 3 *Volvox* Sphaericum
- 4 *Enchelis* Cylindraceum
- 5 *Vibrio*. Elongatum

Membranacea

- 6 *Cyclidium* Ovale
- 7 *Paramoecium*. Oblongum
- 8 *Kolpoda* Simatum
- 9 *Gonium*: Angulatum
- 10 *Bursaria* Cavum

Two of these genera contained organisms now included among the bacteria. In the genus *Monas* were ten species and in *Vibrio* thirty-one species. Of these very probably *Monas termo*, *M. punctum* and *M. lens*, *Vibrio lineola*, *V. rugula*, *V. bacillus*, *V. undula*, *V. serpens* and *V. spirillum* were bacteria. The monads were spherical or ovoid in shape, the vibrios longer, and in most species spiral.

Within the next two decades, no advance was made in the classification of the forms now known as bacteria. However, three generic names were created by as many writers for organisms supposed to be fungi, but which many decades later were shown to be bacteria. The first of these was named by Link (1795), who described *Polyangium vitellinum*, an organism in its fruiting stages bearing a close resemblance to a mold, but which was later shown by Thaxter (1892) to belong to the *Mycobacteriaceae*. The second name was applied to the species *Serratia marcescens* by Bizio (1823). This investigator had undertaken to study the cause of an outbreak of "bloody" polenta which was causing much superstitious fear among the peoples of northern Italy. He attributed the difficulty to a fungus which developed a red pigment on farinaceous foods. This was probably the organism now more commonly termed *Bacillus prodigiosus*. Somewhat later Sette (1824, p. 51) named what is apparently the same organism *Zoogalactina metroptha*, also regarding it as one of the fungi.

Persoon (1822, p 96) used the designation *Mycoderma mesentericum* for the membrane which appears upon the surface of fermenting alcoholic solutions, consisting largely of acetic acid bacteria and yeasts. The organism was classed with the fungi, its relationship to the *Infusoria* of Mueller was not at the time apprehended.

Bory de St. Vincent (1824) created the genus *Melanella* to include all non-flexible organisms of the group of *Vibriomides* whether straight or spiral. To this genus he transferred four of Mueller's species of *Vibrio*. (*V. lineola*, *V. rugula*, *V. bacillus* and *V. spirillum*)

Bory de St Vincent (1826) prepared a "Tableau des Ordes, des Familles et des Genres de Microscopiques". This was a dichotomous key to genera. A translation of the key carried out to include only the forms now listed with the bacteria is as follows:

Bory de St Vincent's Classification (1826)

- a Without Testa
 - b Perfectly smooth, without cilia or other vibratile organs
 - Order *Gymnode*
 - c Without appendages
 - d Never linear or wormlike (5 families)
 - dd Linear or wormlike
 - Family *Vibriomides*
 - e Body opaque
 - f Coiled in a discoid spiral
 - Spirulina*
 - ff Not coiled in a discoid spiral
 - Melanella*
 - ee Body transparent (diaphanous)
 - f Attenuate at both tips
 - Vibrio*
 - ff Obtuse
 - g With one end enlarged
 - Lacrimatoria*
 - gg Not enlarged at one end
 - Pupella*

Von Baer (1827, p 748) proposed to establish a genus *Lineola* to include all of the extremely minute *Infusoria*.

The first considerable contributions to our knowledge of bacteria and their classifications, following Mueller, were those of Ehrenberg (from 1828-1838). Certain of his genera of the *Infusoria* were made up almost entirely of organisms now included with bacteria. The following key will indicate the principal groups of the *Infusoria* and the position of the "bacterial" genera in the system.

Ehrenberg's Classification of the Infusoria (1838)

- a Forms which ingest solid particles, as carmine
Polygastrica
- aa Forms which never ingest particles
 - b Family *Monadina* Genus *Monas*, of which *Monas crepusculum* was probably a bacterium
 - bb Family *Vibrionia* Forms which tend to form filaments as a result of incomplete fission
 - c Cells not flattened
 - d Cells not flexuous
 - e Forming straight rods *Bacterium*
 - ee Forming spiral rods *Spirillum*
 - dd Cells flexuous
 - e Straight rods *Vibrio*
 - ee Spiral rods *Spirochaeta*
 - cc Cells in form of a flattened spiral *Spirodiscus*

He also described as a member of the diatom genus *Gaillonella*, a *G. ferruginea* now generally included among the bacteria. Later he changed the spelling to *Gallionella*. In spite of the fact that it is that of a recognized diatom genus, this name is still frequently incorrectly used as a bacterial generic name in the designation of this iron organism. In the form finally assumed by the Ehrenberg classification (1838) the bacterial species recognized were as follows: *Bacterium triloculare*, *Vibrio lineola*, *V. tremulans*, *V. subtilis*, *V. rugula*, *V. prolifer* and *V. bacillus*, *Spirillum undula*, *S. tenue*, *Spirochaeta plicatilis* and *Spirodiscus fulvus*. Later (1840, p. 202) he described *Vibrio synxanthus* and *V. synxanthus*.

Kützing (1833, p. 385) described as a fresh water alga a form which he termed *Sphaerotilus natans*. This is now generally included among the filamentous water or iron bacteria.

Dujardin (1841, p. 209) used the family designation *Vibrionia* to include "Animaux sans organes locomoteurs visibles, se mouvent par l'effet de leur contractibilité générale". The genera recognized may be differentiated as follows:

Dujardin's Classification of *Vibrionia* (1841)

- A Cells straight, non-flexuous, more or less definitely jointed, with slow wave-like motility *Bacterium*
- B Cells straight or bent, more or less definitely jointed, flexuous *Vibrio*
- C Filamentous forms spiral or screw shaped, never straight, and revolving rapidly on the axis *Spirillum*

Trevisan (1842, p 56) described an organism which he placed among the algae as closely related to *Oscillatoria*. To it he gave the name *Beggiatoa punctata*. This genus is now generally recognized as belonging among the sulphur bacteria. In the same year Goodsir (1842, p. 432) described an organism which he named *Sarcina ventriculi*. It occurred as masses of spherical cells forming regular packets in the vomit of a patient. The relationship of this organism to the types included by previous writers among the Infusoria was not recognized.

Kuetzing (1843, p 198) described one of the iron bacteria, *Leptothrix ochracea*, which may be regarded as the type species of this genus. Later Robin (1853, p 345) named one of the mouth bacteria *Leptothrix buccalis*. We know now that the two species named are not at all closely related, but the double use of this designation has led to much confusion in literature in recent years. Sanitary bacteriologists in general use the genus *Leptothrix* for iron bacteria, while many medical bacteriologists use the name for certain parasitic mouth forms.

Leidy (1849-1852) described several genera of "*Entophyta*" from the intestines of arthropods. Several of these, probably *Arthromitus*, *Cladophytum*, *Corynocladus*, *Cryptodesma*, *Eccrina*, and *Enterobryus*, may tentatively be included with the bacteria. It is noteworthy that in 1849 this author called attention to the probable relationships of the *Vibrio* and *Bacterium* to plants, that is to the *Entophyta*.

Perty (1852) in his "*Zur Kenntniss kleinster Lebensformen*" divided the *Infusoria* into *Ciliata* and *Phytozoidia*, the latter in turn he divided into three sections, the third of which he termed *Lampozoridae*. The family *Vibrionida* is included under this. The subfamilies and genera of the *Vibrionida* may be differentiated as follows

Perty's Classification of the Vibrionida (1852)

- | | | |
|---------------------------------------|-------------|--------------------|
| a. Chains or filaments spirally wound | | |
| | Subfamily 1 | <i>Spirillum</i> |
| 1 Cells not flexuous | | <i>Spirillum</i> |
| 2 Cells flexuous | | <i>Spirochaeta</i> |
| aa Filaments flexuous or straight | Subfamily 2 | <i>Bacterina</i> |
| 1 Cells flexuous | | <i>Vibrio</i> |
| 2 Cells not flexuous, cells isolated | | <i>Bacterium</i> |
| 3 Cells not flexuous, in filaments | | <i>Metallacter</i> |
| 4 Cells not flexuous, filaments | | <i>Sporonema</i> |

Three species of *Spirillum*, one of *Spirochaeta*, three of *Vibrio*, one of *Bacterium*, one of *Metallacter* and one of *Sporonema* were described.

Griffith (1853, p 438) recognized the fact that the *Gallionella ferruginea* of Ehrenberg was not a diatom, and created for it a new genus *Didymohelia*

Ferdinand Cohn (1854, p 123) described as a genus the slimy bacterial growth which develops in certain solutions, giving it the name *Zoogloea*. Later (1872) he abandoned this term recognizing this to be simply a growth stage in the developmental cycle Cohn's most important conclusion at this time was to the effect that the bacteria (*Vibrionien*) belong in the plant kingdom rather than with the true *Infusoria*

Henfrey (1856, p 53) described as an alga a form which he termed *Clathrocystis roseo-persinica* This was later placed among the sulphur bacteria by Winogradsky (1888)

Naegeli (1857, p 760) first created a definite group in the plant kingdom for the bacteria He united the genera *Bacterium*, *Vibrio*, *Spirillum*, *Sarcina*, *Umbina* and *Nosema* under the group *Schizomycetes*, the name by which the bacteria have in general since been known

Berkeley (1857, p 313) in his Introduction to Cryptogamic Botany included two new genera among the molds, *Chondromyces* and *Stigmatella* The work of Thaxter has shown these forms to be the fruiting bodies of myxobacteria

In 1865 (p 156) Trecul studied carefully certain vegetable cells which found developing in putrefying plant tissues To the various shapes he gave the names *Urocephalum*, *Amylobacter* and *Clostridium* These terms have been sometimes used by subsequent writers as generic names and ascribed to Trecul, but this ascription is not, strictly speaking, correct, for this author named no species, his names can be regarded only as pseudogeneric

Rabenhorst (1865) in Sectio II, Algas Phycochromaceas Complectens of his Flora Europaea Algarum included a number of bacterial genera with the blue green algae All are included in his order *Nematogenae* Family *Oscillariaceae* The following key to the three subfamilies and to the genera of the subfamily *Spirillineae* will serve to show this author's conception of relationships

Rabenhorst's Classification of Bacteria (1865)

- A Filaments more or less destitute of color, flexuous or spirally bent, not sheathed, often embedded in mucus

Subfamily 1 *Spirillineae*

- I Filaments short, cylindric or filiform 1 *Vibrio*
- II Filaments filiform, articulated, spirally twisted 2 *Spirillum*
- III Filament spiral, much elongated 3 *Spirochaeta*
- B Filaments colored (often only pale blue) sheathed, usually not moving, rarely slowly oscillating Subfamily II *Leptothricheae*
One genus of the two included, *Leptothrix*, is often classed with the bacteria
- C Filaments more or less colored, sheathed, often showing active oscillation Subfamily III *Oscillariaceae*
One genus, *Beggiatoa* is now generally classed with the bacteria

As a footnote under the heading "*Fungorum genera Leptothricheis et Oscillarins quasi affinis*" the following genera are included *Cryptococcus*, *Ulvina*, *Sphaerotilus* and *Hygroscopicus*

In the genus *Vibrio* the following species are listed, *V. lineola*, *V. tremulans*, *V. subtilis*, *V. rugula*, *V. profluer*, *V. serpens*, and *V. bacillus*. In the genus *Spirillum* are included *Sp. undula*, *Sp. tenue*, *Sp. volutans* and *Sp. rufum*. The single species of *Spirochaeta* is *Sp. plautii*.

Hallier (1866) introduced the theory of pleomorphism as applied to microorganisms. As a result of his work no additional knowledge of bacterial classification was developed, indeed it was probably somewhat instrumental in retarding advance. No generic names are to be ascribed to him, although he introduced certain names to designate what he believed to be growth forms, and which were later used by various authors as genera.

Davaine (1868) concluded as a result of his study of the anthrax organism that this microbe was closely related to the other bacteria, but unlike those previously described, non-motile. To this type he gave the generic name *Bacteridium*. A key to the genera of bacteria which he recognized is as follows:

Davaine's Classification of Bacteria (1868)

- a Filaments straight or bent, never spiral
 - b Motile
 - c Cells rigid *Bacterium*
 - cc Cells flexuous, sinuous *Vibrio*
 - bb Non-motile *Bacteridium*
- aa Filaments spiral *Spirillum*

Cohn (1870, p 108) described one of the filamentous iron bacteria as *Crenothrix polyspora*. Klebs (1871) in conformity with his ideas on bacterial pleomorphism gave the name *Microsporion septicum* to the pleomorphic organism which he associated with pus infections.

Cohn (1872, p 229) named a spherical organism which he found in smallpox lymph the *Microsphaera vaccinae*. Later the fact that this generic name had been previously used for a genus of *Ascomycetes* led him to change to the generic name *Micrococcus*.

The first measurably adequate classification of bacteria following the recognition of the fact that they were plants, not animals, was that of Cohn (1872, p 127). He defined the bacteria as having cells free from chlorophyll, spherical, oblong or cylindric, straight or bent, multiplying exclusively by fission and either isolated or vegetating in cell families. Four tribes with a total of six genera were recognized. They may be differentiated by reference to the following key:

Cohn's Classification of Bacteria (1872)

A Cells spherical	Tribe I	SPHAEROBACTERIA
One genus only		1 <i>Micrococcus</i>
B Cells isolated short rods	Tribe II	MICROBACTERIA
One genus only		2 <i>Bacterium</i>
C Cells elongate or in filaments	Tribe III	DESMOBACTERIA
1 Cells not flexuous		3 <i>Bacillus</i>
2 Cells flexuous		4 <i>Vibrio</i>
D Cells spiral	Tribe IV	SPIROBACTERIA
1 Cells not flexuous		5 <i>Spirillum</i>
2 Cells flexuous		6 <i>Spirochaete</i>

It will be noted that one new genus, *Bacillus*, was created with *B. subtilis* as the type. Two other species, *Bacillus ulna* and *B. anthracis* were included.

Billroth (1874) in a volume entitled "Untersuchungen über die Vegetationsformen von *Coccobacteria septica*," revived in a modified form the Haller theory of pleomorphism. As the result of extended studies he came to the conclusion that various bacteria which he observed were different growth forms of the organism which he termed *Coccobacteria septica*. To the various growth forms he gave names, such as *Micrococcus*, *Mesococcus*, *Megacoccus*, *Microbacteria*, *Mesobacteria*, *Megabacteria*, *Monococcus*, *Monobacteria*, *Petalococcus*, etc. With a modified spelling some of these names have been used by subsequent writers as generic names for bacteria, and by some have been credited to Billroth. This is incorrect as the names were not used by this author in a generic sense, and had no species named.

Cohn (1875, p 141) later greatly extended and improved his classification of bacteria. He concluded that they belonged with the blue green algae, differing principally in the absence of chlorophyll. His complete classification is as follows:

Cohn's Classification of the Schizophyta (1875)

a Cells free or united into slimy families by an intercellular substance

Tribe I *Gloeogenae*

b Cells free or in twos or fours

c Cells spherical

Chroococcus Naeg

cc Cells cylindrical

Synechococcus Naeg

bb Cells in resting stage united into amorphous slimy families

c Cell membranes continuous with intercellular substance

d Cells without phycochrome, very small

e Cells spherical

Micrococcus Hall emended

ee Cells cylindrical

Bacterium Duj

dd Cells with phycochrome, larger

e Cells spherical

Aphanocapsa Naeg

ee Cells cylindrical

Aphanothece Naeg

cc Intercellular substance formed of layers of cell membranes

d Cells spherical

Gloeocapsa Kg Naeg

dd Cells cylindrical

Gloeothece Naeg

bbb Cells united into definite slimy families

c Cell families of a flat single layer of cells

d Cells in fours in one plane

Merismopedia Meyen

dd Cells irregularly grouped on surface of sphere

e Cells spherical, families broken into nets

Clathrocystis Henfr

ee Cells cylindric

Coelosphaerium Naeg

cc Cell families, many layered, united into spherical cell bodies

d. Number of cells definite

e Cells spherical, in tetrads, colorless

Sarcina Goods

ee Cells cylindric, not regularly arranged, with phycochrome

Gomphosphaeria Kg

dd Number of cells indefinite, very large

e Cells colorless, very small

Ascococcus Billr emended

ee Cells with phycochrome, larger

Polycystis Kg*Coccochloris* Spr*Polycoccus* Kgaa Cells united into filaments Tribe II *Nematogenae* Rab

b Cell filaments always unbranched

c Filaments free or felted together

d Filaments cylindric, colorless, indefinitely jointed

e Filaments very slender, short

Bacillus Cohn

ee Filaments very slender, long

Leptothrix Kg emeee Filaments thicker, long *Beggiatoa* Trev

- dd Filaments cylindric, with phycochrome, definitely jointed.
Reproductive cells unknown
Hypheothrix Kg
Oscillaria Bosc u a
- 3d Filaments cylindric, jointed, forming gonidia
e Colorless *Crenothrix* Cohn
ee With phycochrome *Chamaesiphon* u a
- 4d Filaments spiral
e Without phycochrome
f Filaments short, slightly wavy
Vibrio Ehr em
ff Filaments short, spiral, stiff
Spirillum Ehr
fff Filaments long, spiral, flexuous
Spirochaete Ehr
Spirulina Link
- ee With phycochrome
- 5d Filaments rosary like
e Colorless *Streptococcus* Billr
ee With phycochrome *Anabaena* Bory
- 6d Filaments tapered to tip *Spermosira* Kg u a
- cc Filaments united together into slimy families
d Filaments cylindric, colorless *Myconostoc* Cohn
dd Filaments cylindric, with phycochrome
Chthonoblastus-Limnochlide
3d Filaments rosary like *Nostoc, Hormosiphon* u a
4d Filaments tapering to tip *Rwularia* Roth
- bb All filaments with false branches
c Filaments cylindric, colorless *Cladothrix* Cohn
Streptothrix Cohn
- cc Filaments cylindric, with phycochrome
Calothrix Ag
3c Filaments rosary like *Merizomyria* Kg
4c Filaments tapering *Schizosiphon* Kg

It will be noted that the following bacterial genera were new, *Ascococcus* (species *A. Billrothii*), *Streptococcus* (erroneously ascribed to Billroth, even here without a species designation and probably therefore invalid) *Myconostoc* (*M. gregarium*), *Cladothrix* (sp *C. dichotoma*, name invalid because earlier used for another genus of plants), *Streptothrix* (*Str. Foersteri*, generic name invalid because of previous use of this generic name for a genus of fungi)

It has been suggested that if a date for a point of departure for the classification of bacteria is to be fixed it might well date from this 1875 classification of Cohn

In 1877 the veterinarian Bollinger sent to the botanist Harz purulent material from a case of "lumpy jaw" in a cow Harz found present

an organism which he named *Actinomyces bovis*, the type of a group of organisms now recognized as showing many intermediate characters between the true bacteria and the fungi. Rivolta (1878, p. 169) objected to this name, and proposed in its place *Discomyces bovis*.

McNab (1877, p. 340) outlined a classification of the "Vegetable Kingdom." His lowest group was the Subkingdom *Thallophyta*, with several classes. The first of these he termed *Schizophyta* or *Protophyta*. The orders and families were as follows:

McNab's Orders of the Schizophyta (1877)

- Order I *Cyanophyceae* or *Phycochromaceae*
Families *Chroococcaceae*, *Nostocaceae*, *Oscillatoriaceae*, *Rivulariaceae*, *Scytonemaceae*
- Order II *Chlorophyllophyceae*
Families *Palmellaceae*, *Euglenaceae*
- Order III *Schizomycetes*
Family *Bacteriaceae*
- Order IV *Saccharomycetes*
Family *Saccharomycaceae*

Van Tieghem (1878, p. 198) introduced the generic name *Leuconostoc* to designate a streptococcus-like form occurring in large gelatinous masses in syrup, an organism previously described by Cienkowski (1878, p. 12) as *Ascococcus mesenteroides*.

Magnin (1878) in his text "Les Bacteries" reviewed at length the classification of bacteria, and adopted the scheme of Cohn.

Miquel (1879-1881) published a series of articles on the organisms which produce ammoniacal fermentation in urine. He introduced the generic names *Urobacillus*, *Urococcus* and *Urosarcina* for bacilli, cocci and sarcinae which could bring about this change.

Trevisan (1879, p. 13) created several new genera of bacteria, and evolved the following classification:

Trevisan's Classification of Bacterial Genera (1879)

- a Unicellular bacteria, globose or cylindric. Spores solitary within the cells.
Multiplication by division of the cells in one, two, or all directions.
Tribe I *Bacterieae*
- b Cells free or when quiescent associated in indeterminate mucous colonies. Subtribe I *Eubacterieae*
- c Cells globose or globose ovoid. Multiplication by vegetative division of the cells in all directions.
Genus 1 *Micrococcus*

Winter's Classification of Bacteria (1880)

- a. Cells spherical or oval
 - b Cells isolated, in chains or embedded irregularly in mucus
 - 1 *Micrococcus* Cohn
 - bb Cells united into definite colonies
 - c Colony hollow, with peripheral layer of cells
 - 2 *Cohnia* Winter
 - cc Colony solid, filled with cells
 - d Cells united in small but definite numbers into regular families
 - 3 *Sarcina* Goodsir
 - dd Cells in large and indefinite numbers united into irregular colonies
 - 4 *Ascococcus* Cohn
 - aa Cells cylindric, short or long
 - b Cells short, cylindric, single or bound loosely together in pairs
 - 5 *Bacterium*
 - bb Cells longer, cylindric, united into filaments
 - c Filaments isolated or felted
 - d Unbranched
 - e Cells straight
 - f Short, definitely jointed
 - 6 *Bacillus* Cohn
 - ff Long, indefinitely jointed
 - g Very slender
 - 7 *Leptothrix* Kutzing
 - gg Thicker
 - 8 *Beggiatoa* Trevisan
 - ee Cells spiral or bent
 - f Cells short, with few turns, not flexuous
 - 9 *Spirillum* Ehr
 - ff Cells longer, with numerous turns, flexible
 - 10 *Spirochaeta* Ehr
 - dd With apparent branches
 - 11 *Cladothrix* Cohn
 - cc Cells embedded in rounded gelatinous masses
 - 12 *Myconostoc* Cohn
 - aaa. Cells lancet-like, flat bands, spirally wound
 - 13 *Spiromonas* Perty

In an "Anhang" he includes two other genera, *Sphaerotilus* Kutzing and *Crenothrix* Cohn. The name *Cohnia* used as generic designation by Winter was preoccupied, and changed later by Schroeter to *Lamprocystis*.

Prazmowski (1880) in a study of certain anaerobic bacteria revived Trecul's casual name *Clostridium*, and created a genus with this designation to include those organisms which produce spores in more or less spindle-shaped cells.

De Lanessan (1880) gave an outline of a classification of the genera of bacteria as follows

de Lanessan's (1880) Classification of Bacteria

- A Cells not united in filaments, separating immediately after division, or in colonies (zoogloea) united by a gelatinous substance
 - I Cells dividing in one direction only
 - a Cells spherical *Micrococcus*
 - b Cells elliptical or short cylindrical *Bacterium*
 - II Cells dividing regularly in three directions of space, forming cubical colonies *Sarcina*
- B Cells cylindrical in filaments
 - I Filaments straight, imperfectly segmented
 - a Filaments very slender and short, small rods *Bacillus*
 - b Filaments slender and long *Leptothrix*
 - c Filaments broader and long *Beggiatoa*
 - II Filaments undulant or spiral
 - a Filaments short
 - 1 Filaments slightly undulant *Vibrio*
 - 2 Filaments spiral, stiff *Spirillum*
 - b. Filaments long, flexible, with rapid undulations, spiral throughout length, slender, actively motile *Spirochaeta*

Bergonzini (1881) named a violet organism which he studied *Cromobacterium violaceum*

Kern (1882, p 135) studied the organism producing a Caucasian fermented milk beverage. Because it apparently produced two spores to a cell he named it *Dispora caucasica*. Either Kern mistook other cell inclusions for spores or confused spore-bearing bacteria with the sour milk form. His organism cannot be satisfactorily identified from his description, and the genus name has not been used by other writers.

Zopf (1883) in the first edition of his "Die Spaltpilze" formulated a somewhat new characterization of bacterial genera. The families and genera recognized may be separated as follows

Zopf's Classification of Bacteria (1883)

- a Cells spherical, sometimes in filaments made up of coccoid elements. One genus only *Leuconostoc*
- aa Organisms showing coccus and rod shapes in growth stages
 - b Spiral forms lacking as a growth stage. Filamentous stage showing no differentiation of base and tip
 - c Not becoming spindle or club shaped when sporulating *Bacterium*
 - cc Rods becoming spindle shaped or ellipsoidal or club shaped when sporulating *Clostridium*

bb Possessing spiral stage in the growth

c Not forming false branches

Leptothrix, *Beggiatoa*, *Crenothrix* and *Phragmidiothrix*

cc Forming false branches

Cladothrix

Balcke (1884) according to Lindner named an organism from fermenting wort *Pediococcus cerevisiae*. This generic name has been commonly used since as a designation for organisms which occur in tetrads but do not form regular *Sarcina* packets.

De Bary (1884) introduced a new principle into the classification of bacteria by his separation of these organisms into two groups based upon the presence of endospores in the one and of arthrospores in the other. The following key will illustrate this method of differentiation.

De Bary's Classification of Bacteria (1884)

- a Producing endospores
 - b Rods straight or slightly bent (including all non-arthrosporous types of *Clostridium*, *Bacteridium*, *Vibrio*, etc.)
 - 1 *Bacillus*
 - 2 *Spirillum*
 - bb Rods spiral
- aa Producing arthrospores
 - b Rosaries of cocci, embedded in firm masses of gelatin
 - 3 *Leuconostoc*
 - bb Simple rods
 - 4 *Arthrobacterium*
 - bbb Filamentous forms
 - 5 *Crenothrix*
 - 6 *Beggiatoa*
 - 7 *Cladothrix*
 - 8 *Leptothrix*

The classification used by Grove (1884) was an exact translation of that of Winter.

Van Tieghem (1884) agreed with Cohn in including the bacterial genera among the blue-green algae. The following key, carried to genera only for the bacteria, will illustrate the rather indefinite grouping used by this author.

Van Tieghem's Classification of Bacteria (1884)

- A Family I *Nostocaceae* Producing cysts. With many algal genera and the bacterial genera *Beggiatoa* and *Leuconostoc*
- B Family II *Bacteriaceae* Producing spores
 - I Cell division in one direction of space
 - a Cells spherical *Micrococcus*
(In gelatin, *Ascococcus*, Naked, *Punctula*)

- b Cells elongate
 - 1 Cells not spiral
 - (a) Cells separating when formed
Bacterium
 - (b) Cells forming a chain or filament
Bacillus
 - (c) Filaments relatively long
 - (1) Sheathless *Leptothrix*
 - (2) Sheathed *Crenothrix*
 - (d) Filaments branched *Cladothrix*
 - (e) Enclosed in gelatin *Ascobacteria*
 - (f) Naked cells *Polybacteria*
 - 2 Cells spiral
 - (a) Cells separating quickly *Vibrio*
 - (b) Spiral, like a spring *Spirillum*
 - (c) Cells longer and more slender
Spirochaete
 - (d) Cells embedded in gelatin
Myconostoc
- II Cell division in two directions of space, giving rise to tetrads
Merista
- III Cell division in three directions of space forming cubical packets
Sarcina

The genera *Streptococcus* for the chain cocci and *Staphylococcus* for the irregularly grouped cocci from pus were first adequately defined by Rosenbach (1884). Both of these names had previously been used by other writers, but Rosenbach was apparently the first to associate them definitely with species.

Zopf (1884) in the third edition of *Die Spaltpilze* materially modified and enlarged the bacterial classification used in the first and second editions. The four main groups were recognized as before, but the number of genera increased. Following is the classification used.

Zopf's Classification of Bacteria (1885)

- a Cells producing only a spherical stage
 - b Not embedded in firm masses of gelatin
 - c Cells regularly arranged
 - d Cells in chains *Streptococcus*
 - dd Cells not in chains
 - e Cell division in two planes, forming flat plates of cells *Merismopedra*
 - ee Cell division in three planes, forming regular cubes and packets *Sarcina*
 - cc Cells irregularly arranged *Micrococcus*

- bb Cells embedded in thick gelatinous masses
Ascococcus
- aa Consisting of cocci, rods and filaments, or the latter two forms only, in filaments no differentiation of base and tip
 - b Spore formation unknown or not occurring
 - c Cells not spiral *Bacterium*
 - cc Cells spiral *Spirillum*
 - bb Spore formation known
 - c Filaments spiral *Vibrio*
 - cc Filaments not spiral
 - d Forming cocci and rods, spore formation in the former *Leuconostoc*
 - dd Forming cocci and rods or latter only Spores in rods or cocci *Bacillus*
 - ddd As in dd, but spores in swollen cells *Clostridium*
- 3a Consisting of coccus, rod and filament forms, the latter with a differentiation of base and tip
 - b Filaments containing sulphur, sheathless *Beggiatoa*
 - bb Filaments without sulphur
 - c Filaments sheathed *Crenothrix*
 - cc Filaments not sheathed, cell divisions ultimately both longitudinal and transverse *Phragmidiothrix*
 - ccc Filaments with or without sheaths, no longitudinal cell division *Leptothrix*
- 4a Consisting of cocci, rods, filaments and spirals False branching occurs
Spore formation is unknown *Cladothrix*

Trevisan (1885) described several new genera of bacteria, including *Kurthia*, *Zopfella*, *Bacteriopsis*, *Klebsiella* and *Gaffkya*

In the same year he also created the genus *Pacima* to include organisms of the type of the Asiatic cholera organism

Hauser (1885) described a new genus *Proteus* to include certain putrefactive bacteria which formed characteristic amoeboid colonies on gelatin and which exhibited a high degree of pleomorphism, varying from cocci to rods and spirals

Lutz (1886) proposed the generic name *Coccothrix* to designate the acid fast bacteria of the types of the organisms of tuberculosis and of leprosy. He also made this genus the type of the family *Coccothrichaceae*. In the same year Weichselbaum used the designation *Diplococcus pneumoniae* for the causal organism of pneumonia

Flügge (1886) outlined a classification of bacterial genera which has been extensively followed by medical bacteriologists. A key to his genera may be constructed as follows

GENERAL SYSTEMATIC BACTERIOLOGY

Flügge's Classification of Bacteria (1886)

- a Cells spherical
 - b. Cells isolated, or bound in chains united into amorphous slimy families
 - 1 *Micrococcus*
 - bb. Cells united into definite slimy families
 - c Colonies solid (filled with cells)
 - d Cells united in large and indefinite numbers into irregular colonies
 - 2 *Ascococcus*
 - dd Cells in small but definite numbers united into regular families
 - 3 *Sarcina*
 - cc Colonies with peripheral layers of cells, no cells in the interior
 - 4 *Clathrocystis*
- aa Cells cylindric
 - b Cells short cylindric, single or loosely hanging together, or united into amorphous slimy masses
 - 5 *Bacterium*
 - bb Cells longer cylindric, bound into filaments
 - c Filaments isolated, in filaments or in bundles
 - d Filaments unbranched
 - e Filaments straight
 - f Filaments short, definitely jointed
 - 6 *Bacillus*
 - ff Filaments long, indefinitely jointed
 - g Filaments very slender
 - 7 *Leptothrix*
 - gg Filaments thicker
 - 8 *Beggiatoa*
 - ee Filaments undulating or spiral
 - f Short, stiff
 - 9 *Spirillum*
(*Vibrio*)
 - ff Long, flexible
 - 10 *Spirochaeta*
 - dd Filaments with false branches
 - 11 *Streptothrix*
 - 12 *Cladothrix*
 - cc Filaments enclosed in round gelatinous masses
 - 13 *Myconostoc*

Genera doubtfully belonging here *Crenothrix*, *Sphaerotilus*, *Spiromonas*, *Rhabdomonas*, *Monas*

Schroeter (1886) outlined the most complete and elaborate classification of bacterial genera and species that had been prepared up to that time. He introduced a number of new genera. His generic key is as follows:

Schroeter's Classification of Bacteria (1886)

- I Order *Coccobacteria* Cells in all stages of development, spherical-elliptical, never motile. Spores, if present, developing through transformation of an entire cell. Family *Coccaceae* Zopf

A Cells or colonies free or embedded in slime, without definite gelatinous membranes

1. Cells single or arranged irregularly *Micrococcus* Cohn

2 Cells united in chains *Streptococcus* Billroth

3 Cells united in a stratum

a In tabular form *Lamprospira* n g

b In hollow balls *Lamprocystis* n g

B Single cells or cell colonies with a definite firm gelatinous covering (cyst)

1 Cells single or irregularly grouped

a Each cell or small cell family included in a simple gelatinous cyst *Hyalococcus* n g

b Cells or few celled families included in a many layered slime capsule *Leucocystis* Schroeter

c Cell families of many cells enclosed in an almost solid gelatinous mass *Ascococcus* Billroth

2 Cell families regularly formed

a Cell families in packet-shaped bundles, each single cell with a capsule *Sarcina* Goodsir

b Cells in chains, each family in a capsule *Leuconostoc* Van Tieghem

II Order *Eubacteria* Cells rod shaped, shorter or longer

Family *Bacteriaceae*

A Cells or cell groups without definite gelatinous capsule

1 Cells straight or slightly bent, not spiral

a Cells very small, elliptical, motile or non-motile, spores unknown *Bacterium* Ehr

b Cells cylindrical, motile or non-motile

(1) Cells relatively very broad Red, with sulfur

Chromatium Perty

(2) Cells in vegetative condition narrow rods Endogenous spores

(a) Spore bearing cells cylindric

Bacillus Cohn

(b) Spore bearing cells spindle shaped

Clostridium Praz

2 Cells and cell chains spiral, usually actively motile

a Spirals stiff (not flexuous) Endogenous spores

Spirillum Ehr

b Spirals flexuous

(1) Cells known only in the form of a long flexuous spiral

Spores unknown *Spirochaete* Ehr

(2) Cells in greater part of life only with short ($\frac{1}{2}$) spiral, later growing to screw form Spores spherical

Arthrospores *Microspira* n g

B. Cells enclosed in definite gelatinous masses

1 Cells wound in small numbers in a transparent capsule

Myconostoc Cohn

- 2 Cells rod shaped, in large numbers in a brown thick capsule
Cystobacter n. g.

III Order. Desmobacteria Long filaments, usually surrounded by a definite sheath

A Filaments unbranched Family *Leptotrichaceae* Zopf

1 Filaments always non-motile

- a Filaments with very thin sheaths Multiplication by falling apart into two segments *Leptothrix* Kutzing

- b With thick sheaths Multiplication by spores and separation of single joints *Crenothrix* Cohn

2 Filaments motile

Beggiatoa Trevisan

B Filaments branched

1 Filaments not club shaped or thickened at end

- a Filaments free *Cladothrix* Cohn

- b Filaments bound in bundle by slimy capsule
Sphaerotilus Kutzing

2 Filaments club thickened at end

Actinomyces Harz

The new genera described were *Lampropedia*, *Lamprocystis*, *Hyalococcus*, *Leucocystis*, *Microspira* and *Cystobacter*

Trevisan (1887) outlined the following classification of bacterial genera

Trevisan's Classification of Bacteria (1887)

1 *Bacteriaceae trichogenae* Three stages of vegetative development

1 Filaments (typical) 2 Rods 3 Cocci

A Filaments fixed, with base distinct from tip

(*Leptotrichineae*)

Genus 1 *Cladothrix* Cohn

Genus 2 *Crenothrix* Cohn

Genus 3 *Sphaerotilus* Kutzing

Genus 4 *Leptotrichia* Trevis (1879)

Genus 5 *Beggiatoa* Trevis (1842)

Genus 6 *Phragmidiothrix* Engler (1881)

B Filaments free, no distinction between base and tip

1 Filaments spiral, not embedded in a mucous capsule (*Spirilleae*)

Genus 7 *Kurthia* Trevis (1885)

Genus 8 *Vibrio* Cohn (1872)

Genus 9 *Spirillum* Ehrenb

Genus 10 *Spirochaete* Ehrenb

2 Filaments spiral, embedded in mucous capsule (*Myconostocaceae*)

Genus 11 *Myconostoc* Cohn

3 Filaments straight (*Zopfelleae*)

Genus 12 *Zopfella* Trevisan (1885)

II *Bacteriaceae baculogenae* Three stages of vegetative development

1 Rods (typical) 2 Filaments (abnormal or transitory stage) 3 Cocci

- A Not capsulated (*Bacilleae*)
 - Genus 13 *Bacillus* Cohn
 - Genus 14 *Pacima* Trevis (1885)
 - Genus 15 *Clostridium* Prazmowski
 - Genus 16 *Pasteurella* Trevis
 - Genus 17 *Bacterium* Cohn (1872)
 - Genus 18 *Bacteriopsis* Trevis (1885)
- B Normally capsulated (*Klebsielleae*)
 - Genus 19 *Klebsiella* Trevis (1885)
- III *Bacteriaceae coccigenae* One stage only of development, coccus
 - A Cocci enclosed in common membranous mucous capsule (*Ascococceae*)
 - Genus 20 *Leuconostoc* Van Tieghem
 - Genus 21 *Ascococcus* Cohn
 - Genus 22 *Gaffkeya* Trevis (1885)
 - B Cocci in packets, chains, or isolated No capsules
 - Genus 23 *Sarcina* Goodsir
 - Genus 24 *Staphylococcus* Ogston
 - Genus 25 *Streptococcus* Zopf
 - Genus 26 *Micrococcus* Cohn (emend)

Maggi (1887) in his "Acque potabili" outlined a bacterial classification as follows

Maggi's Classification of Bacteria (1887)

- A Order I *Spirobacteria* Cohn
 - I Genus 1 *Spirochaeta* Ehr
 - II Genus 2 *Spirillum* Ehr
 - III Genus 3 *Vibrio* Auct emend
- B Order II *Desmobacteria* Cohn
 - I Genus 1 *Cladothrix* Cohn
 - II Genus 2 *Crenothrix* Cohn
 - III Genus 3 *Beggiatoa* Trevisan
 - IV Genus 4 *Leptothrix* Kutzing
 - V Genus 5 *Bacillus* Cohn
- C Order III *Microbacteria* Cohn
 - I Genus 1 *Bacterium* Dujardin emend
 - II Genus 2 *Zygobacterium* Maggi
- D Order IV *Sphaerobacteria* Cohn
 - I Genus 1 *Micrococcus*

The new genus *Zygobacterium* with its single species *Z. nitrosum* does not seem to be adequately described. The figures show slender rods lying parallel to each other in pairs.

Metschnikoff (1888) described as *Pasteuria ramosa* an organism which he concluded was a bacterium which divided longitudinally rather than transversely. Although this generic name has been used by subsequent writers as a bacterial genus, it is probable that it is best regarded as protozoan.

Winogradsky (1888) reported the results of a study on the morphology and physiology of the sulphur bacteria. A number of new genera were created, and several old genera brought definitely into this group. To the present time the account given by this author is the only recorded observation of certain of these forms. His key for the separation of these sulphur genera may be translated as follows

Key to Winogradsky's Genera of Sulphur Bacteria (1888)

A Colorless sulphur bacteria

I Filaments uniform in thickness, always free, motile, forming no gonidia *Beggiatoa*

II Filaments of variable thickness, attached forming motile rod gonidia. *Thiothrix*

B Red sulphur bacteria

I Cells united into families

a Division of cells in three planes

1 Families of cells small, solitary or many surrounded by a gelatinous cyst, swarming *Thiocystis*

2 Families of cells spread out on a substrate, spherical cells embedded in a common gelatin, not swarming *Thiocapsa*

3 Families packet shaped *Thiosarcina*

b Division of cells first in three planes, then in two Families first solid, then hollow *Lamprocystis*

c Division of cells in two directions of space Families tabular, tetrads, swarming *Thiopedia*

d Division of cells in one plane

1 Families amoeboid motile Cells united by plasma threads *Amoebobacter*

2 Families with thick gelatinous cysts Cells in common gelatin, very loosely embedded, swarming *Thiothece*

3 Families of rods united into a net *Thiodictyon*

4 Families solid, non-motile, of small cells closely appressed *Thioplycoccus*

II Cells free, always swarming

a Cells cylindric-elliptic *Chromatium*

b Cells rod and spindle shaped *Rhabdochromatium*

c Cells spiral *Thiospirillum*

Hansgurg (1888, p 229) called attention anew to the parallelism existing in morphology in the bacteria and the algae. He gives as an illustrative table the following

Hansgirk's Comparison of Bacteria and Blue-green Algae (1888)

Bacteria	Blue Green Algae
<i>Micrococcus-Hyalococcus</i>	<i>Aphanocapsa-Chroococcus</i>
<i>Bacterium-Streptococcus</i>	<i>Aphanothece-Synechococcus</i>
<i>Leucocystis-Mycethece</i>	<i>Gloeocapsa-Gloeothece</i>
<i>Ascococcus-Leuconostoc</i>	<i>Polycoccus-Anacystis (Polycystis)</i>
<i>Lamprocystis (Cohnia)-Sarcina</i>	<i>Clathrocystis-Merismopedium</i>
(<i>Lampropedia</i>)	<i>Iyngbya exp (Leptothrix exp Hypheothrix)</i>
<i>Bacillus-Leptothrix exp</i>	<i>Oscillaria-Spirulina</i>
<i>Beggiatoa-Spirochaete (Spirillum)</i>	<i>Nostoc</i>
<i>Myconostoc</i>	<i>Chamaesiphon</i>
<i>Crenothrix</i>	<i>Plectonema (Glaucothrix)</i>
<i>Cladothrix (Streptothrix)</i>	

He also prepared an outline of the groups of bacteria with particular reference to the forms to be found in damp cellars and other subterranean habitats. While adequate descriptions are not given by Hansgirk, the following key is believed to represent the principal distinctions among the various groups

Hansgirk's Classification of Bacteria (Schizomycetes) (1888)

- A. Filamentous
 - Order I *Desmobacteria*
 - Family *Cladothricaceae*
 - One genus only Genus *Cladothrix* Cohn
 - II No false branching
 - a Gonidia formed
 - Family *Crenothricaceae* Hansg
 - One genus only Genus *Crenothrix* Cohn
 - b No gonidia
 - Family *Leptothricaceae*
 - 1 Filaments immobile
 - Subfamily *Leptothriceae*
 - One genus only Genus *Leptothrix* Ktz ex p
 - 2 Filaments oscillating
 - Subfamily *Beggiatoeae* Hansg
 - One genus only Genus *Beggiatoa* Trevisan
 - (a) Filaments not colored
 - Sec 1 *Eubeggiatoa* Hansg
 - (b) Filaments colored purple
 - Sec 2 *Chromobeggiatoa* Hansg
- B. Cells elongate, but not filamentous
 - Order II *Eubacteria*
 - Family *Bacteriaceae*
 - I Cells not encysted
 - a Cells spirally bent
 - Subfamily *Spirobacteria* Cohn
 - 1 Cells flexuous Genus *Spirochaete* Ehrb
 - 2 Cells not flexuous Genus *Spirillum* Ehrb
 - (a) Colorless, without sulphur
 - Section 1 *Vibrio* (Cohn) Hansg
 - (b) Rose colored, with sulphur

- Section 2 *Ophidomonas* (Ehr) Hansg
- b Cells not spirally bent
- Subfamily *Microbacterina* Cohn
- 1 Cells elongate Genus *Bacillus* Cohn
- (a) Without sulphur inclusions
- Section 1 *Eubacillus* Hansg
- (b) With sulphur inclusions
- Section 2 *Chromobacillus* Hansg
- 2 Cells short Genus *Bacterium* Ehrb
- II Cells encysted Family *Myconostocaceae* Hansg
- One genus only Genus *Myconostoc* Cohn
- C Cells spherical Order III *Sphaerobacteria*
- One family only Family *Mycococcaceae*
- a Cells encysted Subfamily *Cystococcaceae* Hansg
- 1 Cells irregularly grouped
- a Each cell or each group of a small number of cells in common cyst Genus *Hyalococcus* Schröter
- b As (a), but with many-layered cysts Genus *Leucocystis*. Schroter
- c Cell families of very numerous individuals in a common cyst Genus *Ascococcus* Billroth
- 2 Cells regularly grouped
- a Cells in chains, encapsulated Genus *Leuconostoc* Van Tieghem
- b Cells in twos and fours, in many layered capsules Cells elongate Genus *Mycotheca* Hansg

Metschnikoff (1888) suggested the generic name *Sclerothrix* for the acid fast bacteria of the type of the bacillus of tuberculosis. Evidently he was not aware of the previous use of this name by Kuetzing (1849) for a genus of algae.

Bollinger (1888) suggested the name *Botryomyces* as a designation for the genus of cocci found in the so-called botryomycomata of the horse. In this year Gamaleia used the name *Coccobacillus avicidus* for the organism of chicken cholera.

Bejerinck (1889) proposed the generic designation *Photobacterium* to include all of the photogenic forms. This is perhaps the first genus of bacteria proposed based upon a physiologic character. Frank (1889) used *Rhizobium* as the generic name for the bacteria found in the root nodules of leguminous plants. *Spirobacillus* was named by Metschnikoff in the same year.

Probably the most elaborate classification of bacteria based entirely upon morphologic characters is that of De Toni and Trevisan (1889) published in Saccardo's *Sylloge Fungorum*. Many new genera were

recognized, and specific descriptions given of over 650 species. Many of the species and even some of the genera cannot at present be identified with certainty, but this classification has nevertheless been entirely too much ignored by bacteriologists.

De Toni and Trevisan's Classification of Bacteria (1889)

Key to Subfamilies of Schizomycetaceae Naeg

- 1 Three stages of development, filaments, rods, and cocci. Filaments (in the primary stage) sheathed or sheathless, usually with base and apex differentiated, attached or radiating from a central point, rarely free. Rods and cocci included in the filaments. I *Trichogenae* Trev
- 2 Three stages of development, individual rods, filaments and cocci. Rods (in the primary stage) forming filaments and cocci, filaments (2'' stage) never sheathed, nor with distinct base, nor fixed, nor radiating from a central point, formed by single bacilli remaining attached, due to imperfect division. II *Baculogenae* Trev
- 3 One stage only, cells spherical. III *Coccigenae* Trev

Key to Tribes and Genera of the Trichogenae Trev

- a Spores (arthrospores) produced in special parts of the filaments (pseudosporangia). Tribe I *Crenotricheae* Trev
Filaments sheathed unbranched. *Crenothrix* Cohn
- aa Spores (arthrospores) produced in normal filaments. Tribe II *Cladotricheae* Trev
 - b Filaments pseudoramose
 - c Filaments sheathed
 - d Filaments subequal from base to apex. Arthrospores numerous, dividing in three directions of space. *Sphaerotilus* Kuetz
 - dd Filaments from base to apex more and more thickened. Arthrospores formed, dividing only transversely. *Cladotrix* Cohn
 - cc Filaments not sheathed. Arthrospores coccus-shaped, division transverse only. *Nocardia* Trev
- bb Filaments unbranched. Tribe III *Kurthiae* Trev
 - c Arthrospores rod shaped. *Detonella* Trev
 - cc Arthrospores cocci
 - d Filaments fixed at base in zoogloeal masses. *Rasmussenia* Trev
 - dd Filaments free. *Kurthia* Trev
- aaa. Spores absent, at least never detected. Filaments unbranched. Tribe IV *Leptotricheae* Trev
 - b Filaments sheathed. Base and tip unequal. Rod-shaped gonidia. *Leptotrichia* Trev
 - bb Filaments, not sheathed, equal, no rod gonidia

- | | | | |
|----|--|---|-------------------------------|
| c | Filaments fixed | Cell division transverse and longitudinal | <i>Phragmidiothrix</i> Engler |
| cc | Filaments free, division transverse only | | <i>Beggiatoa</i> Trev |

Key to Tribes and Genera of the Subfamily Baculogenae Trev

- a Bacilli and cocci naked, never with a tegument
 Tribe I *Bacillae* Trev
- b Producing endospores
 c Division longitudinal Subtribe I *Pasteureae* Trev.
 One genus only *Pasteuria* Metch
 cc Divisions transverse
 d Rods united in a reticulate coenobium
 Subtribe II *Thiodictyeae*
 One genus only *Thiodictyon* Winogradsky
 dd Rods not united in reticulate coenobium
 e Rods straight or slightly bent, never spiral
 Subtribe III *Eubacillae* Trev
 f Spores not greater in diameter than normal vegetative rods
 g Spores forming in unchanged rods
 h Plasma uniformly diffused in rods
 i Rods fusiform
Mantagozzaea Trev
 ii Rods cylindrical or cylindraceous
Bacillus Cohn
 hh Plasma bipolar in distribution
Pasteurella Trev
 gg Spores with special integuments, cells becoming ellipsoid-fusiform
Clostridium Praz
 ff Spores greater in diameter than the normal vegetative rods
 g Spores median in unchanged rods
Cornuha Trev
 gg Spores apical in clavate rods
Vibrio Zopf
- ee Rods spirally bent
 Subtribe IV *Spirilleae* Trev
 f Rods circular in cross section
Spirillum Ehr
 ff Rods compressed in cross section
Spiromonas Perty
- bb Producing arthrospores
 c Cylindrical, straight or slightly curved Filaments undulate—flexuous or irregularly pseudospiral
Pacima Trev

- cc Rods ellipsoidal, straight, filaments never pseudospiral
Bacterium Ehr
- aa Bacilli and cocci surrounded by a special membranous-gelatinous capsule
Tribe *Klebsiellae* Trev
 - b Bacilli straight or curved, never true spirals
Subtribe *Euklebsiellae* Trev
 - c Capsule repeatedly ramose *Winogradskya* Trev
 - cc Capsule simple, never ramose
 - d Plasma uniformly diffused *Klebsiella* Trev
 - dd Plasma condensed at poles *Dicoccia* Trev
 - bb Rods spirally twisted Subtribe *Myconostocae* Trev
One genus only *Myconostoc* Cohn
Anhang Cystobacter

Key to Tribes and Genera of Subfamily Coccogenae

- a Cocci associated in families in bladder-like gelatinous masses or cysts
Tribe *Ascococceae* Trev
- b Cocci separated in mucus of the matrix
 - c Cocci destitute of special cysts, in families in common cysts
Subtribe *Euscocceae*
 - d Cocci numerous, associated in large families
 - e Cysts homogeneous, not lamellose
 - f Families at first solid, then irregularly clathrate
Division at first in three directions, then in two directions *Lampiocystis* Schroet.
 - ff Families in solid mass, cocci dividing in one direction
Ascococcus Cohn
 - ee Cysts lamellose Solid colonies Cocci divide in three planes *Bollingeria* Trev
 - dd Cocci not numerous, associated in small families
 - e Cysts many layered Division in three directions
Leucocystis Schroet
 - ee Cysts homogeneous, not lamellose
 - f Cocci all in periphery Cocci divide at first in three directions in two
Cenomesia Trev
 - ff Cocci few and scattered Divide in one direction
Thiothece Winogr
 - fff Cocci in solid families, cocci divide in three directions
Thiocystis Winogr
 - cc Cocci borne in special cysts Common cysts absent
Subtribe *Gaffkyae* Trev
 - d Cysts broad, persistent, numerous in conglobate heap, interior solid, densely aggregated
Chlamydatomus Trev
 - dd Cysts tenuous, at length breaking apart, solitary, never associated in heaps *Gaffkyia* Trev

- bb Cocci in mucoid matrix loosely joined in a plasmic filamentous series.
Cysts very tenuous, at length breaking apart Special cysts none.
Subtribe *Amoebobacterinae* Trev
One genus only *Amoebobacter* Winogr
- aa. Cocci in families of single or many layers, joined by mucus more or less conspicuous No cysts, endospores small, appearing in cocci
Tribe *Sarcinae* Trev
 - b Cocci heaped up closely in firm mucous matrix
 - c Cocci in solid irregular heaped families without order, densely associated Cocci divide in one direction
Thiopolyccus Winogr
 - cc Cocci closely associated in regular cubical families, dividing in three directions
Sarcina Goodsir
 - bb Cocci grouped loosely in mucous matrix
 - c Cocci in fours in one layered families, in firm mucus, forming parallelograms, loosely associated Division in two planes
Lampropedia Schroet
 - cc Cocci few in irregular families in firm mucoid membranes, periphery amorphous, associated without regular order Divide in three directions
Thiocapsa Winogr
 - ccc Cocci in fours in small regular colonies, one layered, tenuous, amorphous, scarcely breaking apart, loosely associated Divide in two planes
Pediococcus Lindner
- aaa Cocci in moniliform chains Arthrospores large in the filaments or on apex
Tribe III *Streptococceae* Trev
 - b Filaments produced in a membranous, gelatinous capsule
 - c Capsules large, broad, lamellose *Leuconostoc* Van Tiegh
 - cc Capsules close, tenuous, not lamellose
Schuetzia Trev
 - bb Filaments in a cylindrical sheath
Peironicota Trev
 - bbb Filaments naked
 - c Pseudodichotomous Arthrospores at tip of filaments
Babesia Trev
 - cc Simple Arthrospores included in filaments
Streptococcus Billroth
emended Zopf
- 4a Cocci not in cysts, capsules or sheaths, nor in chains, with endospores
Tribe IV *Micrococceae* Trev
 - b Cocci in pairs
Neisseria Trev
 - bb Cocci associated in botryoid masses
Staphylococcus Ogston
 - bbb Cocci solitary or associated in amorphous zoogloeal masses
Micrococcus (Hallier) Cohn

Hansgurg (1890) described two new bacterial genera which have been uniformly ignored by subsequent writers They are *Mycacanthococcus* and *Mycotetraedron*, both from wine cellars

Baumgarten (1890) ignored the recognized rules of nomenclature in the following classification

Baumgarten's Classification of Bacteria (1890)

- A Species relatively monomorphic
- I Cells spherical
 - a Cells in pairs Genus *Kokkus*
 - b Cells in chains Subgenus *Diplokokkus*
 - c Cells in tabular masses Subgenus *Streptokokkus*
 - d Cells in regular packets Subgenus *Merista* or *Merismopedia*
 - e Cells in irregular masses Subgenus *Sarcina*
 - II Cells rod shaped Many species with spores
 - Genus *Bacillus*
 - III Cells spiral
 - Genus *Spirillum*
 - Subgenera
 - Subgenus *Vibrio*
 - Subgenus *Spirillum*
 - Subgenus *Spirochaete*
- B Species relatively pleomorphic
- I Rods and spirals Proteus types
 - Genus *Spirulina*
 - II Long filaments
 - a Unbranched
 - Genus *Leptothrix*
 - Subgenera
 - Leptothrix*
 - Beggiatoa*
 - Crenothrix*
 - Phragmidiothrix*
 - b Branched
 - Genus *Cladothrix*

The Cornil-Babes (1890) classification is another example of the carelessness in the definition of genera frequently to be found among the writers on medical bacteriology. The following outline will give the principal groups recognized, although generic names not here mentioned are used in the body of the text.

Cornil-Babes' Classification of Bacteria (1890)

- Group I *Micrococcus* Cells spherical
- 1 Cells in rosary *Streptococcus*
 - 2 Cells in irregular groups *Staphylococcus*
 - 3 Cells in firm gelatinous masses *Ascococcus*
 - 4 Cells in regular packets *Sarcina*
- Group II *Bacteriaceae* Short rods usually, may develop to filaments
- 1 Without capsules
 - a Not polymorphic *Bacterium*
 - b Polymorphic *Proteus*
 - 2 Encapsulated *Ascobacterium*
- Group III *Bacilli* Longer rods, producing spores Genera *Bacillus* and *Vibrio*
- Group IV *Spirobacteria* Cells spiral
Genera *Spirillum*, *Spirochaete*, *Leptothrix*, *Beggiatoa*, *Streptothrix*

In the same year also appeared the first of the classifications of bacteria used by Migula (1890). He gave no summarized statement, but the following outline shows his conception of generic relationships.

Migula's First Classification of Bacteria (1890)

- A Cells spherical
 - I Cells falling apart upon division *Mikrococcus*
 - II Cells remaining united in groups
 - a Forming irregular masses *Staphylococcus*
 - b Forming regular groups
 - 1 Cells in chains *Streptococcus*
 - 2 Cells in regular packets *Sarcina*
 - 3 Cells in flat plates, tetrads *Merismopedra*
 - 4 Cells in gelatinous masses *Leuconostoc*
- B Cells rod shaped
 - I No spores produced *Bacterium*
 - II Spores produced
 - a Cell when sporulating not spindle shaped *Bacillus*
 - b Cell when sporulating spindle shaped *Clostridium*
- C Cells spiral
 - I Cells relatively short
 - a Stiff, not flexuous *Spirillum*
 - b Flexuous *Vibrio*
 - II Cells relatively long and flexuous *Spirochaete*
- D Cells filamentous

Genera *Beggiatoa*, *Crenothrix*, *Cladothrix* and *Sphaerotilus*

Guignard (1890) named an organism responsible for the formation of large slimy masses on rocks on the seashore, *Snieblothrichia Bornetii*.

During the year 1891 at least three generic names were applied to bacteria as new. These were *Micromyces* by Gruber for a form of *Actinomyces*, *Eubacillus* by Dangeard for certain green bacteria (containing chlorophyll) and *Rhodococcus* by Zopf for the red cocci.

Ludwig (1892) outlined the following classification of bacterial genera. The number of genera recognized is smaller than those of some previous authors, but in some respects it represents a real advance.

Ludwig's Classification of the Bacteria (1892)

- I Cocci, with spherical non-motile cells, without endospores, or with a single endospore in the cell
 - A Not embedded in slime
 - 1 Simply or irregularly grouped *Micrococcus*
 - 2 Cells in pairs *Diplococcus*
 - 3 Cells in chains *Streptococcus*

- B Embedded in slime
 - 1 Single cells *Hyalococcus*
 - 2 Packets *Sarcina*
 - 3 Chains *Leuconostoc*
- II Forming rods, not long filaments
 - A Not embedded in slime
 - 1 Small elliptical cells *Bacterium*
 - 2 Small rods with endogenous spores *Bacillus*
 - 3 Spiral
 - a Stiff or rigid *Spirillum*
 - b Long and flexuous *Sprochaete*
 - B Embedded in slime
 - 1 Cells bent or spiral *Myconostoc*
 - 2 Straight rods *Cystobacter*
- III Forming long filaments, usually with a sheath
 - A Unbranched
 - 1 Non-motile and without inner spore formation *Leptothrix*
 - 2 Non-motile with sheath and inner spore formation *Crenothrix*
 - 3 Motile *Beggiatoa*
 - B Branched
 - 1 Branches not swollen at tips
 - a Free *Cladothrix*
 - b In slimy sheath *Sphaerotilus*
 - 2 Branches swollen at tips *Actinomyces*

Sternberg (1892) used a comparatively simple classification, merging genera in several cases. His conception of generic names however is in some cases inconsistent. The following key probably fairly indicates the principal points of difference as recognized by this author.

Sternberg's Classification of Bacteria (1892)

- A Cells spherical
 - I Cells not embedded in gelatinous masses
 - a Cell division in one direction
 - 1 Cells irregularly grouped *Micrococcus*
 - 2 Cells in pairs *Diplococcus*
 - 3 Cells in chains *Streptococcus*
 - b Cell division in two directions, forming tetrads *Merismopedia*
 - c Cell division in three planes, forming regular packets *Sarcina*
 - II Cells embedded in gelatin
 - a Cells separate in globular or lobulate zoogloal masses *Ascococcus*
 - b Cells solitary or in chains, with a thick gelatinous envelop *Leuconostoc*

- B. Rod shaped or filamentous, not spiral, no differentiation between the extremities of the rods
- I With or without spores, if the former not fusiform when sporulating.
 - Bacillus*
 - Clostridium*
 - II With spores, fusiform
- C Curved rods or spiral filaments
- I Not pleomorphic
 - a Cells not sinuous
 - 1 Cells rigid, not flexuous *Spirillum*
 - 2 Cells flexible *Vibrio*
 - b Cells with sinuous motion *Spirochaete*
- D Cells filamentous, with differentiated base and apex
- I Without pseudo-branches
 - Genera *Leptothrix*, *Beggiatoa*, *Crenothrix* and *Phragmidiothrix*
 - II With pseudo-branches *Cladothrix*

Attention should be called to the merging of all rod-genera (except *Clostridium*) into the genus *Bacillus*. This came in time to be the most commonly accepted of the medical definitions of the term.

Thaxter (1892) called attention to the existence of a new family of bacteria which he named the *Myxobacteriaceae*. The organisms belonging to this family show a more complex life history than the forms more commonly studied. The rod shaped cells are associated in a pseudo-plasmodium which in possession of motility resembles the plasmodium of a slime mold. The genera *Myxobacter* and *Myxococcus* were described.

During the same year the work of Winogradsky (1892) called to attention the important group of soil bacteria which oxidize ammonia to nitrites and nitrites to nitrates. To his work we owe the generic names *Nitrobacter*, *Nitrosococcus* and *Nitrosomonas*. The genus *Nevskia* was also described in the same year by Famintzin (1892) to include certain rod shaped bacteria which developed peculiar branched or lobed capsules due to the asymmetric deposition of the gummy material.

During the year 1893 the only generic name proposed was *Achromatum* by Schewiakoff for an organism of relatively monstrous proportions, which in its content of oxalates (?) simulated in appearance some of the sulphur bacteria.

Freudenreich (1894) recognized three genera of *Coccaceae*, *Micrococcus*, *Sarcina* and *Streptococcus* with their usual definitions, and in the *Bacteriaceae* the genera *Bacterium* and *Bacillus*, the latter with endospores, the former without.

Bernhard Fischer (1894) gave the generic name *Halbacterium* to nonphosphorescent bacteria from salt water.

The most important contribution of 1894 was the appearance of the second Migula classification. In practically identical outline but with some elaboration of subgenera this classification appeared the next year (1895) under the heading of *Schizomycetes* in Engler and Prantl's *Pflanzenfamilien*. A key to the families and genera of bacteria as recognized by Migula is as follows

Key to the Families of Bacteria Migula (1894)

- I Cells in free state spherical, not lengthening before division
Cell division in one, two, or three directions of space
Family I *Coccaceae*
- II Cells short or long cylindric, divide only transversely, and before division extending to double length
 - a Cells straight, rod shaped, without sheath, non-motile or motile by means of flagella Family II *Bacteriaceae*
 - b Cells bent, without sheath
Family III *Spirillaceae*
 - c Cells surrounded by a sheath
Family IV *Chlamydo bacteriaceae*
 - d Cells united into a filament, without sheath, motile by means of an undulating membrane Family V *Beggiatoaceae*

Key to Genera of the *Coccaceae*

- I Cells without organs of motion
 - a Division in one direction of space *Streptococcus* Billr
 - b Division in two directions of space *Micrococcus* (Hall) Cohn
 - c Division in three directions of space *Sarcina* Goodsir
- II Cells with organs of motion
 - a Division in two directions of space *Planococcus* Migula
 - b Division in three directions of space *Planosarcina* Migula

Key to the Genera of *Bacteriaceae*

- I Cells without organs of motion *Bacterium* Ehr emend
- II Cells with flagella
 - a Flagella over entire cell body *Bacillus* Cohn emend
 - b Flagella polar *Pseudomonas* Migula

Key to the Genera of the *Spirillaceae*

- I Cells stiff, not bending like a snake
 - a Cells non-motile *Spirosoma* Migula

- b Cells with flagella
 - 1 Cells with 1, rarely 2 or 3 polar flagella *Microspira* Schroeter
 - 2 Cells with tuft of polar flagella *Spirillum* Ehr
- II Cells flexuous *Spirochaeta* Ehr

Key to the Genera of Chlamydobacteriaceae

- I. Cells contents without sulfur granules
 - a Cell filaments unbranched
 - 1 Cell division continually in one direction of space *Streptothrix* Cohn emend
 - 2 Cell division before conidia formation in 3 directions of space
 - x Cells surrounded by a very delicate, scarcely visible sheath (marine) *Phragmidiothrix* Engler
 - xx Sheaths definitely recognizable (fresh water) *Crenothrix* Cohn
 - b Cell filaments branched *Cladothrix* Cohn
- II Cell contents with sulfur granules *Thiothrix* Winogr

Key to the Genera of the Beggiatoaceae

- One genus only *Beggiatoa* Trev

It will be noted that in Migula's classification emphasis is put largely upon the shape of the cell, motility and the distribution of flagella on the bacterial cells. Spores are not regarded as useful in the differentiation of genera. Neither the sulfur bacteria nor the Myxobacteria are recognized as distinct groups.

In 1895 appeared the first classification of Alfred Fischer. The principal differential characteristic will be noted in the family *Bacteriaceae* which is divided into sixteen genera based upon cell form, spore production, and location of the flagella. No species were known for several of the genera which he included. The classification of this family follows.

Fischer's Classification of Bacteria (1895)

Key to Genera of Bacteriaceae

- I Family *Baciller* Non-flagellate
 - a Producing endospores
 - 1 Spore bearing rod cylindric *Bacillus*
 - 2 Spore bearing rod spindle shaped *Paracloster*
 - 3 Spore bearing rod clavate *Paraplectrum*
 - b Producing arthrospores *Arthrobacter*
- II Family *Bactriniae* One polar flagellum
 - a Producing endospores
 - 1 Spore bearing rod cylindric *Bactrinum*

- | | |
|------------------------------------|------------------------|
| 2 Spore bearing rod spindle shaped | <i>Clostridium</i> |
| 3 Spore bearing rod clavate | <i>Plectrillum</i> |
| b Producing arthrospores | <i>Arthrobacterium</i> |
- III Family *Bactriller* With polar tuft of flagella
- | | |
|------------------------------------|-------------------------|
| a Producing endospores | |
| 1 Spore bearing rod cylindric | <i>Bactrillum</i> |
| 2 Spore bearing rod spindle shaped | <i>Clostrillum</i> |
| 3 Spore bearing rod clavate | <i>Plectrillum</i> |
| b Producing arthrospores | <i>Arthrobactrillum</i> |
- IV Family *Bactridae* With diffuse flagella
- | | |
|------------------------------------|-------------------------|
| a Producing endospores | |
| 1 Spore bearing rod cylindric | <i>Bactridium</i> |
| 2 Spore bearing rod spindle shaped | <i>Clostridium</i> |
| 3 Spore bearing rod clavate | |
| (a) Single spored | <i>Plectridium</i> |
| (b) With two spores | <i>Diplectridium</i> |
| b Producing arthrospores | <i>Arthrobactridium</i> |

The year 1896 was notable in the history of bacterial classification for the appearance of the scheme of Lehmann and Neumann. The outlines of classification of Migula, Fischer, and Lehmann and Neumann constitute the bases upon which practically all of the systematic work of the bacteriologists was built during the next two decades. Lehmann and Neumann separated from the true bacteria the bacteria-like *Hyphomycetes* and the filamentous water or algal bacteria. A key to the families and genera of the true bacteria is as follows:

Classification of Lehmann and Neumann (1896)

Bacteria or Schizomycetes

Small cells (almost) always free of chlorophyll, unbranched. Diameter usually not more than 2μ , rarely $3-5\mu$. Spherical, elongate or spiral, without organs other than flagella. Vegetative division usually transverse, very rarely longitudinal.

- I Cells spherical when free. Family I *Coccaceae* Zopf emend Migula
- | | |
|--|-------------------------------|
| a Cells divide only in one direction of space, forming chains if cells remain united | <i>Streptococcus</i> Billroth |
| b Cells divide on suitable media in three directions of space, forming packets | <i>Sarcina</i> Goodsir |
| c Cells divide irregularly, forming isolated cells or irregular masses | <i>Micrococcus</i> Cohn |
- II Cells at least $1\frac{1}{2}$, usually 2-6 times as long as broad, straight or slightly bent, never spiral. Family II *Bacteriaceae* Zopf emend Migula
- | | |
|--|------------------------------------|
| a Without endogenous spores, perhaps often with arthrospores. Rods usually under $0.8-1\mu$ diameter | <i>Bacterium</i> Cohn emend Hueppe |
|--|------------------------------------|

- b With endogenous spores Rods often more than 1μ in diameter
Bacillus Cohn emend Hueppe
- III Cells bent or spiral Family *Spirillaceae* Migula
 - a Cells short, slightly bent, stiff, comma-like, sometimes in spiral cell groupings, usually with one, rarely two terminal flagella
Vibrio O F Mueller emend Löffler
 - b Cells long, spirally bent, corkscrew like, stiff usually with a cluster of polar flagella
Spirillum Ehrenb emend Löffler
 - c Cells flexuous, long, producing spirally wound filaments Flagella unknown Motile by means of an undulating membrane
Spirochaete Ehrenb

As "Anhang I" three genera of the bacteria-like *Hyphomycetes* are included They may be differentiated as follows

Lehmann and Neumann's Classification of Hyphomycetes (1896)

- A Cultures showing ordinary bacterial character, flat or medium Microscopically rods with swollen ends *Corynebacterium* L & N
- B Cultures on solid media more or less folded, often "cartilagenous"
 - I Usually only short slender rods, seldom short branched filaments without air mycelium and without aerial conidia, acid fast
Mycobacterium L & N
 - II Mycelium filaments long, often bent, without sheath, with true branching Many species produce conidia on aerial hyphae Not acid fast.
Oospora Wallroth

The genera of the fission algae are differentiated as follows.

- A Filaments without sheaths
 - 1 Without sulphur granules *Leptothrix*
 - 2 With sulphur granules, motile *Beggiatoa*
- B Filaments with sheaths
 - 1 Without sulphur granules
 - a Without pseudodichotomous branching *Crenothrix*
 - b With pseudodichotomous branching *Cladothrix*
 - 2 With sulphur granules *Thiothrix*

In many ways this classification of Lehmann and Neumann is one of the most satisfactory that has been proposed

During the year 1896 Beijerinck introduced the generic name *Gramulobacter* for anaerobic spore-bearing bacteria of the butyric acid group, and Forti named an organism from wine *Oenobacillus Abbae*

Alfred Fischer (1897) somewhat modified his earlier classification in his "Vorlesungen über Bakterien" The key to the higher groups and genera is as follows

Fischer's Classification of Bacteria (1897)

Order I *Haplobacterinae*

Vegetative phase unicellular, spherical, cylindrical, or spirally twisted; isolated or united in chains or clusters

Family I *Coccaceae*

Vegetative cells spherical

Sub-Family I *Allococcaceae*

Planes of fission without definite sequence, no pronounced colonies or growth forms, cells isolated or in short chains or irregular clusters

Genus *Micrococcus*, Cohn Non-motile Includes most cocci, the pathological "staphylococci" etc

Genus *Planococcus*, Migula Motile

Sub-family II *Homococcaceae*

Planes of fission in definite sequence

Genus *Sarcina*, Goodsir Three planes of division at right angles to each other Cubical colonies, non-motile

Genus *Planosarcina*, Migula Similar to *Sarcina*, but monotrichous, and motile

Genus *Pedrococcus*, Lindner. Two planes of fission, alternate and at right angles

Genus *Streptococcus*, Billroth Planes of fission parallel, giving rise to chains, the pathological *Streptococci* and *Leuconostoc*

Family 2 *Bacillaceae*

Vegetative cell straight, cylindrical, ellipsoidal or egg-shaped, very short forms difficult to distinguish from cocci Fission always transverse

Sub-family I *Bacilleae*

Spore-forming rods cylindrical, unchanged

Genus *Bacillus*, Cohn Non-motile

Genus *Bactrinum*, A Fischer Motile, monotrichous, with terminal cilium includes provisionally all monotrichous rods whose spores are as yet unknown, e g, *Bac pyocyaneum*

Genus *Bactrillum*, A Fischer Motile, with lophotrichous ciliation Includes provisionally *Bac cyanogenum*, and many other sporeless forms

Genus *Bactridium*, A Fischer Motile, peritrichous, in some spores as yet unknown

Sub-family II *Clostridiae*

Rods spindle-shaped during sporulation

Genus *Clostridium*, Prazmowski Motile, peritrichous, includes some of the butyric bacteria. Genera with monotrichous and lophotrichous ciliation are unknown as yet

Sub-family III *Plectridae*

Genus *Plectridium*, A Fischer Motile, peritrichous, some butyric bacteria, the parasite of tetanus and a methane ferment

Family 3 *Spirillaceae*

Vegetative cell cylindrical but spirally twisted Fission always transverse

Genus *Vibrio* Muller and Löffler Very slightly curved rods, "comma" shaped, motile, monotrichous *Vibrio cholerae asiaticae* and numerous other vibrios of fresh and salt water

Genus *Spirillum*, Ehrenberg Cylindrical cells twisted in an open spiral, motile, lophotrichous *Spirillum undula*, *Sp rubrum*

Genus *Spirochaete*, Ehrenberg Cells long and attenuated, spirally twisted with numerous turns, cilia unknown, the cell membrane is perhaps yielding *Spirochaete Obermeieri* (remittent fever)

Order II *Trichobacterinae*

Vegetative phase an unbranched or branched filament or chain of cells, the individual members of which break off as swarm-spores (gonidia)

Family 1 *Trichobacteriaceae*

(a) Filaments non-motile, rigid, enclosed in a sheath

Genus *Crenothrix*, Cohn Filaments unbranched and devoid of sulphur granules

Genus *Thiothrix*, Winogradsky The same, but containing sulphur granules

Genus *Cladothrix*, Cohn Filaments, branched, false dichotomy (includes *Sphaerotilus*)

(b) Filaments motile, with oscillating and gliding movements, and devoid of a sheath

Genus *Beggiatoa*, Trevisan Containing sulphur

Chester (1897) adopted Migula's (1894) classification of *Coccaceae*, *Bacteriaceae* and *Spirillaceae*, but separated the forms included by Lehmann and Neumann in their *Hyphomycetes*, under the family name *Mycobacteriaceae* His characterization of this family and the genera is as follows

Chester's Classification of the *Mycobacteriaceae* (1897)

Cells short or long, cylindrical or clavate-cuneate in form without a sheath surrounding the chains of individuals Without endospores, with possibly the formation of gonidia like bodies With true dichotomous branching (this forms an important group standing between the true *Bacteriaceae*, and the true fungi on the one hand and *Chlamydobacteriaceae* on the other)

Order *Mycobacteriaceae* Chester

A In cultures possessing the characters of true bacteria Growth on solid media smooth, flat spreading Rods with swollen ends or cuneate or clavate forms

Corynebacterium Lehm-Neum

B Cultures on solid media raised, folded or warty Generally short slender rods, rarely short branched Take the tubercle stain

Mycobacterium Lehm-Neum

The following key to the families *Chlamydobacteriaceae* and *Beggiatoaceae* shows that in the main Chester followed Migula

Chester's Classification of the families *Chlamydobacteriaceae* and *Beggiatoaceae* (1897)

I Thread-like, composed of individual cells, surrounded by a sheath Simple or with true branching Ordinary vegetative growth by division in only one direction of space, i.e., at right angles to the longer axis

- A Cell content without sulfur granules
 - Order *Chlamydobacteriaceae*
 - 1 Filaments unbranched
 - a Cell division only in one direction of space
Streptothrix Cohn
 - b Cell division in gonidia formation in three directions of space.
 - x Cells surrounded by a very delicate, hardly discernible sheath (marine) *Phragmidiothrix* Engler
 - xx Sheath easily discernible (fresh water)
Crenothrix Cohn
 - 2 Filaments branched
Cladothrix Cohn
 - B Cell contents with sulfur granules *Thiothrix* Winogradsky
- II Thread-like, without a capsule, but with undulating membrane and as in *Oscillaria*, motile cell contents shows sulfur granules Formation of conidia not certainly known
 - Order *Beggiatoaceae*
 - A Threads apparently not septate, septa only faintly visible with iodine staining Colorless or faintly rose colored
Beggiatoa Trev

During this year (1897) Miyoshi reported three new genera of sulphur bacteria from hot sulphur springs, *Throderma*, *Throsphaera* and *Throsphaeron*. Ellis named a new iron bacterium *Spirophylum* and Meyer one of the bacteria of the *B subtilis* group *Astasia asterospora*

In this year also Migula published the first volume of his "System der Bakterien" containing a general discussion of bacterial morphology and physiology and an outline of the classification of genera essentially that of 1895

Hewlett (1898) in his Manual of Bacteriology recognized a small group only of genera. Of these the following were used in the sense of Zopf (1885) *Micrococcus*, *Streptococcus*, *Sarcina*, *Clostridium*, *Cladothrix*, *Leptothrix*. The genus *Bacillus* was defined to include all rod forms, *Spirillum* all spiral forms and *Streptothrix* all thread forms

Chester in 1899 revised his classification of the family *Mycobacteriaceae* to read as follows

Mycobacteriaceae Chester (1899)

- A Cells in their ordinary form as short cylindrical rods, often bent and irregularly swollen, clavate or cuneate At times Y shaped forms or longer filaments with true branchings May produce short coccoid elements, perhaps gonidia *Mycobacterium* Lehm-Neum (including, *Corynebacterium* L and N
- B Cells in their ordinary form as long branched filaments Produce gonidia-like bodies Cultures generally have a mouldy appearance due to the development of aerial hyphae
Streptothrix Cohn
Oospora Lehmann-Neumann

In this year a peculiar genus of bacteria was described by Jennings (1899) He used the name *Astrobacter* to designate an organism which multiplied by longitudinal fission, forming radial clusters The organisms were found only in certain stained preparations, it is not entirely certain that he was not observing an artifact or at least some organism not bacterial in nature

Migula (1900) in the second volume of his "System der Bakterien" considerably increased the scope of his classification by dividing the *Schizomycetes* into two classes which he termed the *Eubacteria* and the *Thiobacteria*, the true bacteria and the sulphur bacteria

Migula's Classification of Bacteria (1900)

- I Order *Eubacteria* Cells without any "Centralkorper" and free from sulfur and bacteriopurpurin
 - A Cells spherical Family *Coccaceae* Zopf emend Migula
 - 1 Cells non-motile
 - a Division in one direction of space
Streptococcus Billr
 - b Division in two directions of space
Micrococcus (Hallier) Cohn
 - c Division in three directions of space
Sarcina Goodsir
 - 2 Cells motile
 - a Division in two directions of space
Planococcus Migula
 - b Division in three directions of space
Planosarcina Migula
 - B Cells straight rods, at least never spiral
Family *Bacteriaceae*
 - 1 Non-motile
Bacterium
 - 2 Motile
 - a Flagella monotrichous *Pseudomonas* Cohn emend Smith
 - b Flagella peritrichous *Bacillus* Cohn em Migula
 - C Cells spirally wound or representing a portion of a spiral
 - 1 Non-motile *Spirosoma* Mig
 - a Not enclosed in gelatin
Subgenus *Euspirosoma* Mig
 - b Enclosed in gelatin Subgenus *Myconostoc*
 - 2 Motile
 - a One, two or three polar flagella
Microspira Schroeter
 - b Tuft of polar flagella
Spirillum Ehr
 - c Slender and flexuous
Spirochaete Ehr
 - D Cells cylindric, united into a sheathed thread
Family *Chlamydobacteriaceae*

- 1 Filaments unbranched
 - a No distinction between base and apex
Chlamydothrix Migula
 - b Filaments enlarged toward tips
 - (1) Sheath relatively thick *Crenothrix* Cohn
 - (2) Sheath relatively thin *Phragmidiothrix* Engler
 - 2 Filaments branched dichotomously *Sphaerotilus* Kutzling
- II Order *Thobacteria*, Cells without a "Centralkorper" but with sulfur inclusions Colorless, or rose, red or violet, with bacteriopurpurin
 - A Filamentous, destitute of bacteriopurpurin
Family *Beggiatoaceae*
 - 1 Threads attached, non-motile *Thothrix* Winogr
 - 2 Threads free, motile *Beggiatoa* Trev
 - B Containing bacteriopurpurin
Family *Rhodobacteriaceae*
 - 1 Cells united in families Division of cells in three directions of space
Subfamily *Thiocapsaceae*
 - a Families small compact, enclosed singly or several together in a cyst Capable of swarming
Thiocystis Winogr
 - b Non-swarming, globose cells spread out upon the substratum in flat families, loosely enveloped in a common gelatin
Thiocapsa Winogr
 - c Non-swarming cells arranged in packets like *Sarcina*
Thiosarcina Winogr
 - 2 Cells united into families Division of cells first in three, then in two, directions of space
Subfamily *Lamprocystaceae*
Families first solid, then globose hollow, perforated net
Lamprocystis Schroeter
 - C Cells united into families Divisions in two directions of space
Subfamily *Thiopediaceae*
Families tubular formed of cells arranged in fours
Thiopedia Winogr
 - D Cells united in families Division in one direction of space
Subfamily *Amoebobacteriaceae*
 - 1 Cells connected by plasma threads Families amoeboid motile
Amoebobacter Winogr
 - 2 Not as "1"
 - a Cells arranged in network, united by their ends
Thiodictyon Winogr
 - b Cells not arranged in network
 - (1) Capable of swarming Cells loosely aggregated in gelatin
Thiothece Winogradsky
 - (2) Non-motile, cells closely appressed in colony
Thiopolyoccus Winogr
 - E Cells free, capable of swarming at any time
Subfamily *Chromatiaceae*

- | | | | |
|--|-------------------|----------------|--------------------------------|
| 1 Cells cylindric-elliptic or elliptic | Moderately thick | Polar flagella | <i>Chromatium</i> Perty |
| 2 Cells rod shaped or spindle form, | flagella at poles | | <i>Rhabdochromatium</i> Winogr |
| 3 Cells spirally twisted | | | <i>Thiospirillum</i> |

Bejerinck (1900) suggested the generic name *Aerobacter* to include the gram negative lactic acid and gas producing rods of the *B. lactis aerogenes* and *coli* types Hartleb (1900) designated an organism from a root nodule as *Pseudorhizobium*, and Lagerheim named a bacterium found parasitic upon certain sea weeds, *Sarcinastrum*

Chester (1901) in his Manual of Determinative Bacteriology used the following family and generic keys

Chester's Key to the families of Bacteria (1901)

- I Cells unbranched, or show only a false branching in *Cladothrix*
 - A Cells in their free condition globular, becoming slightly elongated before division Cell division in one, two, or three directions of space *Coccaceae*
 - B Cells short or long, cylindrical, straight, curved, or spiral Without a sheath surrounding the chain of individuals, motile or non-motile, endospores present or absent *Bacteriaceae*
 - C Cells surrounded by a sheath, and arranged in elongated filaments *Chlamydobacteriaceae*
 - D Cells not surrounded by a sheath, arranged in elongated filaments, and motile by means of undulating membrane *Beggiatoaceae*
- II Cells short or long, cylindrical or filaments, often clavate, cuneate or irregular in form Without endospores, but with the formation of gonidia-like bodies due to the segmentation of the cells Without flagella Division at right angles to the axis of a rod or filament Filaments not surrounded by a sheath as in *Chlamydobacteriaceae* With true dichotomous branching. *Mycobacteriaceae*

Key to Genera of the Coccaceae

- A Cells without flagella
 - 1 Division in one direction of space only *Streptococcus* Billroth
 - 2 Division in two directions of space *Micrococcus* Hallier
 - 3 Division in three directions of space *Sarcina* Goodsir
- B Cells with flagella
 - 1 Division in two directions of space *Planococcus* Migula
 - 2 Division in three directions of space *Planosarcina* Migula

Key to the Genera of the Bacteriaceae

- A Flagella absent, endospores present or absent
Bacterium Ehr
- B Flagella present
 - 1 Flagella arising from any part of the body, peritrichic
Bacillus Cohn
 - 2 Flagella attached to one or both poles
Pseudomonas Migula

Key to the Genera of the Spirillaceae

- I Cells stiff, not flexile
 - A Without flagella
Spirosoma Migula
 - B With flagella
 - 1 With 1, rarely with 2 or 3 polar flagella
Microspira Schroeter
 - 2 With a bundle of polar flagella
Spirillum Ehr
- II Cells flexile
Spirochaeta Ehr

Key to the Genera of Mycobacteriaceae

Cells either short or long, cylindrical—clavate—cuneate in form, which at times may show true branching, or as long-branched mycelial-like filaments. Filaments not surrounded by a sheath as in *Chlamydobacteriaceae*. Without endospores, but with the formation of gonidia-like bodies due to a segmentation of the cells. Division at right angles to the axis of a rod or filament.

- A Cells in their ordinary form short cylindrical rods, often bent and irregularly swollen, clavate or cuneate. At times Y-shaped forms or longer filaments with true branchings. May produce short coccoid elements, perhaps gonidia *Mycobacterium* (Lehmann-Neumann), including *Corynebacterium* (Lehmann-Neumann).
- B Cells in their ordinary form as long-branched filaments. Produce gonidia-like bodies. Cultures generally have a mouldy appearance due to the development of aerial hyphae *Streptothrix* (Cohn), *Oöspora* (Lehmann-Neumann).

Key to the Genera of Chlamydobacteriaceae

Filamentous bacteria composed of rod-shaped cells, and surrounded by a distinct sheath. Division of the cells at right angles to the axis of the filaments. In *Phragmidiothrix* and *Crenothrix*, however, in the formation of gonidia, a division of the cells in three directions of space takes place. Reproduction by means of gonidia, which are either motile or non-motile.

- I Cell contents without sulfur granules
 - A Filaments unbranched
 - 1 Cell division takes place in one direction of space
Leptothrix

- 2 Cell division, before the formation of gonidia, takes place in three directions of space
 - a Filaments surrounded by a delicate, scarcely discernible sheath *Phragmadiothrix*
 - b Filaments surrounded by a plainly discernible sheath. *Crenothrix*
- B Filaments show false branchings *Cladothrix*
- II Cell contents contain sulfur granules *Thiothrix*

Lehmann and Neumann (1901) in the second edition of the *Grundriss* replace the designation of the Anhang I *Hyphomycetes* with the name *Actinomyces* and the generic name *Oospora* with *Actinomyces*. Otherwise the classification remained the same. During this year the generic name *Azotobacter* was suggested by Beijerinck for certain non-symbiotic nitrogen fixing bacteria of the soil. He also used the name *Urococcus* as a generic designation of certain urea-hydrolyzing cocci. Bongert used *Corynethrix* to designate an organism associated with mouse pseudotuberculosis.

Kendall (1902) proposed a revision of the Migula classification by the addition of certain genera to the family *Bacteriaceae*. The new genera were based entirely upon flagella distribution. A key to the genera recognized by this author follows.

Kendall's Classification of the Bacteria (1902)

- A *Coccaceae* Cells spherical in a free state, not elongate in any direction before division
 - 1 Non-motile
 - a Cells in chains, planes of fission parallel *Streptococcus* Billr
 - b Planes of fission without definite sequence *Micrococcus* Cohn
 - c Planes of fission three, at right angles *Sarcina* Goodsir
 - 2 Motile
 - a As in *Micrococcus* *Planococcus* Migula
 - b As in *Sarcina* *Planosarcina* Migula
- B *Bacteriaceae* Cells elongated, cylindrical, cells elongated before division
 - 1 Non-motile
 - a Not branching *Bacterium* Ehr
 - b Branching *Mycobacterium* Leh -Neum
 - 2 Motile
 - a Flagella peritrichic *Bacillus* Cohn
 - b Monotrichic *Bactrillus* Kendall

- c. Lophotrichic *Bacterium* Kendall
- d. Unknown *Bacterium* Kendall
- C *Spirillaceae* Cells cylindrical, spirally twisted
 - 1 Cells rigid, non-motile *Spirosoma* Migula
 - 2 Cells motile.
 - a Cell rigid
 - (a) with one or two polar flagella. *Microspira* Schroeter
 - (b) Lophotrichic *Spirillum* Ehr
 - b Cells flexuous *Spirochaete* Ehr
- D *Chlamydo bacteriaceae* Cells enclosed in a sheath
 - 1 Cell division always in one plane *Streptothrix* Cohn
 - 2 Cell division in three planes, very delicate sheath *Phragmidiothrix* Engler
 - 3 As "2", but sheath clearly visible *Ctenothrix* Cohn
 - 4 Cells more or less branched *Cladothrix* Cohn
- E *Beggiatoaceae* Cells without a sheath, motility by means of an undulating membrane Cells contain sulfur granules *Beggiatoa* Trev

Allan J Smith (1902) follows the Chester (1901) classification consistently

Matzuschita (1902) in his Bakteriologische Diagnostik adopts the following simple scheme

Matzuschita's Classification of Bacteria (1902)

- Coccaceae* Cells spherical
 - 1 Cells in chains *Streptococcus*
 - 2 Cells irregularly grouped *Micrococcus*
 - 3 Cells in packets *Sarcina*
- Bacillaceae* Rod shaped *Bacillus*
- Spirillaceae* Spiral or bent rods
 - Short *Vibrio*
 - Longer, relatively thick *Spirillum*
 - Long, very slender *Spirochaeta*

Klocker (translation 1903) in general employed the generic name applied by the one who first named the organism He recognizes the genera *Micrococcus*, *Pediococcus*, *Sarcina*, *Bacterium*, *Bacillus*, *Terrobacterium*, *Saccharobacillus*, *Clostridium* and *Granulobacter*

The classification used by Alfred Fischer (1903) in the second edition of his Vorlesungen is given below

Fischer's Classification of Bacteria (1903)

- A Order I *Haplobacterinae* Vegetative body, one-celled, spherical, cylindrical or spiral, single or united to unbranched chains and other growth forms

GENERAL SYSTEMATIC BACTERIOLOGY

I. Family I *Coccaceae* Vegetative cells spherical

- a. Subfamily, *Allococcaceae* Division in three directions of space, no sharply marked growth form, short chains, masses, in twos or singly

1. Non-motile *Micrococcus* Cohn
2. Motile *Planococcus* Migula

- b Subfamily *Homococcaceae* with definite division sequence for each genus

- 1 Division in three directions of space, forming packets of cells
 - (a) Non-motile *Sarcina* Goodsir
 - (b) Motile *Planosarcina* Migula
- 2 Division in two directions of space, forming fours or platelets *Pediococcus* Lindner
- 3 Division walls always parallel, grow in chains, pairs and singly, no plates or packets *Streptococcus* Billroth

II Family II *Bacillaceae* Vegetative cells cylindric, ellipsoid, oval, straight, during rapid multiplication, difficult to differentiate from cocci

- a Subfamily *Bacilleae* Spore forming rods unchanged, cylindric
Includes all forms not producing spores

- 1 Non-motile *Bacillus* Cohn
- 2 Motile
 - (a) Monotrichous *Bactrinum* A Fischer
 - (b) Lophotrichous *Bactrillum* A Fischer
 - (c) Peritrichous *Bactridium* A Fischer

- b Subfamily *Clostridiaceae* Spore forming rods, spindle shaped

- 1 Non-motile *Paracloster* A Fischer
- 2 Motile, peritrichous *Clostridium* Prazmowski

- c Subfamily *Plectridiaceae* Spore bearing rods, drumstick shaped

- 1 Non-motile *Paraplectrum* A Fischer
- 2 Motile, peritrichous *Plectridium* A Fischer

III Family III *Spirillaceae* Vegetative cells cylindric, but spiral

- a Slightly curved rods, motile, monotrichous

Vibrio (Muller-Loeffler)

- b Decidedly spiral in broad turns, motile, lophotrichous

Spirillum Ehr

- c Very slender, many turns, flagella unknown, perhaps cell wall flexible

Spirochaete Ehr

B Order II *Trichobacterinae* Branched or unbranched cell filaments, whose segments are separated off as swarm cells (gonidia) or homogones One family *Trichobacteriaceae*

- I Non-motile, rigid, enclosed in a sheath

- a Unbranched

- 1 Without sulfur inclusions

- (a) Not attached, cylindric gonidia, swarming
Chlamydothrix Migula

- (b) Attached, spherical gonidia not observed to be motile
Crenothrix Cohn

- 2 With sulfur granules, attached

Thiothrix Winogradsky

b Dichotomously pseudobranchied, lophotrichous cylindric gonidia

Cladothrix Cohn

II Filaments slowly moving, without sheath Containing sulphur

Beggiatoa Trev

During this year the name *Throphysa* was given by Hinze (1903) to a new genus of the sulphur bacteria, and Troih-Petersson named the lactic acid rods *Brachybacterium*

Migula in 1904 again outlined his classification promulgated in 1900 and vigorously defended it from the criticisms that had been made against it. New genera described during the year were *Hypnococcus* (Bettencourt, Kopke, Renzende and Mendes) for an organism found associated with sleeping sickness, *Clonothrix* (Schorler) for one of the iron bacteria, and *Thiobacillus* (Bejerinck) for organisms which could reduce nitrates, oxidize sulphur and utilize CO₂ in the manufacture of plant food.

Winslow and Rogers (1905) as the result of a careful study of many types and races of the cocci made some rather radical rearrangements of the groups, and created the new subfamilies *Paracoccaceae* and *Metacoccaceae*. New genera were *Aurococcus*, *Albococcus* and *Rhodococcus*. This classification was later further elaborated and published in 1910. A key to the families and genera recognized will be found under that year.

One of the most comprehensive analyses of the classification of bacteria which has been published in English is that of Erwin F. Smith (1905) in the first volume of his *Bacteria in Relation to Plant Diseases*. For his family and generic names he follows Migula (1900) consistently, except for several genera for which he substitutes older names. The genus *Pseudomonas* he concludes is a synonym of *Bacterium* Cohn. For *Bacterium* of the Migula system he proposes *Aplanobacter*. The genus *Microspira* Schroeter he concludes is a synonym of the older *Vibrio*. Furthermore he recognizes as a third order coordinate with *Eubacteria* and *Thiobacteria* the order *Myxobacteria*. His key to the genera of the *Myxobacteria* follows.

Smith's Classification of the Myxobacteria (1905)

- III Order *Myxobacteria* Motile rod-like organisms, multiplying by fission, secreting a gelatinous base, and forming pseudoplasmodium-like aggregations before passing into a more or less highly developed cyst-producing resting stage, in which the rods become encysted in groups without modification, or may be converted into spore masses.

- 1 Rods forming free cysts, in which they remain unmodified Cysts various, sessile or borne on a more or less highly developed cystophore
Chondromyces B and C
- 2 Rods forming large rounded cysts, one or more free without gelatinous matrix raised above the substratum *Polyangrium* K
- 3 Rods slender, swarming together after a vegetative period to form definite more or less encysted, sessile or stalked masses of coccus-like spores *Myzococcus* Thaxter

Particular attention was called to the group of spirochetes by the discovery by Schaudinn (1905) of the causal organism of syphilis. It was first named *Spirochaeta*, later Vullermin (1905) proposed *Spiro-nema*. This name was invalid under the rules of priority, so Schaudinn (1905) proposed *Treponema* and Stiles and Pfender (1905) a little later the name *Microspironema*. Two other new generic names were proposed during this year, *Erythrobaillus* by Fortineau for a red pyogenic rod, and *Semulostridium* by Maassen for a spore-producing, gelatin-forming bacterium from sugar vats.

Blanchard (1906) discussed the relationships of the spiral organisms, and concluded that they should be placed in two groups, the plant forms in the *Spirobacteria* and the forms more closely related to the protozoa under *Trypanosomidae*. The following key gives the principal points of differences recognized.

Blanchard's Classification of Spiral Organisms (1906)

- I *Spirobacteria*, Cohn 1875, Bacteria more or less curved, sometimes the curve forming only an arc of a circle, at other times forming spirals of several or many turns. Organisms little or not at all flexible, multiplying by transverse division. Endospores formed in a number of species.
 - A Non-motile *Spirosoma* Migula 1900
 - B Motile
 - 1 Cells rigid, forming a segment of a circle or united into spirals, 1, 2 or 3 polar flagella *Vibrio* Ehrenberg 1838
 - 2 Cells definitely spiral
 - a With endogenous spores. Flagella not polar *Spirobacillus* Metchnikoff 1859
 - b With or without endogenous spores. Flagella polar *Spirillum* Ehrenberg 1838
- II *Trypanosomidae* Doflein 1901. Flagellum continuous in the form of a spiral with turns more or less numerous. Organisms flexible, form more or less fixed, multiplying by longitudinal division. No endogenous spores. Gram —, not cultivable by ordinary means in laboratory.
 - A Form constant, greatly elongate
 - 1 Undulating membrane present, cells flattened *Spirochaeta* 1838 Ehrenberg

- b Cells straight rods, non-motile.
Genus *Rhodobacterium*
- c Cells motile rods
Genus *Rhodobacillus*
- d Cells short, bean or comma shaped, monotrichous, actively motile.
Genus *Rhodovibrio*
- e Cells spiral, actively motile by means of a cluster of polar flagella
Genus *Rhodospirillum*

This work of Molisch is practically the only work of importance on these bacteria since the first formulation of description and the grouping together of these organisms by Winogradsky

Other genera of bacteria described in 1907 include *Pedioplana*, a motile organism occurring in tetrads named by Wolff; *Thioploca* a filamentous sulphur organism related to *Beggiatoa*, named by Lauterborn; *Myrobacillus*, *Myxobacterium* and *Planobacterium*, organisms producing gums and slimes in sugar vats, named by Gonnermann; and *Borrelia* for the spirochetes of relapsing fever named by Swellengrebel. The last named classifies the spiral organisms as follows

Swellengrebel's Classification of the Spirillaceae (1907)

Cells with rounded tips, constituting any part of a spiral

- I Subfamily *Spirillaceae* (nov fam) Cells are not flexible Genera, *Spirillum* and *Vibrio* with diagnoses of Migula
- II Subfamily *Spirochaetaceae* (nov fam) Cells are flexible
 - a Cells without cilia with periplastic appendage well developed and often with an alveolar structure Sometimes there is a band of myonemes in the appendage
Spirochaeta Ehrenb
Type *Spirochaeta plicatilis* Ehrenb *S. balbiani* placed here provisionally
 - b Cells with one cilium at one end, sometimes at both extremities which is a prolongation of the 'calotte' the appendage has been demonstrated sometimes
Treponema Schaudinn
Types (*S. buccalis* Tr *pallidum*)
 - c Cells with peritrichous cilia
Borrelia n gen
Type, *Sp. gallinarum*

Ellis in 1907 as a result of studies on the iron bacteria came to the conclusion that they should be separated into two principal groups in accordance with the following scheme

Ellis' Grouping of the Iron Bacteria '1907'

- I. Reproduction by external abstriction of conidia
 - Leptothrix ochracea*
 - Gallionella ferruginea*
 - Spirophyllum ferrugineum*

II Reproduction by separation of internally produced cells

*Crenothrix polyspora**Cladothrix dichotoma**Clonothrix fusca*

The Winslows (1908) presented the final results of their studies on the *Coccaceae*. A key to the subfamilies and genera recognized follows

Winslows' Classification of the Coccaceae (1908)

Cells spherical Family *Coccaceae*

- I Parasites Growth not abundant (or one species, zooglea-forming saprophytes growth abundant in saccharose media) Generally gram+ Acid formers
 - Subfamily *Paracoccaceae*
 - A Cells in capsulated pairs Parasites Growth very meagre Inulin fermented No pigment
 - Diplococcus* (Weichselbaum) Winslow and Rogers
 - B Cells in chains occurring in masses of zoogloea in sugar refineries Aberrant saprophytic forms Growth abundant in saccharose media No pigment
 - Ascococcus* (Cohn) Winslow and Rogers
 - C Cells in chains Parasites Growth meagre Inulin not fermented No pigment
 - Streptococcus* (Billroth)
 - D Cells in irregular groups
 - Staphylococcus*
 - 1 Parasites Growth fair Orange pigment
 - Aurococcus* Winslow and Rogers
 - 2 Parasites Growth good White pigment
 - Albococcus* Winslow and Rogers
 - II Saprophytes Growth abundant No zoogloea Gram—generally Not acid formers
 - Subfamily *Metacoccaceae*
 - A Cells in irregular groups Pigment generally yellow
 - Micrococcus* (Hallier, Cohn), Winslow & Rogers
 - B Cells in packets Pigment yellow
 - Sarcina* (Goodsir) Winslow and Rogers
 - C Cells in irregular groups or packets, pigment red
 - Rhodococcus* Winslow and Rogers

Flügge (1908) in the classification of the pathogenic bacteria given in his *Grundriss der Hygiene* (sixth edition) made a distinct contribution to systematic bacteriology in the groupings which he created. From a strict nomenclatural point of view his terminology was in many cases technically incorrect, but the groups are very useful. His classification is as follows

Flügge's Classification of the Pathogenic Bacteria

(Lehrbuch der Hygiene, 6th ed., 1907, Quoted in Kolle and Hetch, 3rd ed. 1911, p. 18)

I Coccaceae Cells spherical

A Family *Streptococcus* Grow only in one direction, are Gram +, non-motile and grow only sparingly on artificial media

1 *Diplococcus* type, forms in many media only round or lancet shaped diplococci, in others (bouillon) short chains Pathogen *Diplococcus lanceolatus capsulatus*

2 *Streptococcus* type In bouillon generally forms long chains Saprophytes, *Streptococcus mesenteroides*

B Family, *Sarcina* Numerous species, in part motile and flagellated, dividing in three directions of space, forming packets, Gram - on solid media, growing usually in the form of dry layers, often colored, often in dust

Pathogens None

C Family, *Micrococcus* Numerous species, of which many are saprophytic, found in dust Cells divide irregularly in different directions of space and are often united in bunches of 2, 4, 8 or irregularly

1 *Diplococcus* type Elongated cocci with the long side appressed Gram negative

Pathogens *Micrococcus gonorrhoeae*, *Micrococcus intracellularis meningitidis*, *Micrococcus catarrhalis*

2 *Tetragenus* type Forms cocci that remain in twos or more, frequently remain in fours

Pathogen *Micrococcus tetragenus*

3 *Staphylococcus* type Spherical Grows on solid and fluid nutrient media in regular masses with white yellow or orange pigment

Pathogen *Staphylococcus pyogenes aureus*

II Bacillaceae Rod shaped Not spiral

A Family, *Bacillus* All rods which produce endogenous spores

1 Aerobic forms

a *Hay bacillus* group Relatively large bacilli widely distributed, grow well on solid and liquid media, on the latter with membrane formation Usually motile Non-pathogens

b *Anthrax* group As "a" but non-motile and pathogenic

2 Anaerobic forms Includes all the spore bearing types of butyric acid bacteria

Pathogens *B. botulinus* Bacillus of blackleg of edema of tetanus

B Family *Bacterium* All rods that produce no spores

1 *Fluorescent* phosphorescent, and pigment producing forms Gram negative, may or may not liquefy gelatin

2 *Colon bacillus* group possesses flagella more or less motile Gram negative

Pathogens *Bact. typhi*, and *Bact. paratyphi*

- 3 *Bact aerogenes* group, very near *B coli* (but non-motile?)
Pathogen *Bact dysenteriae*
- 4 Plague and septicemia group Non-motile, polar staining Gram negative, do not liquefy gelatin
Pathogen *Bact pestis*
- 5 *Influenza* group Very small Gram—, growing only on media containing hemoglobin
Pathogen *Bact influenzae*
- 6 *Swine erysipelas* group Very small rods, do not liquefy gelatin. Gram positive
Pathogen *Bact Rhusiopathiae*, Rabbit septicemia
- 7 *Glanders* group Very slender rod Gram negative
Pathogen *Bact mallei*
- 8 *Diphtheria* group Gram positive Tendency to the production of club shaped involution forms and to granular disintegration.
Pathogen, *Bact diphtheriae*, *Bact zeroeosis*
- 9 *Acid fast* group Producing slimy wrinkled layers on the surface of solid and liquid media Gram positive and acid fast
Pathogen *Bact tuberculosis*

III *Spirillaceae* Cells spiral

- A Family *Vibrio* All comma-shaped, bent, fission fungi with polar flagella Often united to spirals Gram negative
Pathogen *Vibrio cholerae asiaticae*
- B Family *Spirillum* Several corkscrew turns Polar flagella Many saprophytes
Pathogens None known

The classification of the *Trichomycetes* or thread bacteria suggested by Jordan (1908) is worthy of note He diagnosed the genera as follows

Jordan's Classification of the *Trichomycetes* (1908)

- 1 Filaments unbranched *Leptothrix*
- 2 Filaments with pseudobranches *Cladothrix*
- 3 Filaments with true branches
 - a Reproductive elements, spores observed *Nocardia*
 - b No spores observed *Actinomyces*

Conn (1909) suggested the following classification

Conn's Classification of Bacteria (1909)

Spherical Bacteria

- Dividing in *one plane*, so as to form chains, *Streptococcus*
- Dividing in *two planes*, and not forming chains, *Micrococcus*
- Dividing in *three planes*, and forming cubical masses, *Sarcina*

Rod Shaped Bacteria

- With flagella, and consequently motile, *Bacillus*

Without flagella and consequently non-motile, *Bacterium*

With a single flagellum, *Pseudomonas*

Spiral Bacteria: *Spirillum*

Higher Bacteria (*Cladothrix*, *Leptothrix*, *Streptothrix*, *Actinomyces*)

The genus *Bacillus* is further divided as follows

Bacilli with *one flagellum* are named *Monotrichic* Bacilli or *Pseudomonas*

Bacilli with *one flagellum* at each end, *Microsporon*

Bacilli with a *tuft of flagella* at one end are called *Lophotrichic* Bacilli

Bacilli with flagella *over the whole body* are called *Peritrichic* Bacilli

Orla-Jensen in 1909 proposed a totally new classification of bacteria. He divided the bacteria into two orders based upon primitive or advanced characteristics, distribution of flagella, etc. In many cases his published descriptions of genera are not very clear, and the construction of a key for the differentiation of the forms is correspondingly difficult. The following key will indicate differences between most of the genera described

Orla-Jensen's Classification of Bacteria (1909)

Key to Orders and Families

- I Cells spherical, rod-shaped or spiral Endospores formed only in a few sulfur-free spirilla When motile, mono- or lophotrichous Typically water bacteria Secure energy almost exclusively by oxidative processes Many develop poorly on laboratory media because organic substances are present in too concentrated a form

Order I *Cephalotrichinae*

- A Bacteria not in thread like filaments, no tendency to branch or produce mycelial growth

- 1 Bacteria not containing sulfur or bacteriopurpurin

- a Obligate aerobes oxidizing carbon hydrogen, or nitrogen compounds without production of noteworthy amounts of unoxidized split products

Family 1 *Oxydobacteriaceae*

- b Luminous and fluorescent bacteria

Family 6 *Luminibacteriaceae*

- c Active reducing organisms, facultative, all spiral forms reducing sulfates

Family 7 *Reduibacteriaceae*

- 2 Bacteria containing sulfur or bacteriopurpurin

- a Containing no bacteriopurpurin

Family 3 *Thiobacteriaceae*

- b Containing bacteriopurpurin

Family 4 *Rhodobacteriaceae*

- B. Bacteria tending to branch, to form filaments or to develop a mycelium

- 1 Not typically water forms Cells tending to branch and to form branching hyphae Not forming filaments usually
Family 2 *Actinomycetes*
 - 2 Typically water forms Cells not tending to form mycelium
Typically filamentous
Family 5 *Trichobacteriaceae*
- II Cells spherical or rod shaped, never spiral Either peritrichous or non-motile Not typically the water bacteria Do not secure growth energy solely by oxidative processes
- Order II *Peritrichinae*
- A Typically not obligate anaerobic or microaerophilic
 - 1 Usually bringing about transformation in carbohydrates with the development of acids
Family 1 *Acidobacteriaceae*
 - 2 Usually bringing about an alkaline reaction by the development of ammonia
Family 2 *Alkalibacteriaceae*
 - B Typically obligate anaerobic or microaerophilic
 - 1 Acting largely upon carbohydrates, developing acids, particularly butyric acid
Family 3 *Butyrbacteriaceae*
 - 2 Acting largely upon proteins, bringing about putrefactive changes
Family 4 *Putribacteriaceae*

Order I *Cephalotrichinae*

Key to the Genera of Family I Orydobacteriaceae

- I Not oxidizing nitrogen compounds nor fixing free nitrogen
 - A Oxidizing methane to CO_2 and H_2O *Methanomonas*
 - B Oxidizing CO to CO_2 *Carboxydomonas*
 - C Oxidizing H_2 to H_2O *Hydrogenomonas*
 - D Oxidizing alcohol to acetic acid *Acetimonas*
- II Oxidizing ammonia or nitrites
 - A Oxidizing ammonia *Nitrosomonas*
 - B Oxidizing nitrites *Nitromonas*
- III Fixing atmospheric nitrogen *Azotobacter*

Key to the Genera of Family II Actinomycetes

- I Not producing a much branched mycelium, individual cells only branching
 - A Symbiotic in the nodules of legumes *Rhizomonas*
 - B Not symbiotic Parasites in mammals
 - 1 Not acid fast *Corynemonas*
 - 2 Acid fast *Mycomonas*
- II Producing a much branched mycelium *Actinomyces*

Key to the genera of Family III Thiobacteriaceae

- I Cells not spiral
 - A Cells rod-shaped, or not spherical
 - 1 Autotrophic, rod-shaped *Sulphomonas*
 - 2 Not autotrophic, oval (?) *Thiomonas*

GENERAL SYSTEMATIC BACTERIOLOGY

- B Cells spherical
- II Cells spherical

Thiococcus
Thiospirillum

Key to the Genera of Family IV *Rhodobacteriaceae*

Uses the following genera with the same meanings as Winogradsky or Molisch.

Rhodomonas (*Chromatium*)
Rhabdomonas (*Rhabdochromatium*)
Rhododictyon (*Thiodictyon*)
Amoebomonas (*Amoebobacter*)
Rhodothece (*Thiothece*).
Rhodopolycoccus (*Thioplococcus*).
Rhodococcus (*Thiopedia*)
Lamprocystis
Rhodocystis (*Thiocystis*)
Rhodocapsa (*Thiocapsa*)
Rhodosarcina (*Thiosarcina*)
Rhodospirillum (*Thiospirillum*)

Key to the Genera of Family V *Trichobacteriaceae*

- I Not containing sulfur granules
 - A. Branched (with pseudobranches) *Cladotrichum*
 - B Not Branched
 - 1 Not spiral
 - a Not attached *Leptothrix*
 - b Attached *Crenothrix*
 - 2 Spiral
 - a Slender and flexuous *Spirochaeta*
 - b Flattened *Sporophyllum*
- II Containing sulfur granules
 - A Not attached Motile *Beggiatoa*
 - B Attached *Thiothrix*

Key to the Genera of Family VI *Luminobacteriaceae*

- I Cells not spiral
 - A Cells rod-shaped
 - 1 Active denitrifiers *Denitromonas*
 - 2 Active liquefiers *Liquidomonas*
 - B Cells spherical
 - 1 Liquefying gelatin *Liquidococcus*
 - 2 Not liquefying gelatin *Solidococcus*
- II Cells bent or spiral *Liquidovibrio*

Key to the Genera of Family VII *Reductobacteriaceae*

- I Short spiral Not liquefying gelatin *Solidovibrio*
- II Longer spirals *Spirillum*

Order II. Peritrichinae

Key to Genera of Family I. *Acidobacteriaceae*

- | | | |
|----|---|--------------------------|
| I | Active denitrifying forms | <i>Denitrobacterium</i> |
| II | Not as in I | |
| | A Rod shaped | |
| | 1 Organisms of the <i>B. coli</i> group | <i>Bacterium</i> |
| | 2 Gram + lactic acid forms | <i>Caseobacterium</i> |
| | 3 Producing propionic acid | <i>Propionibacterium</i> |
| | B Spherical | |
| | 1 In chains | <i>Streptococcus</i> |
| | 2 Irregularly grouped | <i>Micrococcus</i> |
| | 3 In packets | <i>Sarcina</i> |

Key to the Genera of Family II *Allahbacteriaceae*

- | | | |
|-----|--|-------------------------|
| I. | Organisms of the <i>Proteus</i> type | <i>Liquidobacterium</i> |
| II | Organisms of the <i>B. subtilis</i> type | <i>Bacillus</i> |
| III | Producing ammonia from urea | <i>Urobacillus</i> |

Key to the Genera of Family III *Butyribacteriaceae*

- | | | |
|-----|--|------------------------|
| I | Acting primarily upon sugars, forming butyric acid | <i>Butyribacillus</i> |
| II | Acting primarily upon pectin | <i>Pectobacillus</i> |
| III | Acting primarily upon cellulose | <i>Cellulobacillus</i> |

Key to the Genera of Family IV *Putribacteriaceae*

- | | | |
|----|----------------------------|-----------------------|
| I | Not producing an ectotoxin | <i>Putribacillus</i> |
| II | Producing an ectotoxin | <i>Botulobacillus</i> |

It will be noted that Orla-Jensen has created a considerable number of generic names, many times substituting new names for older and apparently perfectly valid terms

Among the new genera described in 1909 by other authors were *Hillhousia* a giant sulphur organism named by West and Griffiths, *Siderocapsa*, a capsulated epiphytic coccus growing in water containing iron and with iron oxid deposition in the capsule, named by Molisch, and *Erysipelothrix* an organism of the general type of the bacterium of swine erysipelas named by Rosenbach

Molisch (1910) in his book "Die Eisenbakterien" recognizes the genera and species separated by the following key

Molisch (1910) Classification of the Iron Bacteria

- | | | |
|---|-------------------------------|---------------------------------|
| A | Filamentous bacteria | |
| | I Filaments always unbranched | |
| | a With brown attaching disks | <i>Chlamydothrix sideropous</i> |
| | b Not as "a" | |

- 1 With definite sheath, often broader toward end Not spirally wound *Crenothrix polyspora*
 - 2 Without definite sheath Spirally wound *Gallionella ferruginea*
- II Filaments unbranched or pseudodichotomously branched
- a Swarm cells developed
 - 1 Regularly pseudodichotomously branched, usually colorless. *Cladothrix dichotoma*
 - 2 Usually unbranched, usually encrusted with iron *Chlamydothrix ochracea*
Clonothrix fusca
 - b Only non-motile gonidia
- B Capsule bacteria
- I Capsule with only a few (1-8) cells *Siderocapsa Treubii*
 - II Capsule with 1-100 cells *Siderocapsa major*

Among the new generic names of bacteria proposed in 1910 were the following *Fusiformis* for the fusiform bacteria of the throat named by Hoelling, *Asterococcus* for the very minute organism responsible for the disease pleuropneumonia in cattle, named by Borrel *et al*, *Actinobacillus* by Brumpt for the causal organism of actinobacillosis of cattle, *Nodofolium* for one of the thread iron bacteria named by Ellis and *Cristispira* given by Gross as designation of the spiral "crested" organism from the crystalline styles of certain classes of mussels

Dobell (1911) included the spirochetes in the group *Spirochaetoidia*, using the following generic differences for the three genera recognized. He emphasizes that this group of *Protista* should be included neither with the bacteria or protozoa

Spirochaetoidia

- 1 Free living forms, fresh water or marine *Spirochaeta* Lhr
- 2 Parasitic in animals (Vertebrates and Invertebrates) *Treponema* Schaudinn
- 3 Parasitic in Lamellibranchia *Cristispira* Gross

In Doflein's *Lehrbuch der Protozoenkunde* (1911) the same problem is handled by considering *Spirochaeta* as a generic name which includes all forms, and regarding *Spirochaeta*, *Cristispira* and *Treponema* as the component subgenera. Gross in this year added the genus *Saprospira* to this group. He included it in the free living marine types.

Frost in Marshall's *Microbiology* (1911) used Migula's (1900) scheme of classification in its entirety. Heim in the 4th edition of his *Lehrbuch der Bakteriologie* gives a very simple classification suitable particularly for the pathogenic bacteria.

Heim's Classification of Bacteria (1911)

- I *Coccaceae* Cells spherical
 - A Division in one direction of space *Streptococcus*
 - B Division in different directions, no chains or packets *Micrococcus*
 - C Dividing in three planes, forming packets *Sarcina*
- II *Bacteriaceae*, Rod-shaped
 - A Without spores *Bacterium*
 - B With spores *Bacillus*
- III *Spirillaceae* Cells spiral
 - A Rigid spirals or sections of spirals *Spirillum* and *Vibrio*
 - B Flexuous, Spirals *Spirochaete*

Kolle and Hetsch (1911) differentiate the bacteria into the fission fungi consisting of cocci, bacilli and spirilla and the fission algae including the sulphur bacteria, the iron bacteria and the *Streptothricaceae*

Bosanquet (1911) in his book on Spirochaetes places all in the genus *Spirochaeta*. During this year two other bacterial generic names were proposed *Paraspirillum* for a flexile spindle formed spiral organism in water (Dobell) and *Cohnistreptothrix* for organisms of the type of *Streptothrix Foersteri* of Cohn (Pino and Morax)

Schneider (1912) follows the classification used by Chester (1901). Benecke (1912) gives one of the most complete of the recent systems of bacterial classification. The orders and families recognized are shown in the following scheme

Benecke's Classification of Bacteria (1912)

- A Comparatively simple organisms, not growing as sheathed filaments, usually as single cells or colonies Order I *Haplobacterinae*
 - I Cells not forming a pseudoplasmodium
 - a Cells not tending to branch or form a mycelium
 - 1 Without bacteriopurpurin
 - (a) Cells spherical Family I *Coccaceae*
 - (b) Cells elongate
 - (1) Cells straight Family II *Bacillaceae*
 - (2) Cells spiral Family III *Spirillaceae*
 - 2 With bacteriopurpurin Family IV *Rhodobacteriaceae*
 - b Cells tending to branch or form a mycelium
 - Family V *Mycobacteriaceae*

GENERAL SYSTEMATIC BACTERIOLOGY

II Cells in vegetative stage forming a pseudoplasmodium

Family IV *Myxobacteriaceae*

B Cells in filaments, frequently sheathed

Order II *Desmobacterineae*

One family only

Family VII. *Desmobacteriaceae*

While no new genera are described, a larger number of genera are included in several of the families than by most other writers

Engler (1912) in the seventh edition of his syllabus gives the following outline of bacterial classification—

Engler's Classification of Bacteria (1912)

Schizomycetes

I Order *Eubacteria* Cells without sulfur and bacteriopurpurin

A Family, *Bacteriaceae* Cells very small, cylindric, straight or slightly bent

1 Without flagella *Bacterium*

2 With flagella

a Peritrichous *Bacillus*

b Polar *Pseudomonas*

B Family, *Spirillaceae* Spiral cells

1 Cells stiff, not serpentine

a Non-motile *Spirosoma*

b Motile

(1) With 1 2 or 3 polar flagella

Microspira

(2) With bunch of polar flagella

Spirillum

2 Cells serpentine *Spirochaete*

C Family, *Phycobacteriaceae* (*Chlamylobacteriaceae*) Filamentous surrounded by a more or less definite sheath and dividing seldom in three directions Multiplication by conidia

Genera *Phragmidiothrix* *Chlamydothrix*, *Gallionella* (*Crenothrix*, *Clonothrix*, *Sphaerotilus* (*Cladothrix*))

D Family, *Actinomycetaceae* Filamentous colonies with true branching radiating non-motile Filaments dividing up into nodules

One genus

Actinomyces

E Family *Coccaceae* Cells spherical

Genera, *Streptococcus* *Micrococcus* *Leuconostoc* *Sarcina* *Azotobacter*, *Planococcus*

F Family, *Myxobacteriaceae* Cells more or less rod shaped, non-flagellate, with slimy membrane moving slowly forward (Pseudoplasmodia) which become a cyst, sessile or stalked

Genera *Polyangium*, *Chondromyces*, *Myxococcus*

II Order, *Thiobacteria* Cells with sulfur inclusions, colorless or red or violet with bacteriopurpurin

A. Family, *Beggiatoaceae* Motile by an undulating membrane, filamentous colonies of colorless, sheathless cells, with sulfur granules.

Genera *Beggiatoa*, *Thiothrix*

B Family *Rhodobacteriaceae* Cell content rose, red, or violet with sulfur granules

Genera, *Lamprocystis*, *Thiosarcina*, *Rhodobacillus*, *Thiospirillum*, *Chromatium*

Meyer (1912, p 2) recognized as an order the *Eubacteria* the genera and generic sections of which may be separated by the following key

Key to the genera and sections of the Eubacteria according to Meyer (1912)

- a Cells straight rods
 - b Not with polar flagella Flagella when present-peritrichous
 - Genus *Bacillus*
 - c Cells containing bacteriopurpurin
 - Section *Rhodobacillus*
 - 2c Cells not containing bacteriopurpurin
 - d Flagella unknown
 - Section *Bacterium*
 - 2d All other species
 - Section *Eubacillus*
 - 2b With polar flagella
 - Genus *Pseudomonas*
- 2a. Cells not straight rods
 - b Cells curved rods
 - c Peritrichous flagella, with spores
 - Genus *Spirobacillus*
 - 2c Polar flagella, spores unknown
 - d With 5-20 polar flagella
 - Genus *Spirillum*
 - e Without bacteriopurpurin
 - f Flagella unknown
 - Section *Spirosoma*
 - 2f Flagella known
 - Section *Euspirillum*
 - 2e With bacteriopurpurin
 - Section *Rhodospirillum*
 - f Sulphur free
 - Subsection α
 - 2f With sulphur
 - Subsection β
 - 2d With 1 rarely 2-3 polar flagella
 - Genus *Microspira*
 - e With bacteriopurpurin
 - Section *Rhodospira*
 - 2e Without bacteriopurpurin
 - Section *Eumicrospira*

2b. Cells spherical

c Cells in chains . Genus *Streptococcus*

d Without bacteriopurpurin.

e With flagella.

Section *Planostreptococcus*

2e Without flagella

Section *Fixostreptococcus*

d. With bacteriopurpurin

Section *Rhodococcus*

e Without sulphur

Subsection α

2e With sulphur.

Subsection β 2c. Cells not in chains Genus *Micrococcus*

d Division in two planes

e Without bacteriopurpurin

f Flagella unknown

Section *Fixococcus*

2f Flagella known

Section *Planococcus*

2e With bacteriopurpurin

Section *Rhodococcus*

f With sulphur

Subsection α

2d Division in three planes

Genus *Sarcina*

e Without bacteriopurpurin

f With flagella

Section *Planosarcina*

2f Without flagella

Section *Fixosarcina*

2e With bacteriopurpurin

Section *Rhodisarcina*

f Without sulphur

Subsection α

2f With sulphur

Subsection β

Lohnis (1913) in his "Vorlesungen über landwirtschaftliche Bakteriologie" gives the following abbreviated classification of genera

Lohnis' Classification of Bacteria (1913)

I. Cells as a rule spherical, seldom assuming an elongated form

a. Single, in pairs, fours, or irregular masses, never in chains

Micrococcus

b In shorter or longer chains

Streptococcus

c. In packets

Sarcina

- II Cells as a rule rod shaped, seldom spherical, or bent
 - a Without endospores *Bacterium*
 - b With endospores *Bacillus*
- III Cells as a rule slightly to decidedly spirally bent
 - a Comma shaped *Vibrio*
 - b Stiff spirals *Spirillum*
 - c Flexose spirals *Spirochaete*

Chatton and Perard (1913) introduced the generic names *Meta-bacterium* and *Oscillospira*

Vuillemin (1913) has given a comprehensive review of bacterial classification with particular reference to the fact that at the Botanical Congress that was to have met in 1915 the date of departure and special rules for bacterial nomenclature were to be considered. He suggested that the date of departure be fixed as 1915 and that the genera and higher groups that follow should be adopted as *genera conservanda*, or "natural history" genera

Vuillemin's Classification of Bacteria (1913)

- I. Plastids rounded (spherical)
 - A Dissociated or grouped without regular order
 - 1 Polar flagella *Planococcus*
 - B Associated in elementary colonies of determined form
 - 1 Divisions parallel
 - a Grouped in rosaries *Streptococcus*
 - b Pairs encapsulated, isolated or united into a rosary Elements sometimes elongated or pointed *Klebsiella*
 - 2 Division in two planes at right angles
 - a Fission centripetal
 - (1) Non-motile *Merista*
 - (2) Flagellated *Planomerista*
 - b Fission excentric *Neisseria*
 - 3 Division in three planes at right angles
 - a Non-motile *Sarcina*
 - b Flagellated *Planosarcina*
- II Plastids rod-shaped
 - A Sporiferous elements specialized
 - 1 Oval Flagella unknown *Metabacterium*
 - 2 Claviform Flagella diffuse *Clostridium*
 - B None of the sporiferous elements distinct from the vegetative rods
 - 1 Flagella diffuse *Serratia*
 - 2 Flagella polar *Bacterium*
- III Plastids sinuous
 - A Flagella polar *Spirillum*

Vuillemin also proposed that the following names be kept as *openera conservanda*," but without nomenclatural status, *Micr. Mantegazzaea*, *Bacillus* and *Spirosoma*.

Gonder (1914) in Prowazek's *Handbuch der pathogenen Protozoen* gives the following classification of the spirochetes

Gonder's Classification of the Spirochetes (1914)

Spirochaeta Ehrb 1838

Type species *Sp. plicatilis* Ehr

Cristispira Gross 1910

Type species, *Cristispira balbiani*

Spironema Vuillemin 1905

Type species *Spironema recurrentis*

Treponema Schaudinn 1905

Type species *Treponema pallidum*

Buchanan (1915) reviewed the terminology for the *Coccaceae* proposed by Winslow and Rogers and suggested certain changes that should be made in names of genera and higher groups the better to recognize priority

Buchanan (1917-1918) published a series of articles under the general heading of "Studies in the Nomenclature and Classification of the Bacteria," in which were listed the various names of genera and higher groups which had been recognized in the literature, and a classification of bacteria into orders, families and genera, with designation of type species for the genera. The dichotomous keys to the groups recognized follow

Key to the Orders of the Class Schizomycetes 'Buchanan' 1917

A Plant-like in the principal characters not protozoan like cells never slender, flexuous spirals, cell divisions never longitudinal

I Not producing a pseudoplasmodium during the vegetative stage without a highly developed cyst-producing resting stage

a Containing neither granules of free sulphur nor bacteriopurpurin nor requiring the presence of hydrogen sulphid for the best development

1 Not typically producing filaments as a regular growth form, though chains of cells may be developed Conidia not developed, spores when formed are endospores

Order I *Eubacteriales*

2 Typically producing true filaments as a regular growth form Conidia may be developed, but never endospores

(a) Alga-like, typically water forms Filaments never showing true branching, false branching may be

present A sheath usually evident, and usually impregnated with iron

Order II *Chlamydobacteriales*

- (b) Mold like, not typically water forms, not with the sheath impregnated with iron True branching often evident

Order III *Actinomycetales*

- b Cells typically containing either granules of free sulphur or bacteriopurpurin or both, usually growing best in the presence of hydrogen sulphid

Order IV *Thiobacteriales*

- II Cells united during the vegetative stage into a pseudoplasmodium which passes over into a highly developed, cyst-producing, resting stage

Order V *Myxobacteriales*

- B. Protozoan-like in many characters Cells usually relatively slender flexuous spirals, multiplication of cells apparently by longitudinal division in some types, by transverse division in others, or both

Order VI *Spirochaetales*

Key to the families of the Eubacteriales

- A. Organisms usually growing more or less readily upon organic media, not securing growth energy primarily by the oxidation of ammonia or nitrites

I Cells typically spherical Family I *Coccaceae*

II Cells not spherical, elongate

a Cells not spiral Family II *Bacteriaceae*

b Cells spiral, or at least curved

Family III *Spirillaceae*

- B Not growing readily or at all on media containing considerable amounts of organic material, nitrifying bacteria, securing growth energy primarily by the oxidation of ammonia or nitrites

Cells may be either spherical or rod-shaped

Family IV *Nitrobacteriaceae*

Key to tribes of the Coccaceae

- I Not epiphytes, not causing the deposition of iron upon capsules

A Commonly parasitic, in some forms saprophytic, usually thriving well under anaerobic conditions, not commonly producing luxuriant growths on artificial media, many forms failing to grow except upon special media Cells in pairs, chains or irregular masses, never regular packets Gram-positive, with exception of a few strict parasites Acid usually formed in dextrose and lactose broth Pigment white, orange or absent

Tribe I *Streptococcae*

B Saprophytes, or facultative parasites Usually grow best under aerobic conditions, producing abundant to luxuriant growth on artificial media Planes of fission often at right angles Cells aggregated in groups, packets or zoogloal masses Usually Gram-negative Pigment as a rule yellow, red or orange

Tribe II *Micrococcae*

- II. Epiphytes, usually on leaves and stems of water plants. Iron oxid deposited upon the capsule Tribe III. *Siderocapsee*

Key to the genera of the Streptococceae

- A Cells occurring normally in chains
 I. Usually parasitic Not forming zoogloal masses in sugar solutions
 Genus 1 *Streptococcus*
 II. Saprophytic, occurring in cane sugar solutions in zoogloal masses
 Genus 2 *Leuconostoc*
 B Cells not occurring usually or characteristically in chains
 I Parasitic Cells in pairs, growth as a rule meagre No pigment formed.
 a Gram-positive Genus 3 *Diplococcus*
 b Gram-negative Genus 4 *Neisseria*
 II Cells in irregular groups, usually parasites, growth as a rule good, pigment usually orange or white
 Genus 5 *Staphylococcus*

Key to the genera of Micrococceae

- A Cells not in regular packets
 1 Pigment generally yellow Genus 1 *Micrococcus*
 2 Pigment red Genus 2 *Rhodococcus*
 B Cells in regular packets Pigment yellow or orange
 Genus 3 *Sarcina*

Key to the genera of Siderocapsee

- One genus only, *Siderocapsa*

Key to the tribes of the Bacteriaceae

- A Cells not acid-fast
 I Typically producing endospores under favorable conditions.
 Tribe I *Bacilleae*
 II Not producing endospores Tribe II *Bacterieae*
 B Cells acid-fast, frequently showing some tendency to branching
 Tribe III *Mycobacterieae*

Key to the genera of Bacilleae

- A Usually a single endospore formed within each cell
 I Aerobic usually Gram-positive, as a rule liquefying gelatin spores usually not distorting rods when formed
 Genus 1 *Bacillus*
 II Anaërobic or microaerophilic usually
 a Spores produced at extreme tip of cells, forming typical drumsticks Genus 2 *Plectridium*
 b Spores not produced at extreme tip of cells at least not forming drumsticks Cells usually somewhat swollen when spores are formed Genus 3 *Clostridium*

- B Usually a number of spores develop within a swollen cell
 Genus 4 *Metabacterium*

Key to the subgenera of Bacillus

- A Spore not barrel-shaped in longitudinal section and not star-shaped in cross section
 I Motile by means of peritrichous flagella
 Subgenus 1 *Eu-Bacillus*
 II Non-motile
 Subgenus 2 *Bacteridium*
 B Spore barrel-shaped in longitudinal section, longitudinal striations evident
 Subgenus 3 *Astasia*

Key to the subtribes of the Bacterieae

- A Cells usually fusiform Subtribe I *Fusiforminae*
 B Cells not fusiform
 I Requiring serum or hemoglobin for development Obligate parasites
 Gram-negative Non-motile Subtribe II *Hemophilinae*
 II Not requiring serum, or at least hemoglobin, for development Gram-negative or positive Motile or non-motile
 a Obligate aerobes, securing growth energy by oxidation, etc
 Subtribe III *Rhizobinae*
 b Not obligate aerobes, not securing growth energy by oxidation of carbonaceous compounds
 Subtribe IV *Bacteriinae*

Key to the Genera of Fusiforminae

One genus only *Fusiformis*

Key to the Genera of Hemophilinae

- A Requiring serum for growth Cells almost ultra-microscopic Stain best by Giemsa Involution forms characteristic
 Genus 1 *Asterococcus*
 B Requiring hemoglobin for growth Stain readily with ordinary aniline dyes
 Involution forms infrequent Genus 2 *Hemophilus*

Key to the genera of Rhizobinae

- A Not fixing atmospheric nitrogen, securing growth energy usually by the oxidation of ethyl alcohol to acetic acid Genus 1 *Mycoderma*
 B Capable of fixing appreciable amounts of atmospheric nitrogen Grow well on nitrogen-free media
 a Small motile rods, with abundant involution forms, frequently living in root nodules of the higher plants (legumes)
 Genus 2 *Rhizobium*
 b Not symbiotic Cells larger, plump, almost spherical in some cases
 Genus 3 *Azotobacter*

Key to the genera of Bacteriinae

- A Producing usually a yellowish or greenish or fluorescent pigment, usually Gram-negative, motile by means of polar flagella, or non-motile
Genus 1 *Pseudomonas*
- B When pigmented not greenish or fluorescent, when motile with peritrichous flagella
- 1 Cells typically pigmented, chromoparous
- a Producing red or pink pigment
Genus 2 *Serratia*
- b Producing a violet pigment
Genus 3 *Chromobacterium*
2. Cells not typically definitely pigmented, or at least not red or violet
- a Cells typically Gram-negative
- (1) Non-motile, showing bipolar staining commonly Never produce gas from carbohydrates Power of acid production low
Genus 4 *Pasteurella*
- (2) Not showing bipolar staining
- (a) Not producing honey-like growth on potato, branching forms uncommon
- x Gelatin not liquefied or liquefied very slowly Motile or non-motile
Genus 5 *Bacterium*
- vx Gelatin liquefied quickly Motile
Genus 6 *Proteus*
- (b) Producing a honey-like growth on potato Branched cells not uncommon Genus 7 *Pfeifferella*
- b Cells typically Gram-positive All non-motile
- (1) Usually microaerophilic Not typically growing well on the surface of laboratory media Without metachromatic granules
- (a) Non-pathogenic lactic acid bacilli
Genus 8 *Lactobacillus*
- (b) Pathogenic, slender small rods not lactic acid formers
Genus 9 *Erysipelothrix*
- (2) Aerobic rods frequently showing metachromatic granules or irregular staining Genus 10 *Corynebacterium*

Key to the subgenera of Bacterium

- A Organisms which show a maximum of fermentative power including fermentation of lactose, rarely pathogenic, some forms slowly liquefy gelatin
Subgenus 1 *Aërobacter* (or *Eu-Bacterium*)
- B Organisms not showing maximum fermentative power, never producing gas in lactose, frequently pathogenic, never liquefying gelatin
- 1 Producing acid and gas from glucose, sometimes other sugars, but none from lactose
Subgenus 2 *Salmonella*
- 2 Producing gas from none of the carbohydrates, acid sometimes formed
Subgenus 3 *Eberthella*

Key to the Genera of Mycobacteriaceae

One genus only, *Mycobacterium*

Key to Genera of Spirillaceae

A. Cells not larger at center, not tapering

1. Cells usually short, only a segment of a spiral One or rarely two or three
polar flagella Genus 1 *Vibrio*

2. Cells longer, usually with a tuft of polar flagella
Genus 2 *Spirillum*

B. Cells enlarged at center and tapering

Genus 3 *Paraspirillum*

Key to the Genera of Nitrobacteriaceae

A. Cells rod shaped

1. Oxidizing ammonia to nitrous acid Motile

Genus 1 *Nitrosomonas*

2. Oxidizing nitrous acid to nitric acid

Genus 2 *Nitrobacter*

B. Cells spherical

Genus 3 *Nitrosococcus*

Key to the Genera of Chlamydo bacteriaceae

1. Filaments not usually permanently attached

a. Filaments straight, or at least not twisted

Genus 1 *Leptothrix*

b. Filaments twisted

Genus 2 *Didymothrix*

2. Filaments attached

a. Filaments unbranched Genus 3 *Crenothrix*

b. Filaments show pseudodichotomous or false branching

(1) Swarm cells developed (motile conidia) Usually without a
deposit of iron oxid in the sheath

Genus 4 *Sphaerotilus*

(2) Spherical, non-motile conidia Usually with iron oxide

Genus 5 *Clonothrix*

Key to the Genera of Actinomycetaceae

A. No evident aerial threads or conidia formed Usually parasitic Often
anaerobic or microaerophilic

1. Threads usually not branched

a. Threads disjoining very readily, long mycelial threads uncommon

Genus 1 *Actinobacillus*

b. Threads longer, not disjoining into short rods

Genus 2 *Leptotrichia*

2. Threads more or less branched, frequently clubbed in tissues

Genus 3 *Actinomyces*

B. Aerial threads and conidia evident on culture media

Genus 4 *Nocardia*

Key to the Families of Thobacteriales

- A Cells containing sulphur granules (or in one species possibly oxalate crystals), but no bacteriopurpurin
 - 1 Unicellular, motile forms Not filamentous
Family I *Achromatiaceae*
 - 2 Filamentous forms Family II. *Beggiatoaceae*
- B Cells containing bacteriopurpurin with or without sulphur granules
Family III *Rhodobacteriaceae*

Key to the Genera of Achromatiaceae

- A Cells spherical or ellipsoidal
 - 1 Cells ellipsoidal (spherical when newly divided) Cells containing granules of calcium oxalate (perhaps sulfur).
Genus 1 *Achromatium*
 - 2 Cells spherical, with sulphur granules in a central vacuole
Genus 2 *Thiophysa*
- B Cells longer, very large (42 to 86 μ) with peritrichous flagella
Genus 3 *Hillhousia*

Key to the Genera of Beggiatoaceae

- A Filament non-motile, with a contrast to base and tip, attached
Genus 1 *Thiothrix*
- B Filaments motile (oscillating) not attached, no differentiation into tip and base
 - 1 Filaments not in bundles nor surrounded by a gelatinous sheath
Genus 2 *Beggiatoa*
 - 2 Filaments in bundles, surrounded by a gelatinous sheath
Genus 3 *Thioploca*

Key to the Subfamilies of Rhodobacteriaceae

- A Cells containing sulphur granules
Subfamily I *Chromatioideae*
- B Cells without sulphur granules Subfamily II *Rhodobacterioideae*

Key to the Tribes of Chromatioideae

- A Cells united, at least during a part of the life history, into families
 - I Cell division such that masses of cells, not merely plates, are formed
 - a Cell division in three directions of space
Tribe I *Thiocapsee*
 - b Cell division first in three, then in two directions of space.
Tribe II *Lamprocystee*
 - II. Cell division in two planes, forming plates of cells
Tribe III *Thiopediace*
 - III Cell division in one plane Tribe IV *Amoebobacteriace*
- B. Cells free, capable of swarming at any time
Tribe V *Chromatiace*

Key to the Genera of the Thiocapsae

- A Cells capable of swarming
 - I Families small, compact, enclosed singly or several together in a cyst
Genus I *Thiocystis*
 - II Cells large, 7 to 8 μ , loosely bound by gelatin into loose families
Genus II *Thiosphaera*
 - III Cells small, united into solid, spherical families
Genus III *Thiosphaerion*
- B Cells not capable of swarming
 - I Spherical cells spread out upon the substratum in flat families, loosely enveloped in a common gelatin
Genus IV *Thiocapsa*
 - II Arranged in regular packets like *Sarcina*
Genus V *Thiosarcina*

Key to the Genera of Lamprocystae

One genus only *Lamprocystis*

Key to the Genera of Thiopediae

- A Cells occurring regularly in fours Genus I *Lampropedia*
- B Cells occurring in a film or membrane, not regularly disposed in tetrads
Genus II *Thioderma*

Key to the Genera of Amoebobacteriae

- I Cells connected by plasma threads, families amoeboid motile
Genus I *Amoebobacter*
- II Cells not as I
 - A Cells arranged in a net, united by their ends
Genus II *Thiodutylon*
 - B Cells not arranged in a net
 - 1 Capable of swarming Cells loosely aggregated in gelatin
Genus III *Thiotheca*
 - 2 Non-motile Cells closely appressed into a colony
Genus IV *Thiopolycoccus*

Key to the Genera of Chromatiae

- A Cells motile by means of polar flagella Elongated
 - I Cells not spiral
 - a Cell cylindric Genus I *Chromatium*
 - b Cells with tendency to spindle shape
Genus II *Rhabdomonas*
 - II Cells spiral Genus III *Thiospirillum*
- B Cells spherical, or little elongate, non-motile
 - I Cells not encapsulated Genus IV *Rhodocapsa*
 - II Cells encapsulated in pairs Genus V *Rhodotheca*

Key to the Genera of Rhodobacterioidae

- A Cells rod shaped, many embedded in the same slimy capsule
Genus I *Rhodocystis*

- B Cells spherical or short rods
 - 1 In chains, each chain surrounded by a capsule
Genus II *Rhodonostoc*
 - 2 Cells free.. Genus III *Rhodosphaera*
- C Cells free and elongate
 - 1 Cells not bent
 - a Non-motile Genus IV *Rhodobacterium*
 - b Motile. Genus V *Rhodobacillus*
 - 2 Cells bent or curved
 - a Cells short, comma shaped, with single polar flagellum
Genus VI. *Rhodovibrio*
 - b Cells spiral, with polar flagella
Genus VII *Rhodospirillum*

Key to the Genera of Myxobacteriaceae

- I Cells not transformed into coccus-like spores when encysted
 - A Rods forming free cysts in which they remain unmodified Cysts
various sessile or borne on a more or less highly developed cystophore
Genus I *Chondromyces*
 - B Rods forming large rounded cysts one or more, free within a gelatinous
matrix raised above the substratum
Genus II *Polyangium*
- II Rods transformed to form definite more or less encysted, sessile or stalked
masses of coccus-like spores Genus III *Myrococcus*

Key to the Genera of Spirochaetaceae

- I Usually saprophytic, free living in water
 - A Protoplasm spirally wound around an elastic axis filament
Genus I *Spirochaeta*
 - B Not as in (A), cross section circular
Genus II *Spiriospira*
- II Usually parasitic
 - A Possessing a "crest" or ridge Parasitic in mussels
Genus III *Cristispira*
 - B Without a crest Parasitic in warm blooded animals
Genus IV *Treponema*

In September, 1917, appeared the Preliminary Report of the Committee of the Society of American Bacteriologists on Characterization and Classification of Bacterial Types (Winslow, Broadhurst, Buchanan, Krumwiede and Smith) An historical review of systems of classification is given, likewise a review of the points found in the International Botanical Code of principal interests to bacteriologists Four orders of bacteria were recognized The subdivisions of the *Eubacteriales* alone were recommended The classification in outline follows

Suggested Committee Classification of the Class Schizomycetes (1917)

- A Order *Myxobacterales*
- B Order *Thiobacterales*
- C Order *Chlamydobacterales*
- D Order *Eubacterales*
 - I Family *Nitrobacteriaceae*
 - 1 Genus *Hydrogenomonas*
 - 2 Genus *Methanomonas*
 - 3 Genus *Carboxydomonas*
 - 4 Genus *Mycoderma*
 - 5 Genus *Nitrosomonas*
 - 6 Genus *Nitrobacter*
 - 7 Genus *Azotobacter*
 - 8 Genus *Rhizobium*
 - II Family *Mycobacteriaceae*
 - 1 Genus *Actinomyces*
 - 2 Genus *Nocardia*
 - 3 Genus *Mycobacterium*
 - 4 Genus *Corynebacterium*
 - 5 Genus *Fusiformis*
 - 6 Genus *Leptotrichia*
 - III Family *Pseudomonadaceae*
 - 1 Genus *Pseudomonas*
 - IV Family *Spirillaceae*
 - 1 Genus *Vibrio*
 - 2 Genus *Spirillum*
 - V Family *Coccaceae*
 - a Tribe *Streptococceae*
 - 1 Genus *Neisseria*
 - 2 Genus *Streptococcus*
 - 3 Genus *Staphylococcus*
 - 4 Genus *Albococcus*
 - b Tribe *Micrococceae*
 - 1 Genus *Micrococcus*
 - 2 Genus *Sarcina*
 - 3 Genus *Rhodococcus*
 - VI Family *Bacteriaceae*
 - 1 Genus *Bacterium*
 - 2 Genus *Erwinia*
 - 3 Genus *Pasteurella*
 - 4 Genus *Hemophilus*
 - VII Family *Lactobacillaceae*
 - 1 Genus *Lactobacillus*
 - VIII Family *Bacillaceae*
 - 1 Genus *Bacillus*
 - 2 Genus *Clostridium*.

GENERAL SYSTEMATIC BACTERIOLOGY

E Organisms intermediate between bacteria and protozoa

- 1 Genus *Spirochaeta*
- 2 Genus. *Cristispira*
- 3 Genus *Saprospira*.
- 4 Genus *Treponema*.

An artificial key to the orders and genera was also published as follows

1. Artificial Key to the Orders of the Schizomycetes

- a. Cells united during the vegetative stage into a pseudoplasmodium
A *Myxobacterales*
- 2a Cells not forming a pseudoplasmodium
 - b Cells free or united in elongated filaments, often with a well-defined sheath Conidia frequently formed Free sulphur, iron or bacteriopurpurin often present
 - c Cells typically containing granules of sulphur or bacteriopurpurin or both
B *Thiobacterales*
 - 2c Sulphur and bacteriopurpurin absent, iron often present
C *Chlamydobacterales*
 - 2b Cells never in sheathed filaments Conidia only in the mycelial *Mycobacteriaceae* Flagella often present Free iron, sulphur, or bacteriopurpurin never present
D *Eubacterales*

2 Artificial Key to the Families of the Eubacterales

- a Cells spiral with polar flagella
IV *Spirillaceae*
- 2a Not as above
 - b Cells spherical, rarely, if ever, motile, spores never produced, never securing growth energy from nitrogen or ammonia
V *Coccaceae*
 - 2b Not as above
 - c Cells short rod-shaped with a single rarely two) polar flagellum, usually forming green or yellow pigment
III *Pseudo nonadaceae*
 - 2c Not wholly as above
 - d Spores formed
VIII *Bacillaceae*
 - 2d Spores never formed
 - e Metabolism simple, securing growth energy from carbon, hydrogen, or their simple compounds, flagella, if present, polar
I *Nitrobacteriaceae*
 - 2e Metabolism complex dependent upon more complex carbohydrate and protein substances, flagella, if present, peritrichic
 - f Cells clubbed, fusiform, filamentous, branching or mycelial, those not distinctly so are either acid-fast or show barred irregular staining
II *Mycobacteriaceae*

2f Not as above

g Gram-positive, non-motile

VII *Lactobacillaceae*

2g Gram-negative, often motile

VI *Bacteriaceae*

3 Artificial Key to the Genera of the Eubacteriales

a. *Nitrobacteriaceae*

b Fixing nitrogen or oxidizing its compounds

c. Fixing nitrogen

d Cells large, in soil 7 *Azotobacter*

2d Rods minute, in roots of leguminous plants

8 *Rhizobium*

2c Oxidizing nitrogen compounds

d Oxidizing ammonia 5 *Nitrosomonas*

2d Oxidizing nitrites 6 *Nitrobacter*

2b Not as above

c Oxidizing hydrogen 1 *Hydrogenomonas*

2c Oxidizing carbon compounds

d Oxidizing alcohol, branching forms common

4 *Mycoderma*

2d Not as above, using simpler carbon compounds

e Oxidizing CO 3 *Carboxydomonas*

2e Oxidizing CH₄ 2 *Methanomonas*

2a *Mycobacteriaceae*

b Slender rods, staining with difficulty and acid-fast

3 *Mycobacterium*

2b Not as above

c Mycelium and conidia formed

d With aerial hyphae and conidia, usually saprophytic soil organisms 2 *Nocardia*

2d Hyphae and conidia not aerial, usually parasitic in animals

1 *Actinomyces*

2c Not as above, cells rod-like, usually somewhat curved, clubbed, fusiform, or even branched, but never mycelial

d Thick, long threads, fragmenting into short, thick rods

6 *Leptotrichia*

2d Not as above

e Cells usually elongate and fusiform, filaments, if formed, not branching, staining somewhat irregularly

5 *Fusiformis*

2e Cells slightly curved, clubbed, or in old cultures even branching, not filamentous, showing definitely barred staining

4 *Corynebacterium*

3a *Pseudomonadaceae*

Generic characters mainly those of family 1 *Pseudomonas*

4a *Spirillaceae*

b Flagellum single (rarely 2 or 3) 1 *Vibrio*

- 2b Flagella tufted (5-20) 2 *Spirillum*
- 5a *Coccaceae*
- b Abundant red-pigmented growth on agar 7. *Rhodococcus*
- 2b Not as above
- c Gram-negative
- d Normally in pairs of flattened cells, growth on plain agar scanty, never bright yellow 1 *Neisseria*
- 2d Normally in plates, packets, or irregular masses, growth on plain agar abundant, pigment definitely yellow
- e Cells in regular packets 6 *Sarcina*
- 2e Cells not in regular packets 5 *Micrococcus*
- 2c Gram-positive (exceptions rare and not easily confused with above genera)
- d Cells normally in chains, sometimes in pairs (especially in acid environment) never in large irregular masses Gelatine rarely liquefied Growth on plain agar usually translucent, never heavy, never yellow or orange 2 *Streptococcus*
- 2d Cells normally in groups or masses (occasionally in plates in *Albococcus?*), chains short and irregular if present Gelatine often liquefied Agar growth abundant white to orange
- e Pigment orange (rarely lacking), gelatine often liquefied actively 3 *Staphylococcus*
- 2e Whitish to porcelain white, liquefaction less vigorous 4 *Albococcus*
- 6a *Bacteriaceae*
- b Plant pathogens 2 *Eruinia*
- 2b Not as above saprophytes or in animal habitats (intestines tissues, etc.)
- c Usually motile and exhibiting active fermentative powers typically parasitic in intestines of man and higher animals, growing well on ordinary media 1 *Bacterium*
- 2c Not wholly as above
- d Growing only in presence of hemoglobin ascitic fluid or serum 4 *Hemophilus*
- 2d Growth on media scanty but less sensitive than the above short rods with tendency to bipolar stain 3 *Pasteurella*
- 7a *Lactobacillaceae*
- Generic characters mainly those of family 1 *Lactobacillus*
- 8a *Bacillaceae*
- b Aerobic, usually saprophytic, cells not greatly enlarged (if at all) at sporulation 1 *Bacillus*
- 2b Anaerobic, often saprophytic cells frequently enlarged at sporulation 2 *Clostridium*

Brief diagnoses were given of all groups recognized, and type species indicated for the genera

Castellani and Chalmers (Manual of Tropical Medicine, 3rd edition, 1919), published an unusually detailed classification of the bacteria, carrying the classification out only to include the most important pathogenic organisms which have been described. They use as a general name for the group as a whole, *Schizomycetacea*. The keys to the various groups which they recognize are as follows

Key to the orders of Schizomycetacea Castellani and Chalmers (1919)

- A Cells without sulphur or bacterioporpurin
Order I *Eubacteriales*
- B Cells containing sulphur
Order II *Thiobacteriales*
- C Motile rods in pseudoplasmodial masses embedded in a gelatinous matrix and forming highly developed cysts
Order III *Myxobacteriales*

Key to the Families of Eubacteriales

- A Cells, in free condition, usually globular, in division somewhat elliptical
Family 1 *Coccaceae* Zopf, 1885, *emendavit* Migula, 1900
- B Cells, long or short, cylindrical, straight, division one direction
Family 2 *Bacillaceae* Fischer, 1894
- C Cells, spirally curved or representing part of a spiral, division in one direction
Family 3 *Spirillaceae* Migula, 1900
- D Cells, surrounded by a sheath and arranged in elongated filaments
Family 4 *Chlamydobacteriaceae* Migula, 1900
- E Cells, short or long, cylindrical or filamentous, often clavate, cuneate, or irregular, with enclosed granules Filaments without a sheath
Family 5 *Mycobacteriaceae* Chester, 1901

Key to the Tribes of Coccaceae

- A Parasitic on plants and animals, often growing best anaerobically but frequently with difficulty and in small amount, or even not at all, on artificial media, in pairs or chains, generally but not always staining by Gram, and often producing acidity in glucose and lactose media, and when pigmented generally white or orange
Tribe I *Streptococceae* Trevisan, 1889, *emendavit* Winslow and Rogers, 1905
- B Facultative parasites or saprophytes growing best under aerobic conditions and well on artificial media, in cell groups, packets or zoogloea masses and often Gram-negative, and when pigmented usually yellow or red
Tribe II *Micrococceae* Trevisan, 1889, *emendavit* Winslow and Rogers, 1905

Key to the Genera of Streptococceae

- A Gram stain negative *Neisseria*
- B Gram stain positive
 - I Pigment absent
 - a Cells in zoogloea masses, *Ascococcus*

b Not in zoogloea masses

1. Pairs or chains encapsulated, soluble in bile and salt solutions *Diplococcus*
2. Chains, not encapsulated, not soluble in bile or salt solutions *Streptococcus*

II Pigment present

- a Orange, *Aurococcus*
- b White, *Albococcus*

Key to the Genera of the Tribe Micrococceae

- A Pigment red *Rhodococcus*
 B Pigment yellow
 I Cells in irregular groups *Micrococcus*
 II Cells in packets *Sarcina*
 C. Pigment black *Nigrococcus*

Key to the Tribe of the Family Bacillaceae

Growth in ordinary laboratory media

A Entirely or almost entirely absent

Tribe 1 *Nitrobactereae*

B Poor, Gram-negative, grow best on blood media

Tribe 2 *Haemophyleae*

C Extremely slow and scanty growth on ordinary and blood media

Tribe 3 *Gracilordeae*

D Growth good

- I Endospores present Tribe 4
- Bacilleae*

II Endospores absent

(a) Fluorescent or chromogenic

Tribe 5 *Bacterideae*

(b) Neither fluorescent nor chromogenic

- 1 Obligatory anaerobes Tribe 6
- Bacterordeae*

- 2 Aerobes often facultative anaerobes

(1) Gelatine liquefiers

Tribe 7 *Proteae*

(2) Gelatine non-liquefiers

(i) Without capsules

(A) With polar staining

Tribe 8 *Pasteurelleae*

(B) Without polar staining

Tribe 9 *Ebertheae*

(ii) With capsules

Tribe 10 *Encapsulateae**Key to the Tribe Encapsulateae*

This tribe contains one genus only

*Encapsulateae**Key to the Genera of the Tribe Ebertheae*

- A Glucose and lactose either not at all or only partially fermented with the production of acid, but no gas

I Milk not clotted

(a) Glucose and lactose not fermented

Genus 1 *Alcaligenes* Castellani and Chalmers, 1918

(b) Glucose partially fermented with the production of acid and no gas, lactose not fermented

1 Motile Genus 2 *Eberthius* Castellani and Chalmers, 1918

2 Non-motile

Genus 3 *Shigella* Castellani and Chalmers, 1918

(c) Lactose and glucose partially fermented with the production of acid, but no gas

Genus 4 *Dysenteroides* Castellani and Chalmers, 1918

II Milk clotted

Glucose partially fermented with the production of acid, but no gas, lactose not fermented (no gas in any sugar)

Genus 5 *Lankorides* Castellani and Chalmers, 1918

B Glucose completely fermented with the production of acid and gas, lactose not fermented

I Milk not clotted

Genus 6 *Salmonella* Lignières, *emendavit* Castellani and Chalmers, 1918II Milk clotted Genus 7 *Balkanella* Castellani and Chalmers, 1918

C Glucose completely fermented with the production of acid and gas, lactose partially fermented with the production of acid and no gas

I Milk not clotted Genus 8 *Wesenbergus* Castellani and Chalmers, 1918

D Glucose and lactose completely fermented with the production of acid and gas

I Milk not clotted Genus 9 *Enteroides* Castellani and Chalmers, 1918II Milk clotted Genus 10 *Escherichia* Castellani and Chalmers, 1918*Key to the Subgenera of the Genus Shigella*A Mannitol fermented Subgenus *Flemerella* (Flexner group *sensu lato*)I Maltose fermented *Flexner group*II Maltose not fermented *Pseudodysenteriae group*B Mannitol not fermented Subgenus *Shigella**Key to the Tribe Pasteurellae*One genus only *Pasteurella**Key to the Genera of the Tribe Proteae*

A Rapid gelatine liquefiers, do not ferment lactose, mostly Gram-positive

Proteus

B Slow gelatine liquefiers, ferment lactose, Gram-negative

*Cloaca**Key to the Tribe Bacteroidae*One genus only *Bacteroides**Key to the Tribe Graciloidae*One genus only *Graciloides*

Key to the Tribe Bacteroidae

One genus only *Bacteroides*.

Key to the Tribe Bacillae

One genus only *Bacillus*

Key to the Genera of the Family Spirillaceae

A Non-motile, comma-shaped, or spirally curved filaments, rigid, without flagella
Genus 1 *Spirosoma* Migula, 1900

B Motile, short, slightly curved, rigid, comma-like, sometimes in chains, with one, rarely more, flagella at one end, seldom at both ends

Genus 2 *Vibrio* O F Muller, 1773, *emendavit* Loeffler

C Motile, long, spirally curved, usually with a bunch of polar flagella composed of long and short forms

Genus 3 *Spirillum* Ehrenberg 1838 *emendavit* Loeffler

Key to the Family Mycobacteriaceae

One genus only *Mycobacterium*

Many of the Hyphomycetes are included under the class *Fungi-Imperfecti* and under the sub-class *Hyphales*

Key to the Orders of Hyphomycetes

A Mycelium composed of fine bacilliiform hyphae in which the nuclei are usually indistinct
Order 1 *Microsiphonales* Vuillemin 1912

B Mycelium not so composed

1 Reproduction by thallospores

Order 2 *Thallosporales* Vuillemin 1910

2 Reproduction by hemispores

Order 3 *Hemisporales* Vuillemin 1910

3 Reproduction by conidia Order 4 *Coniosporales* Vuillemin 1910

Key to the Families of the Order Microsiphonales

A *Nocardiaceae* Castellani and Chalmers 1918 Synonyms — *Actinomycetes* Lachner-Sandoval 1898 *Trichomycetes* Petrusky 1903

Definition — *Microsiphonales* with a mycelium

Type Genus — *Nocardia* Toni and Trevisan 1889

B *Mycobacteriaceae* Miede, 1909 Definition — *Microsiphonales* without a mycelium

Genus 1 — *Mycobacterium* Lehmann and Neumann with the diphtheria bacillus as a type

Genus 2 — *Corynebacterium* Lehmann and Neumann with the tubercle bacillus as a type

Key to the Family Nocardiaceae

A Grows aerobically easy of cultivation and producing arthrospores (Fig 549)

Genus 1 *Nocardia* Toni and Trevisan 1889

- B Grows best anaerobically, but can often grow aerobically, difficult of culture, and not producing arthrospores Genus 2 *Cohnstieptothrix* Finoy, 1911

Mrs Enlows (1920) prepared a most helpful bulletin in which was listed the various names which have been proposed by various writers for bacterial genera. She has in most cases indicated the type species.

Castellani and Chalmers (1920) proposed a classification of the family *Bacillaceae* Fischer 1894 into some 10 tribes differentiated by the following key

Key to Tribes of Bacillaceae

- A No development in ordinary media *Nitrobacterae*
- B Growth poor in ordinary media, better on media containing blood Gram-negative *Hemophaleae*
- C Very poor and slow growth on ordinary media, also on blood media *Gracilordeae*
- D Good development on ordinary media
 - I Endospores present *Bacilleae*
 - II Endospores absent
 - a Fluorescent or chromogenic *Bacteroidae*
 - b Neither fluorescent nor chromogenic
 - 1 Obligate anaerobes *Bacteroidae*
 - 2 Aerobic, often facultative anaerobic
 - (a) Liquefying gelatin *Proteae*
 - (b) Not liquefying gelatin
 - (1) Without capsules
 - x Bipolar staining *Pasteurellae*
 - 2x No bipolar staining *Ebertheae*
 - (2) With capsule *Encapsulateae*

It will be noted that there is an evident misprint, as both tribes 5 and 6 are termed *Bacteroidae*

Three of the tribes are discussed in detail. The generic keys are given below

Tribe Encapsulateae

The type and only genus is *Encapsulatus* Castellani and Chalmers

Tribe Ebertheae

- A. Either do not ferment glucose and lactose at all, or only partially, with production of acid without gas

I Milk not coagulated

- a Ferments neither glucose nor lactose

Alcaligenes Castellani and Chalmers, 1918

- b Ferments glucose partially, with production of acid but no gas, does not ferment lactose
 - 1 Motile *Eberthus* Castellani and Chalmers, 1918
 - 2 Non-motile *Shigella* Castellani and Chalmers, 1918
- c Ferment glucose and lactose partially with production of acid, but without gas *Dysenteroides* Castellani and Chalmers, 1918
- II Milk coagulated Partial fermentation of glucose with production of acid but no gas does not ferment lactose (no gas from any sugar)
 - Lankoides* Castellani and Chalmers, 1918
- B Ferment glucose completely with production of acid and gas, but do not ferment lactose
 - I Do not coagulate milk
 - Salmonella* Lagnières emendav Castellani and Chalmers
 - II Coagulates milk *Balkanella* Castellani and Chalmers
- C Ferment glucose completely with production of acid and gas, lactose partially fermented with acid and no gas, does not coagulate milk
 - Wesenbergus* Castellani and Chalmers, 1918

In May 1920 the final report of the Committee (Winslow *et al.*, was published. The principal changes from the suggested grouping of 1917 were as follows

The group recognized in 1917 as the family *Mycobacteriaceae* has now been elevated to the rank of an order *Actinomycetales*, and divided into two families, *Actinomycetaceae* and *Mycobacteriaceae*. To the former family we have added the genera *Actinobacillus* and *Erysipelothrix*, and we have omitted *Nocardia*, which Breed (1919) has recently shown should be combined for the present with *Actinomyces*. To the second family we have added the genus *Pfeifferella*.

The *Nitrobacteriaceae* have been divided into two tribes, the *Nitrobactereae* and *Azotobactereae*, and the definition of the family has been modified to permit the inclusion of *Rhizobium* which recent investigations have shown to possess peritrichous flagella, but whose general characteristics ally it clearly with *Azotobacter*. The name *Acetobacter* has been substituted for *Mycoderma* to characterize the vinegar organisms.

Among the *Coccaceae* a new tribe was created for the genus *Neisseria*. The genus *Albococcus* is united with *Staphylococcus* and the new genera *Diplococcus* and *Leuconostoc* are added.

The *Bacteriaceae* are divided into seven tribes *Chromobactereae*, *Eruineae*, *Bactereae*, *Lactobacilleae*, *Pasteurelleae*, *Hemophyleae*, and *Zopfiae*, and the new genera *Erythrobacillus*, *Chromobacterium*, *Zopfius*, and *Proteus* are added. The *Lactobacillaceae*, originally recognized as a distinct family, are thus classed as a tribe of the *Bacteriaceae*.

Artificial Key to the Families and Genera of the Actinomycetales and Eubacteriales Winslow et al. (1920)

- A Typically filamentous forms *Actinomycetaceae*
- B Mycelium and conidia formed *Actinomyces*

- BB No true mycelium
 - C Cells show branching
 - D Gram-negative *Actinobacillus*
 - DD Gram-positive *Erysipelothrix*
 - CC Cells never branch Gram-positive threads later fragmenting into rods *Leptotrichia*
- AA Typically unicellular forms (although chains of cells may occur)
 - B Spherical cells *Coccaceae*
 - C Parasitic forms Cells in pairs, chains or irregular groups, never in packets Generally active fermenters
 - D Cells in flattened coffee-bean-like pairs Gram-negative *Neisseria*
 - DD Cells not as above Gram-positive
 - E Cells in lanceolate pairs or chains Growth on media not abundant
 - F Cells in lanceolate pairs Inulin generally fermented *Diplococcus*
 - FF Cells in chains Inulin generally not fermented *Streptococcus*
 - EE Cells in irregular groups Growth on media fairly vigorous
 - White or orange pigment *Staphylococcus*
 - CC Saprophytic forms Chains occurring in zoogeleal masses in sugar solutions *Leuconostoc*
 - CCC Saprophytic forms Cells in irregular groups or packets, not in chains Fermentative powers low
 - D Packets formed *Sarcina*
 - DD No packets
 - E Yellow pigment *Micrococcus*
 - EE Red pigment *Rhodococcus*
 - BB Rods
 - C Curved rods *Spirillaceae*
 - D Short comma-like rods One to three short flagella *Vibrio*
 - DD Long spirals, five to twenty flagella *Spirillum*
 - CC Straight rods
 - D No endospores
 - E Rods of irregular shape or showing branched or filamentous involution forms
 - F Animal parasites Cells of irregular shape Staining unevenly
 - G Acid fast *Mycobacterium*
 - GG Not acid fast
 - H Cells elongate, fusiform *Fusiformis*
 - HH Cells not fusiform, sometimes branching.

I. Gram-positive. Slender, sometimes clubbed rods.

Corynebacterium

II. Gram-negative. Rods sometimes form threads. Characteristic honey like growth on potato. .

Pfeifferella

FF Not animal parasites Cells staining unevenly and with branched or filamentous forms at certain stages Never acid fast

G Metabolism simple, growth processes involving oxidation of alcohol or fixation of atmospheric nitrogen (latter in symbiosis with green plants).

H Cells minute, symbiotes in roots of leguminous plants

Rhizobium

HH Oxidizing alcohol, branching forms common

Acetobacter

GG. Not as above Proteus-like colonies

H Not attacking carbohydrates Gram +

Zopfius

HH Fermenting glucose and sucrose Gram—

Proteus

EE Regularly formed rods

F Metabolism simple, growth processes involving oxidation of carbon, hydrogen or their simple compounds or the fixation of atmospheric nitrogen

Nitrobacteriaceae

G Fixing nitrogen or oxidizing its compounds

H Fixing nitrogen

Cells large, in soil

Azotobacter

HH Oxidizing nitrogen compounds

I Oxidizing ammonia

Nitrosomonas

II Oxidizing nitrites

Nitrobacter

GG Not as above

H. Oxidizing hydrogen

Hydrogenomonas

HH. Not as above, using simpler carbon compounds

I Oxidizing CO

Carboxydomonas

II. Oxidizing CH₄

Methanomonas

- FF Not as above
 - G Flagella usually present, polar *Pseudomonadaceae Pseudomonas*
 - GG Flagella when present peritrichic *Bacteriaceae*
 - H Parasitic forms showing bipolar staining *Pasteurella*
 - HH Not as above
 - I Strict parasites growing only in presence of hemoglobin or ascitic fluid *Hemophilus*
 - II Not as above
 - J Water forms producing red or violet pigment
 - K Pigment red *Erythrobacillus*
 - KK Pigment violet *Chromobacterium*
 - JJ Not as above
 - K Plant pathogens *Erwinia*
 - KK Not as above
 - L Gram positive, forming large amount of acid from carbohydrates and sometimes CO₂ but no H₂ *Lactobacillus*
 - LL Gram negative, forming H₂ as well as CO₂ if gas is produced *Bacterium*
 - DD Endospores present, *Bacillaceae*
 - E Aerobes *Bacillus*
 - EE Anaerobes *Clostridium*

Orla-Jensen (1921) suggested certain revisions in his classification, introducing the new generic names *Fluormonas*, *Photomonas*, *Protonococcus*, *Microbacterium*, *Colbacterium*, *Aerogenesbacterium*, *Butyriclostridium* and *Putriclostridium*

Hilda Hempl Heller (1921, p 549) suggested that *Bacteria* be recognized as a phylum, and the principal groups be differentiated as follows

Key to the Orders of the Bacteria. Heller (1921)

Simple, one-celled plants that multiply typically by binary fission and occasionally by budding. They show no form of sexual multiplication. They rarely contain cellulose and do not contain chlorophyll or phycocyanin

Phylum 1 *Bacteria*

A *Bacteria* which do not form a complex fruiting body

Class I *Eubacteriae*

Eubacteriae whose cells are never in sheathed filaments. Conidia not observed

Free iron sulphur or bacteriopurpurin never present. Multiplication always occurs by transverse fission

Order I. *Eubacteriales*

Sulphur bacteria

Order II *Thiobacteriales*

Iron and manganese bacteria

Order III *Chlamydothales*

B *Bacteria* which join to form a complex fruiting body.

Class II. *Myxobacteriae*

She also added to the "Committee's" list of families of the *Eubacteriales*, the *Clostridiaceae* with two subfamilies, the *Clostridioideae* and *Putrificoideae*. A key to the genera of the *Clostridioideae* was later (1922, p. 5) published as follows

Key to the Genera of the Subfamily Clostridioideae. Heller (1922)

A Do not produce H_2S demonstrable by lead-acetate-paper test were grown in blood-broth

B Do not liquefy gelatin

C Very large rods that form oval spores and store up starch in carbohydrate media

Genus 1 *Clostridium* Prazmowski
emend Heller. Type species *butyricum* as described by Winogradsky (1895)

CC Rods with spherical spores

D Do not grow on media containing much protein

Genus 2 *Omelianskillus* nov. gen.
Type species *hydrogenicus* as described by Omeliansky (1895 and 1904b)

DD Grow on ordinary anaerobe media

E Sides of the bacilli parallel, spores strictly terminal

Genus 3 *Macintoshillus* nov. gen.
Type species *tetanomorphus* (pseudotetanus bacillus, McIntosh (p. 32), *Bacillus tetanomorphus* Committee (p. 41), as described by the Committee

EE Spores not always strictly terminal sides of bacilli may not be parallel

Genus 4 *Douglasillus* nov. gen. Type species *sphenoides* (*Bacillus sphenoides* Douglas, Fleming and Colebrook), as described by the Committee (p. 43).

- CCC Slender rods with oval endospores, usually Gram-negative
 D Clot milk and attack various sugars, produce much acid
 Genus 5 *Henrillus* nov gen Type
 species *tertius* (*Bacillus tertius* Henry)
 as described by Henry
- DD Do not clot milk, attack few or no sugars, produce little acid
 Genus 6 *Flemingillus* nov gen Type
 species *cochlearius* (*Bacillus cochlearius* Douglas, Fleming and Colebrook) as described by the Committee (p 40)
- CCCC Gram-positive rods which are not markedly slender and which produce oval spores
 D Clot milk, saccharolytic
 E Sporulate meagerly, attack a few sugars, occasionally moderately pathogenic tissue invaders
 Genus 7 *Vallorillus*, nov gen Type
 species *fallax* (*Bacillus fallax* Weinberg and Séguin) as described by the Committee (p 27)
- EE Sporulate readily, attack several sugars, not known to be pathogenic
 Genus 8 *Multifermentans* nov gen
 Type species *tenalbus* (*Bacillus multifermentans tenalbus* Stoddard) as described by Stoddard (1919b)
- DD Do not clot milk ellipsoid spores Large Gram-positive rods with long
 Genus 9 *Hiblerillus* nov gen Type
 species *sextus* (*bacillus* VI of von Hibler) as described by von Hibler (1908) (Résumé by Weinberg and Séguin (p 202)
- BB Liquefy gelatin
 C Produce stormy fermentation of milk and sporulate on alkaline media only
 Genus 10 *Welchillus* nov gen Type
 species *aerogenes* (*Bacillus aerogenes capsulatus* Welch and Nuttall), type 1 as defined by Simonds (1915 a and b)
- CC. Do not produce stormy fermentation of milk
 D Do not sporulate Genus 11 *Stoddardillus* nov gen
 Type species *egens* (*Bacillus egens* Stoddard) as described by Stoddard (1919a)
- DD. Sporulate readily
 E Gram-positive, form woolly colonies in deep agar
 Typically highly pathogenic tissue invaders of many species of animals
 Genus 12 *Rivoltillus* nov gen Type
 species *vibron* (the *vibron septique* of Pasteur) as defined in a future paper

EE Gram-negative, may contain Gram-positive granules.
Form smooth lenticular or modified lenticular colonies
in deep agar Typically pathogenic for cattle, sheep,
and guinea-pigs

Genus 13 *Arloingillus* nov gen. Type
species *Chauvoei* (*Bacterium Chauvoei*
Arloing, Cornevin and Thomas) as
described in a future paper

AA Produce H_2S demonstrable by a lead-acetate-paper test when grown in blood
broth

B Produce a large amount of gas from carbohydrates Heavy Gram-
positive rods with little or no tendency to sporulation

Genus 14 *Meyerillus* nov gen Type
species *sadoua* nov sp to be described
in a future paper

BB Produce less gas from carbohydrates Heavy Gram-positive or Gram-
negative rods that form oval spores that are usually subterminal

Genus 15 *Novillus* nov gen Type
species *maligni* (*Bacillus oedematis*
maligni II Novy) as described by
Novy

A committee of the Society of American Bacteriologists (Bergey, Breed, Hammer, Harrison and Huntoon) have recently (1923) prepared a Manual of Determinative Bacteriology In general, the classification suggested by Winslow *et al* has been followed In some groups genera have been added, and several new families created In the treatment of the *Bacteriaceae* they have followed Castellani and Chalmers and in orders other than *Eubacteriales* they have followed Buchanan The keys for differentiation of orders, families and genera follow

The Classification of Bacteria according to Bergey et al (1923)

Key to the Orders of the Class Schizomycetes

- 1 Simple and undifferentiated forms, the true bacteria
Eubacteriales
- 2 Specialized or differentiated forms
 - a Plant like
 - b Mold like *Actinomycetales*
 - bb Not mold like
 - c Sheathed *Chlamydothales*
 - cc Not sheathed
 - d Sulphur bacteria *Thiobacteriales*
 - dd Slime-mold like *Myxomycetales**
 - aa Protozoan like *Spirochaetales*

* A misprint for *Myxobacteriales*

Key to Families of the Order Eubacteriales

- 1 Organisms obligate aerobes, using oxygen for direct oxidation of carbon, hydrogen, or nitrogen or compounds of these Cells usually rod-shaped, occasionally spherical *Nitrobacteriaceae*
- 2 Not securing growth energy as under I
 - a Cells spherical *Coccaceae*
 - aa Cells not spherical
 - b Cells spiral *Spirillaceae*
 - bb Cells straight rods
 - c Not producing endospores Cells motile or non-motile
Bacteriaceae
 - cc Producing endospores *Bacillaceae*

Key to the Tribes and Genera of the Family Nitrobacteriaceae

- 1 Organisms oxidize simple compounds of carbon and nitrogen *Nitrobacteraceae*
 - a Cells capable of securing growth energy by the oxidation of hydrogen to form water *Hydrogenomonas*
 - b Cells oxidize methane to form CO_2 and water *Methanomonas*
 - c Cells oxidize CO to form CO_2 *Carboxydomonas*
 - d Cells oxidize ammonia to form nitrites *Nitrosomonas*
 - e Cells oxidize nitrites to nitrates *Nitrobacter*
 - f Cells oxidize alcohol to form acetic acid *Acetobacter*
 - g Cells oxidize compounds of sulphur *Thiobacillus*
- 2 Organisms capable of fixing free nitrogen of the air *Azotobacteraceae*
 - a Cells capable of fixing free atmospheric nitrogen when growing in solutions of carbohydrates *Azotobacter*
 - b Cells capable of fixing free nitrogen when growing symbiotically on the roots of *Leguminosae* *Rhizobium*

Key to the Tribes of the Family Coccaceae

- I Strict parasites, failing to grow or growing very poorly on usual artificial media Cells normally in pairs, occasionally in tetrads *Neisseriae*
- II Parasites (except *Leuconostoc*) growing best in media containing serum Occur in pairs or chains *Streptococcae*
- III Facultative parasites or saprophytes Cell aggregates of groups, packets or zoogloeal masses Growth abundant *Micrococcae*

Key to the Genera of the Tribe Neisseriae

One genus only *Neisseria*

Key to the Genera of Tribe Streptococceae

- 1 Parasites growing poorly, or not at all, on artificial media
Cells usually in pairs *Diplococcus*
- 2 Saprophytes, usually growing in cane sugar solutions Cells in pairs or chains
Leuconostoc
- 3 Chiefly parasites Normally forming short or long chains, sometimes pairs,
but never packets *Streptococcus*
- 4 Parasites Cells in groups or short chains, rarely in packets
Staphylococcus

Key to Genera of Tribe Micrococceae

- 1 Facultative parasites or saprophytes Cells in plates or irregular masses
(never in long chains or in packets) *Micrococcus*
- 2 Cell division occurs in three planes forming packets
Sarcina
- 3 Saprophytes Cells in groups or packets. Form red pigment on agar
Rhodococcus

Key to the Genera of Family Spirillaceae

- 1 Cells short, bent rods, rigid, single or united into spirals
Vibrio
- 2 Cells rigid, of various thickness and length and pitch of the spiral forming
either long curves or portions of a turn *Spirillum*

Key to the Tribes of Family Bacteriaceae

- 1 Produce pigment on solid media The pigment may be red yellow violet
green or blue *Chromobactereae*
- 2 Without pigment formation on agar or gelatin *Achromobactereae*
- 3 Cellulose digesting organisms occurring in soil *Cellulomonadeae*
- 4 Plant pathogens growth usually whitish *Eruineae*
- 5 Gram-positive rods growing freely on artificial media Do not attack
carbohydrates *Zoppeae*
- 6 Gram-negative rods growing freely on artificial media (generally act on
carbohydrates with formation of acid and gas
 - a Without capsules *Bacteriae*
 - b Encapsulated *Encapsulateae*
- 8 Rods often long and slender Gram-positive Non-motile Usually pro-
duce lactic acid from carbohydrates *Lactotacillae*
- 9 Anaerobic, non-spore-forming rods *Bacteroidae*
- 10 Gram-negative rods showing bipolar staining Parasitic forms
Pasteurellae
- 11 Minute parasitic forms growing only (or best) in the presence of hemoglobin,
ascitic fluid or other body fluids *Hemophileae*

Key to the Genera of Tribe Chromobactereae

- 1 Small, aerobic rods, producing a red or pink pigment on agar or gelatin
Serratia

- 2 Small, aerobic rods, producing a yellow pigment on gelatin or agar
Flavobacterium
- 3 Small, aerobic rods, producing a violet pigment on solid media
Chromobacterium
- 4 Small, aerobic rods, producing a green or blue-green pigment
Pseudomonas

Key to the Genera of the Tribe Achromobactereae

One genus only *Achromobacter*

Key to the Genera of the Tribe Cellulomonadeae

One genus only *Cellulomonas*

Key to Genera of Tribe Erwiniace

- 1 Motile rods
a Flagella peritrichous *Erwinia*
- 2 Rods motile or non-motile Motile forms possess polar flagella
Phytomonas

Key to the Genera of the Tribe Zopfiae

One genus only *Zopfius*

Key to the Genera of Tribe Bactereae

- A Ferment dextrose with production of acid or acid and gas
- 1 Gas formed from dextrose
- a Gas formed from lactose
- b Acetyl-methyl-carbinol not formed from dextrose *Escherichia*
- bb Acetyl-methyl-carbinol formed *Aerobacter*
- aa Gas not formed from lactose
- b Gas formed from sucrose *Proteus*
- bb Gas not formed from sucrose *Salmonella*
- 2 Gas not formed from dextrose
- a Acid formed from dextrose *Eberthella*
- AA Do not form acid or gas from any of the carbohydrates
Alcaligenes

Key to the Genera of the Tribe Encapsulateae

One genus only *Encapsulatus*

Key to the Genera of the Tribe Lactobacillae

One genus only *Lactobacillus*

Key to the Genera of the Tribe Bacteroidae

One genus only *Bacteroides*

Key to the Genera of the Tribe Pasteurellae

One genus only *Pasteurella*

Key to Genera of Tribe Hemophilae

- | | |
|----------------------|-------------------|
| 1. Aerobic species | <i>Hemophilus</i> |
| 2. Anaerobic species | <i>Dialister</i> |

Key to the Genera of Family Bacillaceae

- | | |
|-------------------------------------|--------------------|
| A Aerobic forms, mostly saprophites | <i>Bacillus</i> |
| B Anaerobic forms Often parasitic | <i>Clostridium</i> |

Key to the Families of the Order Actinomycetales

- | | |
|---|-------------------------|
| A Filamentous forms, often branched, sometimes forming mycelia. Conidia sometimes present Some species parasitic. | <i>Actinomycetaceae</i> |
| B Parasitic forms Rod-shaped, rarely filamentous, and with only slight and occasional branching No conidia | <i>Mycobacteriaceae</i> |

Key to the Genera of Family Actinomycetaceae

- | | |
|---|-----------------------|
| A Parasitic forms | |
| a Aerobic | <i>Actinobacillus</i> |
| b Anaerobic or facultative | <i>Leptotrichia</i> |
| B A few parasitic forms, mostly soil forms | |
| a Filamentous, often branched, sometimes forming mycelia | <i>Actinomyces</i> |
| b Rods forming long filaments which rarely show branching | <i>Erysipelothrix</i> |

Key to the Genera of Family Mycobacteriaceae

- | | |
|--|------------------------|
| A Slender rods acid fast | <i>Mycobacterium</i> |
| B Slender often slightly curved rods not acid-fast | <i>Corynebacterium</i> |
| C Obligate parasites Cells frequently fusiform | <i>Fusiformis</i> |
| D Slender Gram-negative rods | <i>Pfeifferella</i> |

Key to the Families of the Order Chlamydoxales

One family only *Chlamydoxaceae*

Key to the Genera of the Family, Chlamydoxaceae

- | | |
|--|---------------------|
| 1 Filaments usually not permanently attached | |
| a Filaments straight or at least not twisted | <i>Leptothrix</i> |
| b Filaments twisted | <i>Didyminella</i> |
| 2 Filaments attached | |
| a Filaments unbranched | <i>Crenothrix</i> |
| b Filaments show pseudodichotomous branching | |
| 1 Swarm cells developed (motile conidia) Usually without a deposit of iron in the sheath | <i>Sphaerotilus</i> |
| 2 Spherical non-motile conidia Usually with iron oxide | <i>Clonothrix</i> |

Key to the Families of the Thioobacteriales

The keys to the subgroups and genera of this order are the same in general as those of Buchanan (1917-18) *q v*

Key to the Families of the Order Myxobacterales

One family only *Myxobacteriaceae*

Key to the Genera of Family Myxobacteriaceae

- I. Rods transformed into coccus-like spores when encysted, sessile or stalked
masses of coccus-like spores *Myxococcus*
- II Cells not transformed into coccus-like spores when encysted
 - A Rods forming large, rounded cysts, one or more within a gelatinous
matrix raised above the substratum *Polyangium*
 - B Rods forming cysts in which they remain unmodified Cysts various,
sessile or borne on a more or less highly developed cystophore
Chondromyces

Key to the Families of the Order Spirochaetales

One family only *Spirochaetaceae*

Key to the Genera of Family Spirochaetaceae

- A Usually saprophytic, free-living in water
 - 1 Protoplasm spirally wound around an elastic axis filament
Spirochaeta
 - 2 Cross section circular *Saprospira*
- B Usually parasitic
 - 1 Possessing a "crest" or ridge Parasitic in mussels
Cristispira
 - 2 Without crest Parasitic in warm blooded animals and in man
Spirochaeta
Treponema
 - 3 Short, sharply twisted spirals Parasitic in rodents and in man
Leptospira

CHAPTER II

CODES OF NOMENCLATURE AND THEIR APPLICATION IN BACTERIOLOGY

If it be taken as a guiding principle that for each kind of organism there should be one valid name, and that there should be some criterion as to the correctness or validity of any name, it is evident that there must be developed and recognized some code of rules or laws. Such so-called "Codes of Nomenclature" have been developed both in Botany and in Zoology by International Congresses called among other things for this purpose. Inasmuch as the bacteria are plants, or at least most closely allied to the plant kingdom, it would be most natural to follow the botanical code. However, an element of confusion in practice is introduced because bacteriologists have shown a tendency in recent years to term themselves *microbiologists* and their subject *microbiology* to make it evident that their studies are not confined to the bacteria, or even to the plant kingdom. This has been fostered by the discovery of many diseases of man and animals caused by protozoa. In many cases these can be best studied by the methods which have been developed in the bacteriological laboratory. It has proved somewhat confusing to use the botanical code for the plant forms and the zoological code for the protozoa.

A careful study of the botanical and zoological codes will show them to be very similar in most essential characteristics. The botanical code is less rigid, and probably has something more of ambiguity in some of its statements than does the zoological. Logically it would seem that we should follow the botanical code with the bacteria and the zoological code with the protozoa.

The questions at once arise

Are these codes really applicable to the nomenclature of micro-organisms? Have they not been planned with higher groups of plants and animals in mind?

A careful study of the provisions of either code will show that there is no inherent difficulty in application to the lower forms of life. In the botanical code there are, perhaps, one or two rules which the bacteriologists might be reluctant to accept, particularly the rule that all

new species of plants (in our case, bacteria) to be recognized as valid, must be published with a Latin diagnosis. From the standpoint at least of the bacteriologist it would be greatly improved by the introduction of the type concept. Furthermore, in both botanical and zoological codes there is a rule that no two genera (or other groups) can exist with the same name, and that duplicate genera in plant and animal kingdoms should be avoided as far as possible, but that such duplicates are not invalid. In other words, two plants can not have the same name, nor can two animals, but a plant and an animal may be named alike. It would appear that for the sake of the microbiologist there might be appended to each code the rule that in those groups whose position in plant or animal kingdom is in doubt there should exist no duplicate names. There is no reasonable question that the zoological code is the clearer.

In a final determination or formulation of the practice which will best serve the bacteriologist, and particularly in the formulation of a statement to govern practice in the present series, it will be necessary to discuss

a The principles suggested by bacteriologists before the latest revision of the botanical code, particularly the Rules of Nomenclature suggested by Lehmann and Neumann

b The Botanical Code of Vienna as modified at Brussels.

c The Zoological Code

d The suggestions and recommendations of the Society of American Bacteriologists

e Suggested revision of the botanical code formulated by a committee of the Botanical Society of America

A. THE RULES OF BACTERIAL NOMENCLATURE ACCORDING TO LEHMANN AND NEUMANN

Apparently the first bacteriologists to give very serious consideration to rules of nomenclature governing the naming of bacteria and to the formulation of such rules were Lehmann and Neumann. In general it may be said that these rules conformed measurably well to the principles of the botanical code. They undoubtedly constituted a very real advance over any previous statement. They are of sufficient importance to warrant their inclusion. The translation of the rules is given in smaller type. The comments are those of the author.

1 Every plant and, therefore, every *Schizomycete* belongs to a species, every species to a genus, and every genus to a family

2. Following the precedent of Linnaeus, every plant or animal, therefore, every microorganism must have two Latin names the first designating the genus to which the organism in question belongs, this name being a substantive, the second indicating the species, which name is an adjective (not two) or the genitive of a substantive, or rarely a substantive in the nominative case.

Comment. The first two rules are basic, and generally accepted in all codes

3 Genera must only be founded upon important morphologic characteristics, so called "biologic genera," such as *Photobacterium* for all light-emitting bacteria, *Pyobacterium* for rods causing suppuration, etc., are only calculated to produce confusion

Comment. Modern tendency in bacteriology is to disregard this rule in part. Many valid genera are recognized at present in which physiologic as well as morphologic characters are used for purposes of diagnosis and differentiation

4 As a designation for species many authors have used, instead of one adjective or substantive, a plurality of adjectives, evidently with the object of furnishing a description through the name *Bacillus rosettaceus metalloides*, *Staphylococcus pyogenes aureus*, *Bacillus pyogenes foetidus*, *Bacillus mesentericus panis viscosi* I and II This effort can be understood, but it has been abandoned as entirely impracticable by all descriptive naturalists since Linnaeus The name of the species should indicate this unequivocally, the characterization belongs to the description It does no harm if two or more organisms possess names which mean the same, if they do not sound alike In addition to a *Micrococcus albus*, a *Micro niveus*, *Micro albusissimus*, *Micro candicans* and a *Micro purus*, may all be quite valid, we must look to the description to determine the exact differences which exist among these white cocci

5 Names which have been formed improperly i.e. contrary to the binomial rule, may be replaced We have done this with the greatest consideration for the existing name whenever possible We have not changed names like *Bacillus acidilactici*, because *acidum lacticum* represents a single idea, and names like *Sempervivum Reginae Amaliae*, *Pedicularis Frederici Augusti*, *Trigonella Foeniculum Graecum*, *Pedicularis Sceptum carolinum* have remained although certainly not convenient, still uncontested

6 If names are properly formed in the binomial manner and correctly published, then they must not be changed by the author himself, much less by others even if subsequently another appears better Furthermore, the fact that a name is philologically incorrect or not beautiful, is no adequate reason for change Even, for example, if it were literally more correct to call the genus we have named "*Mycobacterium*" by the new name "*Tuberculomyces*" such a proposition is absolutely unallowable Renaming is required only when the name given was employed earlier with another signification Thus, Cohn founded upon a certain organism the new genus *Streptothrix*, without knowing that Corda, about thirty years previously had conferred this name upon a fungus totally different

from the one which he described. His new species, therefore, must receive a new genus name which he who first observed Cohn's oversight is justified in establishing.

7 It may happen that an author differs from his predecessor as to the boundaries of certain genera, and he, therefore, transfers a species from one genus into another, either one which previously existed or which is newly formed by himself. This is permissible, nevertheless the designation of the species must not be changed. So we had the right when we broke up the very large genus *Bacillus*, following the suggestion of Huppe, into the two genera *Bacillus* and *Bacterium*, to rename a number of species (for example, *Bacillus pyocyaneus* renamed *Bacterium pyocyaneum*), but we did not have the right, however much we disliked the species name *pyocyaneum*, to rename the organism *Bacterium coerulesco-viride* or *Bacterium gessardi* or anything else.

8 The author who names the genus places his name after the name given. We speak of *Bacillus* Cohn, and mean the genus *Bacillus* as Cohn established it, of *Vibrio* Ehrenberg emend. Loeffler, and mean the genus *Vibrio* as established by Ehrenberg and afterward more accurately described by Loeffler.

9 Whoever discovers a new species or names one not previously named gives it a genus and a species name, and places his own name after the latter. Flugge, who first named a large number of bacteria, gave, for example, the name *Bacillus pyocyaneus* Flugge, to the long known cause of bluish green suppuration.

10 When one places a species in a new genus he puts his own name after the new name, thus, *Bacterium pyocyaneum* Lehmann and Neumann, but it is always to be recommended that he add in parentheses, the name of the author who first named the species. Therefore we always write, where it does not become too cumbersome (in titles, etc.), *Bacterium pyocyaneum* (Flugge) Lehmann and Neumann.

While we believe that all names which express the systematic position of the organisms shall conform to the general rules of nomenclature, still we believe that names currently used in bacteriologic literature, as gonococcus, pneumococcus, staphylococcus, tubercle bacillus, diphtheria bacillus, can still be used, but as so-called ordinary names. Thus the technical botanist, if not speaking in a strictly systematic sense, often speaks of oak instead of *Quercus*, and strawberry instead of *Fragaria*. We must, however, strictly avoid smuggling into the literature as names of genera, such names as gonococcus, etc.

Comment. Migula in his monumental treatise on bacteria published in 1900, and Chester in his Manual of Determinative Bacteriology published in 1901 have both followed with a measurable degree of success, the rules which have been so clearly summarized in the above statement by Lehmann and Neumann. These rules are, of course, as the authors have indicated, only a condensed summary of international rules with a few interpolations of the authors themselves. In an extensive discussion or consideration of nomenclature, questions are constantly arising which can not be answered by this summary. It is necessary for sufficient and adequate information as to their solution, to turn to the international rules which govern botanical nomenclature.

B. INTERNATIONAL RULES OF BOTANICAL NOMENCLATURE

On the following pages the botanical code (or international rules) as revised by the Brussels International Botanical Congress of 1910, is given in detail. In some cases the discussions given in the original rules under the heading of Examples, has been abbreviated, as many of the examples deal with problems not faced by the bacteriologist. In certain cases rules and recommendations have been omitted for the same reason. A paragraph headed "Comments" is added in many cases to point out certain applications in the field of bacteriology. In some cases comparisons with similar or antagonistic provisions in the zoological code are made. It will be evident as one studies these rules that the botanical code was drafted with the vascular plants in mind and most of the illustrations in consequence have been drawn from this field. However, it has been emphasized previously that a study of the various provisions will show that with but few exceptions they are equally applicable to the conditions which confront the bacteriologist.

Chapter I. General considerations and leading principles

Article 1 Natural history can make no progress without a regular system of nomenclature, which is recognized and used by the great majority of naturalists in all countries.

Comment Apparently about 900 names have been proposed by various authors for genera and higher groups of bacteria up to 1924. Several thousand species have been named. However, it is probable that not one species in ten has been adequately described or named. It is evident that unless an effort is made to follow some system that the same name will be applied to many different kinds of bacteria, and many names to a single kind.

Article 2 The prescriptions which govern the exact system of botanical nomenclature are divided into *principles*, *rules* and *recommendations*. The principles (art. 1-9, 10-14 and 15-18) are the foundation of the rules and recommendations. The rules (art. 10-58), destined to put in order the nomenclature which has bequeathed to us, and to form the basis for the future, are always retroactive; names or forms of nomenclature which are contrary to a rule cannot be maintained. Recommendations bear on secondary points, their object being to ensure for the future a greater uniformity and clearness in nomenclature; names or forms of nomenclature contrary to a recommendation are not a model to copy, but cannot be rejected.

Article 3 The rules of nomenclature should neither be arbitrary nor imposed by authority. They must be simple and founded on considerations clear and forcible enough for everyone to comprehend and be disposed to accept.

Comment It may be emphasized that the International Botanical Congresses which have formulated and revised the International Code have no power to enforce these rules. Stiles (1905, p. 9) states the matter succinctly in his discussion of the zoologist's code. He says

Every person is still in a position to follow any code he desires, or to prepare one of his own, but, considering the responsibilities involved, there is undoubtedly a growing tendency to adopt the International Code. Further, it is considered ethical to ignore names used contrary to this and to other standard codes. While not attempting to dictate to men of science what they shall or shall not do, the commission submits the rules to the serious consideration of all workers in the spirit advanced by Strickland (1842), namely, "we offer them to the candid consideration of zoologists in the hope that they may lead to sufficient uniformity of method in future to rescue science from becoming a mere chaos of words."

Article 4 The essential points in nomenclature are (1) to aim at fixity of names, (2) to avoid or to reject the use of forms and names which may cause error or ambiguity or throw science into confusion.

Next in importance is the avoidance of all useless creation of names.

Other considerations, such as absolute grammatical correctness, regularity or euphony of names, more or less prevailing custom, respect for persons, etc., notwithstanding their undeniable importance are relatively accessory.

Article 5 No custom contrary to rule can be upheld if it leads to confusion or error. When a custom offers no serious inconvenience of this kind, it may be a ground for exceptions which we must, however, abstain from extending or copying. Finally in the absence of rule, or where the consequences of rules are doubtful, established custom becomes law.

Comment This general principle is a useful guide in certain puzzling situations, and is a distinct help toward stability. For example, Article 21 of the code specifies that families are to be designated by the name of one of their genera or ancient generic name, with the ending *-aceae*. The family name *Coccaceae* has been quite generally used in bacteriology during the past two decades. There is, however, no valid, or strictly speaking, any ancient, bacterial genus *Coccus* in bacteriology. Nevertheless it is probably advisable to make this family name the basis of an exception, as such exception would quite certainly cause less confusion than the adoption of an entirely new designation.

Article 6. The principles and forms of nomenclature should be as similar as possible in botany and in zoology, but botanical nomenclature is entirely independent of zoological nomenclature.

Comment The zoological code is somewhat more specific than is the botanical code in dealing with this problem. The corresponding article in the zoological rules reads as follows

Zoological nomenclature is independent of botanical nomenclature in the sense that the name of an animal is not to be rejected simply because it is identical to a name of a plant. If, however, an organism is transferred from the vegetable to the animal kingdom its botanical names are to be accepted in zoological nomenclature with the original botanical status, and, if an organism is transferred from the animal to the vegetable kingdom its names retain their zoological status

The following recommendation is also added "It is well to avoid introducing into zoology as generic names, such names as are in use in botany" These points are of interest to the bacteriologist because until the suggestion was made in 1857 by Naegeli that bacteria are plants and not animals, most of the forms up to that time described which we now classify with the bacteria, had been included with the protozoa This means that such names as *Bacterium*, *Spirillum*, *Spirodiscus*, *Spirochaeta*, and *Vibrio* were included. In the classification of Cohn (1872) these forms were definitely transferred to the plant kingdom They have been used in both zoological and botanical terminology, and, therefore, they prevent subsequent use of these names for other forms among either plants or animals. However, the fact that *Coccus* is the name of a genus of insects would not interfere with its use as a genus in bacteriology Neither does the fact that *Spirillum* was used in 1815 as the name of a worm prevent its acceptance as a valid genus of bacteria Numerous examples might be cited of generic names which are identical in zoology and botany It is well, however, to avoid the introduction of names used as genera of animals into botany when possible This is particularly true in bacteriology, for many of the types have an uncertain position The group of spirochetes, for example, is treated both by bacteriologists and protozoologists To introduce a generic name into this group under the impression that it belongs with plants when such a name has previously been used in zoology, is to cause unnecessary confusion and may very possibly invalidate the name In a few instances organisms have been named as plants and later found to be animals Stiles (1905, p. 10) states

In these cases zoologists accept the botanical names, thus *Plasmodium malariae* was originally (1881) classified in a plant genus as *Oscillaria malariae* Now that this organism is classified as an animal the original specific name *malariae* (1881) is accepted in zoology with its original botanical date The generic name *Oscillaria* is not taken over with the specific name because it was not proposed for this organism *Oscillaria* remains as it was prior to 1881, a plant genus

Article 7 Scientific names are in Latin for all groups When taken from another language, a Latin termination is given them, except in cases sanctioned by custom If translated into a modern language, it is desirable that they should preserve as great a resemblance as possible to the original Latin names

Comments. This rule has been too frequently disregarded by bacteriologists. In many cases this has come about through failure of the one who first described an organism to give to it a specific name, or one in the correct form. For example, Duclaux (1882) described an organism which he found associated with the production of slimy or ropy milk. This he placed in his genus *Actinobacter* but without giving it a specific name. In consequence this organism is commonly noted in literature as *Actinobacter du lait visqueux* Duclaux, an impossible combination. German writers furthermore have frequently changed the spelling of Latin words from their Latin form, as *Bacterium* to *Bakterium*, *Bacillus* to *Bazillus*, *Streptococcus* to *Streptokokkus*, in order to use them as vernacular designations. In many cases after having accepted this changed spelling, the fact is ignored that they have lost their status as generic names and the words are used as though they were in correct Latin form, as in the name *Staphylokokkus pyogenes*. German, French and Italian authors have also shown a tendency to change the names of families and of whole groups, from their correct Latin form, and yet continue to use them in a scientific and not merely a vernacular sense, as *Arthrobakteriaceen* for *Arthrobacteriaceae*, *Kokkaceen* and *Coccacées* for *Coccaceae*.

Article 8. Nomenclature comprises two categories of names (1) Names, or rather terms, which express the nature of the groups comprehended one within the other (2) Names peculiar to each of the groups of plants that observation has made known.

Comments. The nature of the names noted under 1 and 2 will be evident after a perusal of Chapters II and III respectively in which these are discussed.

Article 9. The rules and recommendations of botanical nomenclature apply to all classes of the plant kingdom, reserving special arrangements for fossil plants and non-vascular plants.

Comments. A footnote to the code of 1905 states that the special arrangements noted were to be taken up at the next International Botanical Congress in 1910. These arrangements included "(1) Rules bearing on special points in relation to the nature of fossils or lower plants (2) Lists of *nomina conservanda* for all divisions of plants other than Phanerogams." At this meeting several rules relating primarily to paleobotany and to the algae, fungi, hepatics, lichens and mosses were adopted, but the question of the date of departure and other questions relating to the bacteria and certain other groups were reserved for the International Congress which was to have met in London in 1915.

It is evident that this principle makes possible the recommendation and adoption of special rules to govern groups such as the bacteria. Some such special rules have been suggested to the Society of American Bacteriologists through its committee on bacterial classification.

Chapter II. On the manner of designating the nature and the subordination of the groups which constitute the plant kingdom

Article 10 Every individual plant belongs to a species (*species*), every species to a genus (*genus*), every genus to a family (*familia*), every family to an order (*ordo*), every order to a class (*classis*), every class to a division (*divisio*)

Article 11. In many species we distinguish varieties (*varietas*) and forms (*forma*), and in some cultivated species, modifications still more numerous, in many genera sections (*sectio*), in many families tribes (*tribus*)

Article 12 Finally if circumstances require us to distinguish a greater number of intermediate groups, it is easy, by putting the syllable *sub* before the name of a group, to form subdivisions of that group In this way subfamily (*subfamilia*) designates a group between a family and a tribe, subtribe (*subtribus*) a group between a tribe and a genus, etc The arrangement of subordinate groups may thus be carried, for wild plants only, to twenty-one degrees, in the following order Regnum vegetabile, Divisio, Subdivisio, Classis, Subclassis, Ordo, Subordo, Familia, Subfamilia, Tribus, Subtribus, Genus, Subgenus, Sectio, Subsectio, Species, Subspecies, Varietas, Subvarietas, Forma, Individuum

If this list of groups is insufficient it can be augmented by the intercalation of supplementary groups, so long as these do not introduce confusion or error

Example *Series* and *Subseries* are groups which can be intercalated between subsection and species

Recommendation I. In parasites, particularly in parasitic fungi, authors who do not recognize as species forms which are characterized from a biologic point of view, but little or not at all from the standpoint of morphology may differentiate within species special forms (*forma specialis*, *f. sp.*) characterized by their adaptation to their different hosts

Article 13 The definition of each of these names of groups varies, up to a certain point, according to individual opinion and the state of the science, but their relative order, sanctioned by custom must not be altered No classification is admissible which contains such alterations

Examples of inadmissible alterations are—a form divided into varieties, a species containing genera, a genus containing families or tribes

Article 14 The fertilization of one species by another gives rise to a hybrid (*hybrida*), that of a modification or subdivision of a species by another modification to the same species gives rise to a half-breed (*mustus*, *mule* of florists)

Chapter III. On the manner of designating each group or association of plants

Section 1 General principles, priority

Article 15. Each natural group of plants can bear in science only one valid designation, namely, the oldest, provided that it is in conformity with the rules of Nomenclature and the conditions laid down in articles 19 and 20 of section 2

Comment This is the essence of the whole matter. No matter what our individual preferences may be, or how much better another more familiar name may sound, we should choose the oldest name if it is valid. In no other way can nomenclature be stabilized. Many cases in bacteriological literature may be cited in which authors have created new names needlessly, in some cases because the names seemed inappropriate for some reason.

Article 16. The designation of a group by one or several names is not for the purpose of describing the characters or the history of the group, but that we may be understood when we wish to speak of it.

Comment Misunderstanding of this principle has led to many mistakes in literature. There is no good reason why the specific name of an organism must be descriptive of the organism in any way. Some authors have contended that the naming of organisms should in some manner conform to the principles which have been followed in the naming of compounds in the science of chemistry. Others have insisted upon the use of descriptive Latin phrases, such as those which have been approved in anatomy and pathology. As has often been pointed out in taxonomic discussions, there is no more reason why a specific name should be descriptive of the organisms than that the given or Christian name of an individual should indicate his characteristics.

Article 17. No one should change a name or a combination of names without serious motives, based on a more profound knowledge of facts, or on the necessity of giving up a nomenclature that is contrary to rules.

Comment Unfortunately there have been a few botanists, perhaps even some bacteriologists who have coined new names needlessly, evidently with the thought that such could in some way be of personal advantage. It is evident from the statement of this rule that when new names are coined or old names changed, an adequate reason for such new names or such new terms should be given.

Article 18. The form, number and arrangement of names depend on the nature of each group, according to the following rules.

Section 2 Point of departure for nomenclature, limitation of principle of priority

Article 19. Botanical nomenclature begins for the various groups of plants (both living and fossil) with the following dates:

- a *Phanerogamae* and *Pteridophyta*, 1753 (Linné, *Species plantarum*, ed. 1)
- b *Muscineae*, 1801 (Hedwig, *Species Muscorum*)

- c *Hepaticae* and *Sphagnaceae*, 1753 (Linné, *Species plantarum* ed 1)
 - d *Lichenes*, 1753 (Linné, *Species plantarum*, ed 1)
 - e *Fungi Uredinales*, *Ustilaginales* and *Gasteromycetes*, 1801 (Persoon, *Synopsis Methodica Fungorum*)
 - f *Fungi caeteri*, 1821-32 (Fries, *Systema Mycologicum*)
 - g *Algae*, 1753 (Linné, *Species plantarum* ed 1)
- Exceptions *Nostocaceae homocystee*, 1892-93 (Gomont, *Nostocaceae homocystee*), *Nostocaceae heterocystee*, 1886 (Bornet and Flahault, *Nostocaceae heterocystee*), *Desmidiaceae*, 1848 (Ralfs, *British Desmidiaceae*), *Oedogomaceae*, 1900 (Hirn, *Monographie und Iconographie der Oedogoniaceen*)
- h *Myxomycetes*, 1753 (Linné, *Genera plantarum*, ed. 1)

It is agreed to associate genera, the names of which appear in the first edition of *Species Plantarum* of Linné with the descriptions which he has given in his *Genera Plantarum*, ed 5 (1754)

Comment A footnote to this article in the Proceedings of the Brussels Congress in 1910 states, "The points of departure for the nomenclature of the following groups *Schizomycetes* (Bacteria), *Schizophyceae* (excl *Nostocaceae*), *Flagellatae* (incl. *Dinoflagellatae*) and *Bacillariaceae* (*Diatomaceae*) are reserved for the Congress of London in 1915" This Congress was postponed because of the interruption by the European War There is therefore no authoritative starting point for bacterial nomenclature at the present time. At this Congress in Brussels Vuillemin proposed that 1753 be adopted as the point of departure Klebahn suggested that the work of F Cohn (1870-76, *Untersuchungen uber Bakterien*) constituted a more suitable point of departure This was approved by Engler Magnus contended that much work of systematic value had been accomplished before 1870 and proposed that the publications of Ehrenberg (certainly a misprint) in 1786 should be the starting point Vuillemin contended that if a date subsequent to 1753 be taken, it might well be 1910 The whole matter was finally deferred to the next Congress as noted above

In preparation for this Congress Vuillemin (1913) published a paper in which he discussed bacterial classification and nomenclature He concluded that the best thing to do is to determine which generic names are valid, prepare such a list, publish the names as *genera conservanda* and date all bacterial classification from 1915 when the list was to have been adopted by the Congress. A study of Vuillemin's paper despite his criticism of bacteriologists as taxonomists shows that he himself does not formulate tenable bases for differentiation of genera and his reason for choosing certain generic names and abandoning others will scarcely withstand critical analysis

The committee on classification of the Society of American Bacteriologists has suggested that the publication of the third edition of Zopf's "Die Spaltpilze" in 1885 be made the point of departure. Unexpected difficulties in the utilization of 1885 as the point of departure have arisen. It seems probable that the least confusion will eventually develop if 1753 be regarded as the date of departure.

Article 20. However, to avoid disadvantageous changes in the nomenclature of genera by the strict application of the rules of nomenclature, and especially of the principle of priority in starting from the dates given in Article 19, the rules provide for lists of names which must be retained in all cases. These names are by preference those which have come into general use in the fifty years following their publication, or which have been used in monographs and important floristic works up to the year 1890. For the use of paleobotanists two lists are provided: (1) A list of generic names validly published and generally accepted, although they come into conflict with paleobotanic names which are older. (2) A list of generic names of fossil plants, validly published and generally accepted when they come into conflict with older homonyms of living plants which have been discarded into synonymy. In order to avoid using these names anew, these lists are given in an appendix to the rules of nomenclature.

Comments. Two lists of generic names have been proposed for eventual submission to an International Botanical Congress for use in bacteriology. The first of these was published by Vuillemin (1913) and is as follows: *Planococcus*, *Streptococcus*, *Klebsiella*, *Merista*, *Planomerista*, *Neisseria*, *Sarcina*, *Planosarcina*, *Metabacterium*, *Clostridium*, *Serratia*, *Bacterium* and *Spirillum*.

In 1916 the following list of genera was proposed to the Society of American Bacteriologists by its committee on classification. These names were to be considered for two years before a vote should be taken upon them. The list is as follows: *Methanomonas*, *Carboxydomonas*, *Hydrogenomonas*, *Mycoderma*, *Nitrosomonas*, *Nitrobacter*, *Azotobacter*, *Rhizobium*, *Nocardia*, *Actinomyces*, *Mycobacterium*, *Corynebacterium*, *Fusiformis*, *Leptotrichia*, *Pseudomonas*, *Vibrio*, *Spirillum*, *Neisseria*, *Streptococcus*, *Staphylococcus*, *Albococcus*, *Micrococcus*, *Sarcina*, *Rhodococcus*, *Bacterium*, *Erwinia*, *Pasteurella*, *Hemophilus*, *Lactobacillus*, *Bacillus*, *Clostridium*. More recent study shows that probably this list should be revised.

Section 3. Nomenclature of the different kinds of groups

Subsection 1. Names of groups above the family

Recommendations. The following suggestions as to the nomenclature of groups of higher rank than the family will tend to clearness and uniformity.

II. Names of divisions and subdivisions, of classes and subclasses are taken from one of their chief characters. They are expressed by words of Greek or Latin origin, some similarity of form and termination being given to those that designate groups of the same nature.

III. Orders are designated preferably by the name of one of their principal families, with the ending *-ales*. Suborders are designated in a similar manner, with the ending *-ineae*. But other terminations may be retained for these names, provided that they do not lead to confusion or error.

Comment The generally accepted name for the entire group of bacteria is *Schizomycetes*. This may be regarded as a class. The preferable designation for the orders of bacteria would have the ending *-ales*. The committee on classification of the Society of American Bacteriologists has suggested the following four orders: *Myxobacterales*, *Thiobacterales*, *Chlamydobacterales* and *Eubacterales*.

Subsection 2. Names of families and subfamilies, tribes and subtribes.

Article 21. Families (*familiae*) are designated by the name of one of their genera or ancient generic names with the ending *-aceae*.

Examples: *Rosaceae* (from *Rosa*), *Salicaceae* (from *Salix*), *Caryophyllaceae* (from *Dianthus Caryophyllus*), etc.

Comment Among the family names of bacteria which have been proposed and which are in correct form are *Spirillaceae* from *Spirillum*, *Bacteriaceae* from *Bacterium*, *Bacillaceae* from *Bacillus*.

Examples of invalid families in bacteriology are the following: *Thiobacteriaceae*, because there is no genus *Thiobacterium*, *Thiorhodaceae* because there is no genus *Thiorhodum*, *Trichobacteria*, both because it lacks the ending *-aceae* and because it contains no genus *Trichobacterium*.

Article 22. The following names, owing to long usage, are an exception to the rule: *Palmae*, *Gramineae*, *Cruciferae*, *Leguminosae*, *Guttiferae*, *Umbelliferae*, *Labratae*, *Compositae*.

Article 23. Names of subfamilies (*subfamiliae*) are taken from the name of one of the genera in the group, with the ending *-oideae*. The same holds for the tribes (*tribus*) with the ending *-eae*, and for the subtribes (*subtribus*) with the ending *-inae*.

Comment The following subfamilies of bacteria are in proper form: *Chromatioideae* from the genus *Chromatium*, *Rhodobacterioideae* from the genus *Rhodobacterium*. The following are examples of names of subfamilies incorrectly formed: *Paracoccaceae* and *Metacoccaceae* because there are no bacterial genera *Paracoccus* and *Metacoccus*, and also because of an incorrect ending. An example of a tribal name would be *Streptococceae* containing the genus *Streptococcus*, of a subtribe, *Bacillinae* containing the genus *Bacillus*.

Subsection 3 Names of genera and divisions of genera

Article 24 Genera receive names, substantives (or adjectives used as substantives) in the singular number and written with a capital letter, which may be compared with our own family names. These names may be taken from any source whatever and may even be composed in an absolutely arbitrary manner.

Examples *Rosa*, *Convolvulus*, *Hedysarum*, *Bartramia*, *Liquidambar*, *Gloriosa*, *Impatiens*, *Manihot*

Comments. Examples of bacterial genera are *Bacillus*, *Actinomyces* and *Vibrio*.

Not infrequently, particularly in medical writers, one finds a decided tendency to ignore the rule that a genus name must, when used as such, always be written with a capital letter. Many of our medical journals and even some which publish scientific articles on medical bacteriology write *bacillus typhosus* or *bacillus tuberculosis*. It is true that sometimes a word with the same spelling as a generic name may be used as a morphologic designation. It cannot be considered incorrect to speak of *tubercle bacillus*, for this is a common name or vernacular designation. As Stiles (1905, p. 16) says (this same rule holds in zoological nomenclature)

This custom of capitalizing the generic name has come down to us from former centuries, and while it now has its practical basis, its historic basis is probably due to the custom earlier authors had of capitalizing all Latin substantives when used in nomenclature. At present the custom is so thoroughly established that botanists and zoologists experience much the same sensation in seeing "*bacillus tuberculosis*" or "*plasmodium malariae*," instead of *Bacillus tuberculosis* and *Plasmodium malariae*, that they would if they found a reference to "american" authors or to "doctor s. weir mitchell."

It is also considered good form to write names of genera, particularly when associated with a specific name, in *italics*. Some of our medical journals have recently abandoned this practice under the plea that it is an unnecessary expense and that italics should be used only for the sake of emphasis. This is unfortunate as it has in the past served as a satisfactory method to indicate clearly in a text just when names were being used in a strict scientific sense.

Some purists have taken exception to the statement that these names may be taken from any source whatever and may even be composed in an absolutely arbitrary manner. Clements (1902) in his paper on Greek and Latin in biological nomenclature states that vernacular names should be invalid, and that a name or term should be regarded as invalid unless constructed according to the principles of word formation in

classic Greek or Latin, and all terms improperly constructed should be corrected. He further states that anagrams and geographic names should be regarded as invalid for genera.

Below under the heading of "Recommendations" there are many suggestions of the best form to follow in the creation of generic names, but failure to follow such recommendations does not render a term invalid.

Article 25. Subgenera and sections also receive names, usually substantives and resembling the names of genera. Names of subsections and other lower subdivisions of genera are preferably adjectives in the plural number and written with a capital letter, or their place may be taken by an ordinal number or a letter.

Examples Substantives *Fraxinaster*, *Trifolium*, *Adenosella*, *Eubermannia*, *Archieracium*, *Micromelilotus*, *Pseudinga*, *Heterodraba*, *Gymnocimum*, *Neoplantago*, *Stachyotypus*. Adjectives *Pleostylae*, *Fimbriati*, *Bibracielata*, *Pachycladae*.

Comments Among names that have been applied to subgenera of bacteria are *Eusarcina* and *Euspirillum*.

Recommendations IV When the name of a genus, subgenus or section is taken from the name of a person, it is formed in the following manner:

a When the name ends in a vowel, the letter *a* is added (for example *Glazioua* *Glaziou*, *Bureaua* after *Bureau*), except when the name already ends in *a*, in which case *ea* is added (e.g., *Collaea* after *Colla*).

b When the name ends in a consonant, the letters *ia* are added (thus *Magnusia* after *Magnus*, *Ramondia* after *Ramond*), except when the name ends in *er*, in which case *a* is added (e.g., *Kerneia* after *Kerner*).

c The spelling of the syllables unaffected by these finals is retained, even with the consonants *h* and *w* or with groupings of vowels which were not used in classic Latin. Letters which are unknown to botanical Latin must be transcribed, diacritic signs are suppressed. The German *a*, *o*, *u*, become *ae*, *oe*, *ue*, the French, *é*, *è* and *ê* become generally *e*.

d Names may be accompanied by a prefix, or a suffix, or modified by anagram or abbreviation. In these cases they count as different words from the original name. E.g., *Durvillea* and *Urvillea*, *Lapeyrousea* and *Peyrousea*, *Englera*, *Englerastrum* and *Englerella*, *Bouchea* and *Ubochea*, *Gerardia* and *Graderia*, *Martia* and *Martusia*.

Comments Among the generic names of bacteria which have been derived from patronymics are *Pasteurella*, *Salmonella*, *Klebsiella*, *Detoniella*, *Zopfella*, *Mantegazzaea*, *Erwinia*, *Pacima*, *Corneha*, *Newskia*, *Pollendera*, *Macintoshillus*, *Metchnikovillus*, *Zopfius*. It will be noted that authors have frequently given generic names among the bacteria by adding *-ella* in accordance with above. The custom is so well established that there would seem to be little good reason at the present time

for opposing it There have been objections, however, raised by those who contend for the use of classic Latin names for all plants and animals Clements (1902, p 82) says "A single generic name may be founded upon the name of a botanist Such names are only to be formed by adding *-ia* to cognomina ending in a consonant and *-a* to cognomina in a vowel or in *-r* except in the case of names already latinized in which case the termination is first dropped " Apparently Clements possessed a high estimate of the ability of classic Latin and Greek to furnish the tens of thousands of generic names that will still be demanded before all plants and animals have been named and classified The formation of names by anagram is also objectionable to such purists, as witness the following statement from Clements (1902, p 78)

Anagrams if they be considered words at all are vernacular since they are neither Greek or Latin They are the ultimate product of puerility or illiteracy in nomenclature Such a series as *Filago*, *Gifola*, *Ifloga*, *Logfia* and *Oglifa* throws a clear light upon the good sense and linguistic tests of the authors concerned One might better make names after the fashion of Carroll or take names from the "dog latin" of childhood

Views in contradiction to this are generally held, however, by most practical workers in the fields of zoology and botany These workers have a firm conviction that botanical Latin does not need to be classic Latin This is well stated in the following quotation from Stiles (1905, p 12)·

Nothing is stated in the rule as to what particular Latin should be used, but latitude is granted to use Latin or Latinized names, or to use names as if they were Latin This latitude has its practical basis The Latin poets and authors did not foresee the practical scientific use to which their language would be applied, and they failed to leave enough Latin words as names for the millions of animals and plants which come into consideration Hence as a carpenter, a sailor or a manufacturer coins words which are accepted in English as English words of good standing, when applied to carpentry, nautical affairs, and trade, the zoologists and botanists coin words upon Latin precedents, which we adopt as botanical and zoological Latin Not in all cases do our efforts attain the highest standards of classical Latin (*Taenia solium*, and *Gadus tomcod*, for instance), but because of practical considerations such names are retained, although not recommended, as examples to be followed in coining new words

V Botanists who are publishing generic names show judgment and taste by attending to the following recommendations

a Not to make names very long or difficult to pronounce

Comments Clements believes that a generic name should be invalid if it contains more than six syllables Apparently few generic names

have been applied to bacteria which do not conform to this standard, though there are many with six syllables such as *Botulobacillus*, *Actinobacillus*, *Actinocladothrix*, *Aphaneroglia* and *Arthrobacterium*. Some offenders are the following *Actinobacterium*, *Caseobacterium*, *Chromobeggiatoa*, *Corynebacterium*, *Liquidobacterium* and *Nitrosobacterium*. These long names are, of course, valid.

b. Not to use again a name which has already been used and has lapsed into synonymy (homonymy)

Comments. The zoological code goes farther than this and forbids the subsequent use for another group of any name which has previously been used. The rule reads "A generic name is to be rejected as a homonym when it has previously been used for some other genus of animals."

c. Not to dedicate genera to persons who are in all respects strangers to botany, or at least to natural science, nor to persons quite unknown.

d. Not to take names from barbarous tongues, unless those names are frequently quoted in books of travel, and have an agreeable form that is readily adapted to the Latin tongue and to the tongues of civilized countries.

e. To recall, if possible, by the formation or ending of the name, the affinities or the analogies of the genus.

f. To avoid adjectives used as nouns.

g. Not to give a genus a name whose form is rather that of a subgenus or section (e.g., *Eusideroxylon*, a name given to a genus of *Lauraceae*, which, however, being valid, cannot be changed).

Comment. The *Eubacillus* of Dangeard is a similar example from bacteriology.

h. Not to make names by the combination of two languages (*nomina hybrida*)

Comments. This is a common fault among names which have been applied in bacteriology. Among such may be noted *Acetobacter*, *Caseobacterium*, *Coccobacillus*.

VI. Botanists constructing names for subgenera or sections, will do well to attend to the preceding recommendations and also to the following.

a. Give, where possible, to the principal division of a genus, a name which, by some modification or addition, calls the genus to mind (for instance, *Eu* placed at the beginning of the name, when it is of Greek origin, *-astrum*, *-ella* at the end of the name, when Latin, or any other modification consistent with the grammar and usages of the Latin language).

b. Avoid calling a subgenus or a section by the name of the genus to which it belongs, with the final *-oides* or *-opsis*, on the contrary reserve this ending for a

section which resembles another genus, by adding in that case *-oides* or *-opsis* to the name of that other genus, if it is of Greek origin, to form the name of the section

c Avoid taking as the name of a subgenus or section a name which is already in use as such in another genus, or which is the name of an admitted genus

Comment It may be noted that in zoology identical rules are used for the formation of genera and subgenera, because the subgenus of today may be the genus of tomorrow and, from the standpoint of nomenclature, both generic and subgeneric names have the same status. Unfortunately, perhaps, botanists have not seen fit to take this step

Recommendation VII. When it is required to express a subgeneric or sectional name together with the name of the genus and the name of the species, the name of the section is put between the others in a parenthesis. Eg, *Astragalus (Cycloglothis) contortuplicatus*

Subsection 4 Names of species and of subdivisions of species

Article 26 All species, even those that singly constitute a genus, are designated by the name of the genus to which they belong followed by a name (or epithet) termed specific, usually of the nature of an adjective (forming a combination of two names, a binomial, or binary name)

Recommendations VIII The specific name should, in general, give some indication of the appearance, the characters, the origin, the history or the properties of the species. If taken from the name of a person, it usually recalls the name of the one who discovered or described it, or was in some way concerned with it

Comment Apparently this recommendation is sometimes taken to indicate that the specific name should constitute an adequate diagnosis of the organism in question

IX. Names of men and women and also names of countries and localities used as specific names, may be substantives in the genitive (*Clusii, saharae*) or adjectives (*Clusianus, dahuricus*). It will be well, in the future, to avoid the use of the genitive and the adjectival form of the same name to designate two different species of the same genus, for example *Lysimachia Hemsleyana* Maxim (1891), and *L. Hemsleyi* Franch (1895)

X. Specific names begin with a small letter except those which are taken from names of persons (substantives or adjectives) or those which are taken from generic names (substantives or adjectives)

XI. When a specific name is taken from the name of a man, it is formed in the following way—

a. When the name ends in a vowel, the letter *i* is added (thus *Glazouii* from *Glazou*, *Bureauii* from *Bureau*), except when the name ends in *a*, when *e* is added (thus *Balansae* from *Balansa*)

b When the name ends in a consonant, the letters *ii* are added (thus *Magnusi* from *Magnus Ramondii* from *Ramond*), except when the word ends in *er* when *i* is added (ex *Kerneri*, from *Kerner*)

c Syllables which are not modified by these endings retain their original spelling, even in the case of the consonants *k* and *w* or groupings of vowels which are not used in classic Latin Letters foreign to the Latin of botanists should be transcribed, and diacritic signs suppressed The German *a*, *ö*, *u*, become *ae*, *oe*, *ue*, the French *é*, *è* and *ê* become, in general, *e*

d. When specific names taken from the name of a person have an adjectival form a similar plan is adopted (*Geranium Robertianum*, *Carex Hallerana*, *Ranunculus Bureauanus*, etc)

XII The same applies to the names of women These are written in the feminine when they have a substantival form

Example *Cypripedium Hookerae*, *Rosa Beatricis*, *Scabiosa Olga*, *Omphalodes Lucilae*.

XIII In the formation of specific names composed of two or several roots and taken from Latin or Greek, the vowel placed between the two roots becomes a connecting vowel, in Latin *i*, in Greek *o*, thus we write *menthaefolia*, *salviaefolia*, not *menthaefolia*, *salviaefolia* When the second root begins with a vowel and euphony demands, the connecting vowel is eliminated (e g , *calthantha*, *lepidantha*) The connecting *ae* is legitimate only when etymology demands (e g , *caricaeformis* from *Carica*, may be retained along with *cariciformis* from *Carex*)

XIV In forming specific names, botanists will do well to note the following recommendations

a Avoid very long names and those which are difficult to pronounce

Comments A few specific names which have been applied to bacteria seem to transcend good usage, but this of course does not invalidate them Among such long specific names which have been applied may be noted the following *saccharobutyricus*, *heminecrobiphilus*, *diphtheriae-columborum*, *viridi-pallescens*, *endoparagocicum*, *melanoglassophorus*

b Avoid names which express a character common to all or nearly all the species of a genus

Comment Among such names are the following *Streptococcus sphericus* and *Albococcus albus*. These names are, of course, valid

c Avoid names taken from little known or very restricted localities, unless the species be very local

Comment. *Azotobacter vinelandii* named after a locality in New Jersey is scarcely in conformity with this recommendation, but is of course, not invalidated by this fact.

d. Avoid, in the same genus, names which are very much alike, especially those which differ only in their last letters

e. Adopt unpublished names found in travellers' notes and herbaria, attributing them to the authors concerned, only when those concerned have approved the publication

f. Avoid names which have been used before in the genus, or in any closely allied genus, and which have lapsed into synonymy (homonyms).

Comment The Zoological Code reads: "A specific name is to be rejected as a homonym when it has previously been used for some other species of the same genus" This recommendation should as far as practicable be given the force of a rule

g. Do not name a species after a person who has neither discovered, nor described, nor figured, nor in any way studied it

h. Avoid specific names formed of two words

Comment *Bacillus acidilactici* is probably valid inasmuch as the specific name refers to a single concept, but it is not a form to pattern after This, of course, does not validate such a name as *Bacillus coli communis*, which is a trinomial

Article 27 Two species of the same genus cannot bear the same specific name, but the same specific name may be given in several genera

Example *Arabis spathulata* DC and *Lepidium spathulatum* Phil are valid as two names of Crucifers, but *Arabis spathulata* Nutt in Torr and Gray cannot be maintained, on account of the existence of *Arabis spathulata* DC, a name previously given to another valid species of *Arabis*

Article 28 Names of subspecies and varieties are formed like specific names and follow them in order, beginning with those of the highest rank The same holds for subvarieties, forms, and slight or transient modifications of wild plants which receive a name or numbers or letters to facilitate their arrangement Use of a binary nomenclature for subdivisions of species is not admissible

Examples *Andropogon ternatus* subsp *macrothrix* (not *Andropogon macrothrix* or *Andropogon ternatus* subsp *A. macrothrix*), *Hernaria hirsuta* var *diandra* (not *Hernaria diandra* or *Hernaria hirsuta* var *H. diandra*), forma *nanus*, forma *maculatum*

Recommendation. XV. Recommendations made for specific names apply equally to names of subdivisions of species These agree with the generic name when they have an adjectival form (*Thymus Serpyllum* var *angustifolius*, *Ranunculus acris* subsp *Friesianus*)

Special forms are named by preference by means of the names of the host species In making such names one may employ double names *Examples* *Puccinia Hieraci* f sp *villosi*, *Pucciniastrum Epilobi* f sp *Abietis-Chamaenerii*

Article 29 Two subspecies of the same species cannot have the same name A given name can only be used once for a variety of a given species, even when dealing with varieties which are classed under different subspecies The same holds for subvarieties and forms

On the other hand the same name may be employed for subdivisions of different species, and the subdivisions of any one species may bear the same name as other species

Examples: The following are admissible: *Rosa Jundzillii* var. *leioclada* and *Rosa glutinosa* var. *leioclada*; *Viola tricolor* var. *hirta*, in spite of the existence already of a different species named *Viola hirta*. The following are incorrect: *Erysimum hieracifolium* subsp. *strictum* var. *longisiliquum* and *E. hieracifolium* subsp. *pannonicum* var. *longisiliquum*—a form of nomenclature which allows two varieties bearing the same name in the same species.

Recommendation. XVI. Botanists are recommended to use as little as possible the privilege granted in the second part of article 29, in order to avoid confusion and mistakes and also to reduce to a minimum the necessary changes of name when the subdivisions of species are raised to specific rank or vice versa.

Article 30. Forms and half-breeds among cultivated plants should receive fancy names, in common language, as different as possible from the Latin names of the species or varieties. When they can be traced back to a species, a subspecies or a botanical variety this is indicated by a succession of names.

Example: *Pelargonium zonale* Mrs. Pollock.

Subsection 5. Names of hybrids and half-breeds (mules)

This subsection contains four Articles and one Recommendation. It is not included here as it deals with problems which are not met in bacterial nomenclature.

Section 4. The publication of names and of the date of each name or combination of names

Article 35. Publication is effected by the sale or public distribution of printed matter or indelible autographs.

Communication of new names at a public meeting, or the placing of names in collections or gardens open to the public, do not constitute publication.

Examples: Effective publication without printed matter: *Salvia oxyodon* Webb and Heldr. was published in July, 1850, in an autograph catalogue and put on sale (Webb and Heldreich, *Catalogus plantarum hispanicarum, etc., ab. A. Blanco lectarum*, Parisiis, Jul., 1850, in folio).—Non-effective publication at a public meeting: Cusson announced his establishment of the genus *Physospermum* in a memoir read at the Société des Sciences des Montpellier in 1773, and later in 1782 or 1783 at the Société de Médecine de Paris, but its effective publication dates from 1787, in the Mémoires de la Soc. Roy. de Médecine de Paris, vol. V, 1^{re} partie.

Comment. The statement that the communication of new names at a public meeting does not constitute valid publication, is of particular interest to bacteriologists. Priority is always determined from date of publication and not from date of presentation. A committee appointed by the Society of American Bacteriologists on the classification of bacteria at a meeting of the society recommended the use of the generic name *Erwinia* to include certain bacteria producing diseases of plants. This did not constitute valid publication.

Article 36. On and after January 1, 1908, the publication of names of new groups of living plants will be valid only when they are accompanied by a Latin diagnosis. On and after January 1, 1912, the names of new groups of fossil plants will not be considered valid unless accompanied by a Latin diagnosis and by illustrations or figures showing the essential characteristics.

Comment. Bacteriologists in general have never subscribed to this rule. It is unusual to find modern bacteriologists giving Latin description to new species, in fact the rule may be said to be quite commonly ignored by botanists. The committee on classification of the Society of American Bacteriologists has recommended that bacteriologists follow the International Botanical Code in the nomenclature of bacteria with the exception of article 36.

Article 37. A species or a subdivision of a species, announced in a work, with a complete specific or varietal name, but without diagnosis or reference to a former description under another name, is not valid. Citation in synonymy or incidental mention of a name is not effective publication, and the same applies to the mention of name on a ticket issued with a dried plant without printed or autographed diagnosis.

Plates accompanied with analyses are equivalent to a description, but this applies only to plates published before January 1, 1908.

Article 38. A genus or any other group of higher rank than a species, named or announced without being characterized conformably to article 37 cannot be regarded as effectively published (*nomen nudum*). The mere indication of species as belonging to a new genus or of genera as belonging to a higher group, does not allow us to accept the genus or group in question as characterized and effectively published. An exception is made in the case of the generic names mentioned by Linnaeus in the *Species Plantarum* ed. 1, 1753, names which we associate with the descriptions in the *Genera Plantarum* ed. 5, 1754 (see article 19).

Article 39. The date of a name or of a combination of names is that of their effective publication. In the absence of proof to the contrary, the date placed on the work containing the name or combination of names is regarded as correct. On and after January 1, 1908, the date of publication of the Latin diagnosis only can be taken into account in questions of priority.

Recommendations. Botanists will do well, in publishing, to conform to the following recommendations.

XVIII. Not to publish a name without clearly indicating whether it is the name of a family or a tribe, a genus or a section, a species or a variety, briefly, without expressing an opinion on the nature of the group to which they give the name.

When names of new groups are published the author should designate the subdivision which he considers as the type for the nomenclature of the group, he should indicate the generic type in a family, the species type in a genus, the subspecies or variety type in a species. This precaution will prevent difficulties in nomenclature if it should become necessary later to dissociate such groups.

Comment The Zoological Code not only recommends that specific types for genera and subgenera should be designated, but requires that the names of families and orders should be derived from the names of the type genera and type families respectively. This rule would be of assistance in botany and in bacteriology

Recommendation. XIX To avoid publishing or mentioning in their publications unpublished names which they do not accept, especially if the person responsible for these names have not formally authorized their publication (see Rec XIV, e)

Recommendation XX When publishing new names in works written in a modern language (floras, catalogues, etc.) to publish simultaneously the Latin diagnoses and in paleobotany the figures which will make the names valid from the point of view of scientific nomenclature

In view of the particular difficulties indicated in the identification of fossil plants, in addition to the Latin diagnosis, there should be included a detailed description in French, German or Italian

When new groups of the lower Cryptogames especially the fungi or other microscopic plants are described, the description should be accompanied with a figure or figures of these plants with the microscopic details which will aid in their identification

The description of a parasitic plant should always be followed by a designation of the host, particularly in the parasitic fungi. The hosts should be designated by their scientific Latin names and not by vernacular names in a modern language for these frequently have more than one meaning

Recommendation XXI To give the etymology of new generic names and also of specific names when the meaning of the latter is not obvious

XXII To indicate precisely the date of publication of their works and that of the placing on sale or the distribution of named and numbered plants when these are accompanied by printed diagnoses. In the case of a work appearing in parts, the last published sheet of a volume should indicate the precise dates at which the different fascicles or parts of the volume were published as well as the number of pages in each

XXIII When works are published in periodicals to require the editor to indicate on the separate copies the date (year and month) of publication and also the title of the periodical from which the work is extracted

XXIV. Separate copies should always bear the pagination of the periodical of which they form a part, if desired they may also bear a separate pagination

Section 5 On the precision to be given to names by the citation of the author who first published them

Article 40 For the indication of the name or names of a group to be accurate and complete, and in order that the date may be readily verified, it is necessary to quote the author who first published the name or combination of names in question

Examples *Simarubaceae* Lindley, *Simaruba* Aublet, *Simaruba laevis* Grisebach, *Simaruba amara* Aublet var. *opaca* Engler

Comment It is evident from the reading of this rule that the author of the name is frequently not the individual who first described the organism. The name of the author is not quoted following the species in order to designate the discoverer of the organism, or even one necessarily who has studied it in detail, but merely to show who first used the particular combination of letters or words constituting the name under consideration. Any other procedure would be making the discoverer of an organism responsible for a name with which he has never had anything to do. As stated by Stiles (1905, p. 21) "This point of view lays stress upon holding an author responsible for the names he publishes, rather than upon 'giving him credit' for these names. The chief idea we have in citing the author of a name is to aid in tracing it." For example, one of the cocci associated with pus production was first adequately studied by Rosenbach and named by him *Staphylococcus albus*. Later it was renamed *Albococcus albus* by Winslow and Rogers. Certainly Rosenbach is no way responsible for the latter generic name. Welch discovered an organism which produced edema in man. He did not give it a name. Later the organism was termed *Bacillus Welchii*. We certainly could not write *Bacillus Welchii* Welch without giving the impression that Welch had named an organism after himself.

Article 41. An alteration of the constituent characters or of the circumscription of a group does not warrant the quotation of another author than the one who first published the name or combination of names.

When the changes have been considerable, the words *mutatis caract*, or *pro parte*, or *excl gen*, *excl sp*, *excl var*, or some other abridged indication, are added after the citation of the original author, according to the nature of the changes that have been made, and of the group in question.

Examples *Phyllanthus* L. em. (emendavit) Mull. Arg., *Myosotis* L. pro parte, R. Br., *Globularia cordifolia* L. excl. var., etc.

Comment The generic name *Bacterium* was first used by Ehrenberg in 1828. His definition was somewhat modified by Dujardin in 1841. It was again modified by Cohn in 1872 and by Migula in 1894. The name should always be written therefore, *Bacterium* Ehrenberg. This does not indicate that we accept Ehrenberg's diagnosis of the genus but merely that he first used this particular designation.

Article 42. When a manuscript name has been published and referred to its author, the name of the person who published it should be appended to the citation. The same rule should be followed for names of garden origin when they are cited as "Hort."

Article 43. When, in a genus, a name is applied to a group which is moved into another group where it retains the same rank, or to a group which becomes of

higher or lower rank than before, the change is equivalent to the creation of a new group and the author who has effected the change is the one to be quoted. The original author can be cited only in parenthesis.

Examples *Cheiranthus tristis* L. when moved into the genus *Matthiola* becomes *Matthiola tristis* R. Br., or *Matthiola tristis* (L.) R. Br. — *Medicago polymorpha* L. var. *orbicularis* L. when raised to the rank of a species becomes *Medicago orbicularis* Ait. or *Medicago orbicularis* (L.) Ait.

Comment. The corresponding zoological rule requires that the original author shall be cited in parenthesis instead of leaving it optional as in the above article.

Recommendation. XXV. Author's names put after names of plants are abbreviated, unless they are very short.

Section 6 On names that are to be retained when a group is divided, remodelled, transferred, or moved from one rank to another, or when two groups of the same rank are united, or when they concern groups possessing a pleomorphic cycle

Article 44. A change of characters, or a revision which involves the exclusion of certain elements of a group or the addition of new elements, does not warrant a change in the name or names of a group, except in cases provided for in article 51.

Examples The genus *Myosotis* as revised by R. Brown differs from the original genus of Linnaeus, but the name has not been changed, nor is any change allowable. Various authors have united with *Centaurea Jacea* L. one or two species which Linnaeus had kept distinct, the group thus constituted must be called *Centaurea Jacea* L. (sensu ampl.) or *Centaurea Jacea* L. (em Visiani, em Godron, etc.), the creation of a new name such as *Centaurea vulgaris* Godr. is superfluous.

Article 45. When a genus is divided into two or more genera, the name must be kept and given to one of the principal divisions. If the genus contains a section or some other division which, judging by its name or its species, is the type or the origin of the group, the name is reserved for that part of it. If there is no such section or subdivision, but one of the parts detached contains a great many more species than the others, the name is reserved for that part of it.

Examples The genus *Helianthemum* contained according to Dunal (in DC Prodr. I 266-284), 412 well-known species distributed in nine sections, several of these sections have since been raised to generic rank (*Fumana* Spach, *Tuberaria* Spach) but the name *Helianthemum* has been kept for the divisions grouped round the section *Euhelianthemum*. The genus *Convolvulus* L. em Jacq. was divided into two by Robert Brown in 1810 (*Prodr. fl. Nov. Holl.*, p. 482-484), who gave the name *Calystegia* to one of the genera which at the time contained only four species and reserved the name *Convolvulus* for the other genus which contained a much larger number of species. In the same way Salisbury (in *Trans. Linn. Soc.* VI, 317, 1802) in separating *Erica vulgaris* L. from the genus *Erica*, under the name *Calluna*, kept the name *Erica* for the large number of species left.

Comment Winslow and Rogers in 1905 divided the genus *Staphylococcus* of Rosenbach into two genera *Aurococcus* and *Albococcus*, the name *Staphylococcus* should have been retained for one of these groups

Article 46. When two or more groups of the same nature are united, the name of the oldest is retained. If the names are of the same date, the author chooses and his choice cannot be modified by subsequent authors

Examples Hooker f and Thomson (Fl Ind p 67, 1855) united the genera *Wormia* Rotth and *Capellia* Bl, they gave the name *Wormia* to the genus thus formed because the last name dates from 1783 while *Capellia* dates from 1825—In case of union of the two genera *Cardamine* and *Dentaria*, which were founded at the same time by Linnaeus (*Sp Pl* ed 1, p 653 and 654, 1753, *Gen Pl* ed 5, n 726, 727) the collective genus must be called *Cardamine* because that name was chosen by Crantz (*Class Crucif*, p 126, 1769), who was the first to suggest the union

Recommendations XXVI Authors who have to choose between two generic names should note the following recommendations

1 Of two names of the same date to prefer the one which was first accompanied by the description of a species

2. Of two names of the same date, both accompanied by descriptions of species, to prefer the one, which, when the author made his choice, included the larger number of species

3 In cases of equality from these various points of view to prefer the more correct and appropriate name

XXVII When several genera are united as subgenera or sections under one generic name, that subdivision which was first distinguished or described may retain its name (ex *Saxifraga aspera* subsp *aspera*) or bear a prefix (*Alchemilla alpina* subsp *eualpina*) or be designated by some customary title (*normalis*, *genuinus*, *typicus*, *originarius*, *verus*, *veridicus* etc) These prefixes or terms lapse when the subdivisions are raised to specific rank

Article 47 When a species or subdivision of a species is divided into two or more groups of the same nature, if one of the two forms was distinguished or described earlier than the other, the name is retained for that form

Article 48 When a subgenus or section or species is moved into another genus, when a variety or other division of a species is moved into another species, retaining there the same rank, the original name of the subgenus or section, the first specific epithet, or the original name of the division of the species must be retained or must be reestablished, unless, in the new position there exists one of the obstacles indicated in the articles of section 7

Article 49 When a tribe becomes a family, a subgenus or a section becomes a genus, a subdivision of a species becomes a species, or the reverse of these changes takes place, and speaking generally when a group changes its rank, the earliest name (or combination of names) received by the group in its new position must be regarded as valid, if it is in conformity with the rules, unless there exist any of the obstacles indicated in the articles of section 7

Article 49a. Fungi having a pleomorphic evolutionary cycle in different successive states of the same species carry only a single generic and specific name (binôme), the oldest which has been given beginning with Persoon (Synopsis)

or Fries (Systema), to the state containing the form which is generally regarded as being perfect, on condition that in other respects it conforms to the rules

The ascus stage is considered the perfect state in the *Ascomycetes*, the basidia as the stage in the *Basidiomycetes*, the teleutospore or its equivalent in the *Uredinales*, and the spore in the *Ustilaginales*

Generic and specific names given to other stages possess only a temporary value. They are not to be used to replace a generic name already existing and applied to one or more species, any one of which contains the form termed perfect

The nomenclature of fungi with an evolutionary cycle, but not pleomorphic follows the ordinary rules of nomenclature

Recommendations. XXIX 1° When a sub-tribe becomes a tribe, when a tribe becomes a subfamily, when a subfamily becomes a family, etc., or when the inverse changes occur, do not alter the root of a name but only the termination (-inae, -eae, -oideae, -ineae, -ales, etc.), unless in the new position, one of the obstacles indicated in the articles of section 7, supervenes, or the new designation becomes a source of error, or there is some other serious reason against it

2° When a section or a subgenus becomes a genus, or the inverse changes take place, retain the old names, unless this results in two genera of plants having the same name, or the existence of two subdivisions of the same name in the same genus, or one of the obstacles indicated in the articles of section 7 supervenes

3° When a subdivision of a species becomes a species or the inverse change occurs, retain the original epithets, unless this results in two species bearing the same name in the same genus, or two subdivisions bearing the same name in the same species, or unless any of the obstacles indicated in section 7 supervenes

Section 7 On names that are to be rejected, changed or modified

Article 50 No one is authorized to reject, change or modify a name (or combination of names) because it is badly chosen, or disagreeable, or another is preferable or better known, or because of the existence of an earlier homonym which is universally regarded as non-valid, or for any other motive either contestable or of little import (See also art. 57)

Recommendations See on the subject of homonyms recommendations Vb and XIV which suggest that cases of this kind should be avoided for the future

Comment A large proportion of the names which have been suggested for genera and species of bacteria must be rejected because of violation of this article. Orla-Jensen (1909) for example, created a large number of new genera, in order to make them conform to his ideas of fitness. Where such names are applied to groups which have had no previous generic designation they are, of course, valid, but he was not justified in substituting names which he preferred for old and valid designations.

The expression "because of the existence of an earlier homonym which is universally regarded as non-valid" is one which permits of a variety of interpretations. The zoologists insist upon the rejection of any name which is a homonym of an earlier term. In this respect the two codes are at variance.

Article 51 Every one should refuse to admit a name in the following cases

1. When the name is applied in the plant kingdom to a group which has an earlier valid name
2. When it duplicates the name of a class, order, family or genus, or a subdivision or species of the same genus, or a subdivision of the same species
3. When it is based on a monstrosity
4. When the group which it designates embraces elements altogether incoherent, or when it becomes a permanent source of confusion or error
5. When it is contrary to the rules of sections 4 and 6

Comments 1 The generic name *Pneumococcus* of Schmidlechner (1905) is an invalid synonym because of the older name *Diplococcus* given to the same group by Weichselbaum (1886)

2 The generic name *Myxococcus* of Gonnermann (1907) duplicates the name of an entirely different group termed *Myxococcus* by Thaxter (1892)

4 Much difficulty is found in bacteriology in determining the proper designation for some of the older species of bacteria because of the fact that some of the older authors were dealing with mixed or impure cultures

Article 53 When a subgenus, a section or a subsection, passes as such into another genus, the name must be changed if there is already in that genus a valid group of the same rank, under the same name

When a species is moved from one genus into another, its specific epithet must be changed if it is already borne by a valid species of the genus. Similarly when a subspecies, a variety, or some other subdivision of a species is placed under another species, its name must be changed if borne already by a valid form of like rank in that species

Article 54 Names of genera must be rejected in the following special cases

1. When they coincide with technical term currently employed in morphology unless they are accompanied by specific names
2. When they express uninomial nomenclature
4. When they are formed of two words, unless these two words were from the first united or joined by a hyphen

Examples 1° Generic names such as *Lignum*, *Radix*, *Spina*, etc., would not now be admissible, on the other hand a generic name like *Tuber* should not be rejected when it has been published with specific names (*Tuber cibarium*, etc.) — 2° Ehrhart (*Phytophylacium* (1780)) and Beirag (IV, 145-150) made use of a uninomial nomenclature for species known at that time under binary names (*Phaeocephalum*, *Leptostachys*, etc.) These names, which resemble generic names, must not be confused with such and are to be rejected, unless a subsequent author has given them the value of a generic name for example *Baeothryon*, a uninomial expression of Ehrhart's has been applied to a genus characterized by A. Dietrich *Spec. Pl.* II, 89 (1833) — Names like *Quisqualis* (a single word from the first), *Sebastiano-Schaueria* and *Neves-Aimondia* will stand.

GENERAL SYSTEMATIC BACTERIOLOGY

Article 55. Specific names must also be rejected in the following cases

1°. When they are ordinals serving for purpose of enumeration

2°. When they merely repeat the generic name

Examples 1° *Boletus vicesimus sextus*, *Agaricus octogesimus nonus* 2°
Linaria Linaria, *Raphanistrum Raphanistrum*, etc

Comments. 1 Many bacterial species have been named by the use of numbers or letters Where these can be clearly differentiated, new specific names must be given

2 This rule in the botanical code is contrary to that in the Zoological Code Names such as *Apis apis* may be used in zoological literature, but not in botany, hence not in bacteriology A varietal name may be the same as its species but the species name must be different from its genus In zoology on the other hand, variety, genus and species may all have the same name

Article 56. In the cases foreseen in articles 51 to 56, the name to be rejected or changed is replaced by the oldest valid name in the group in question, and in default of such a one a new name (a new binomial) must be made

By a valid name is meant a name or more particularly a combination of names created in conformity with the whole of the rules of nomenclature The author of a new combination ought to borrow the specific epithet of an old non-valid binomial, or employ a new name

Article 57 The original spelling of a name must be retained, except in case of a typographic or orthographic error When the difference between two names, especially two generic names, lies in the termination these names are to be regarded as distinct even though differing by one letter only

Examples *Rubia* and *Rubus*, *Monochaete* and *Monochaetum*, *Peponia* and *Peponum*, *Iria* and *Iris*

Recommendations XXX The liberty of making orthographic corrections must be used with reserve, especially if the change effects the first syllable, and above all the first letter of a name

XXXI Many names differ by a single letter without risk of confusion (ex *Durvillea* and *Urvillea*) In cases where a close approach to identity is a source of error (ex *Astrostemma* and *Asterostemma* in one and the same family, *Asclepiadaceae*, *Pleuropetalum* and *Pleuropetalum* in *Orchidaceae*) only one, the older, of the names should be kept, in accordance with article 51, 4°

Chapter IV. Modification of the rules of botanical nomenclature

Article 58 The rules of botanical nomenclature can only be modified by competent persons at an International Congress convened for the express purpose

C. THE INTERNATIONAL RULES OF ZOOLOGICAL NOMENCLATURE

Although the bacteria are plants, and the bacteriologist is for the most part dealing with bacteria, it must be recognized that the bacteria

intergrade with the protozoa. The bacteriologist must frequently deal also with problems in the nomenclature of protozoa, or even of higher animals. He is therefore vitally interested in the rules used by the zoologist in his systematic nomenclature.

In many respects the zoological rules are more explicit than those of the botanist. They constitute primarily a type basis code. To many bacteriologists this appears to be highly desirable in the field of bacteriology. While the Zoological Code therefore cannot govern bacterial nomenclature, nevertheless an understanding of its rules and recommendations would be of great value in determining good usage on points not covered in the Botanical Code.

The code given below is that adopted by the Ninth International Zoological Congress at Monaco, 1913.

Article 1. Zoological nomenclature is independent of botanical nomenclature in the sense that the name of an animal is not to be rejected simply because it is identical with the name of a plant. If, however, an organism is transferred from the vegetable to the animal kingdom its botanical names are to be accepted in zoological nomenclature with their original botanical status, and if an organism is transferred from the animal to the vegetable kingdom its names retain their zoological status.

Recommendation. It is well to avoid introducing into zoology as generic names such names as are in use in botany.

Article 2. The scientific designation of animals is uninominal for subgenera and all higher groups, binominal for species, and trinominal for subspecies.

Article 3. The scientific names of animals must be words which are either Latin or Latinized, or considered and treated as such in case they are not of classic origin.

Family and subfamily names

Article 4. The name of a family is formed by adding the ending *idae*, the name of a subfamily by adding *inae* to the stem of the name of its type genus.

Article 5. The name of a family or subfamily is to be changed when its type genus is changed.

Generic and subgeneric names

Article 6. Generic and subgeneric names are subject to the same rules and recommendations, and from a nomenclatural standpoint they are coordinate, that is, they are of the same value.

Article 7. A generic name becomes a subgeneric name, when the genus so named becomes a subgenus, and *vice versa*.

Article 8. A generic name must consist of a single word, simple or compound, written with an initial capital letter, and employed as a substantive in the nominative singular. Examples: *Cens*, *Porca*, *Ceratodus*, *Hymenolepis*.

GENERAL SYSTEMATIC BACTERIOLOGY

Recommendation. Certain biological groups which have been proposed distinctly as collective groups, not as systematic units, may be treated for convenience as if they were genera, but they require no type species. Examples *Agamodistomum*, *Amphistomulum*, *Agamofilaria*, *Agamomermis*, *Sparganium*

Recommendations The following words may be taken as generic names

a Greek substantives, for which the rules of Latin transcription (transliteration (see Appendix F)) should be followed. Examples *Ancylus*, *Amphibola*, *Aplysia*, *Pompholyx*, *Physa*, *Cylindrina*

b. Compound Greek words, in which the attributive should precede the principal word. Examples *Stenogyra*, *Pleurobrachius*, *Tylodina*, *Cyclostomum*, *Sarcocystis*, *Pelodytes*, *Hydrophilus*, *Rhizobius*.

This does not, however, exclude words formed on the model of *Hippopotamus*, namely, words in which the attributive follows the principal word. Examples *Philydrus*, *Biorhiza*

c Latin substantives. Examples *Ancilla*, *Auricula*, *Dolum*, *Harpa*, *Oliva*. Adjectives (*Prasina*), past participles (*Productus*) are not recommended

d Compound Latin words. Examples *Staliger*, *Dolabrifer*, *Semifusus*

e Greek or Latin derivatives expressing diminution, comparison, resemblance, or possession. Examples *Dolum*, *Dololum*, *Strongylus*, *Eustrongylus*, *Limaz*, *Lamacella*, *Limacia*, *Lamacina*, *Limautes*, *Lamacula*, *Lingula*, *Lingulicella*, *Lingulepis*, *Lingulina*, *Lingulops*, *Lingulopsis*, *Neomema*, *Proneomema*, *Buteo*, *Archibuteo*, *Gordius*, *Paragordius*, *Polygordius*

f Mythological or heroic names. Examples *Osiris*, *Venus*, *Brisinga*, *Velleda*, *Crimora*. If not Latin, these should be given a Latin termination (*Aegirus*, *Göndulha*)

g Proper names used by the ancients. Examples *Cleopatra*, *Belisarius*, *Melania*

h Modern patronymics, to which is added an ending to denote dedication

α Names terminating with a consonant taking the ending *ius*, *ia*, or *ium*. Examples *Stelysius*, *Lamarchia*, *Kollikeria*, *Mulleia*, *Stalia*, *Kryeria*, *Ibanezia*

β Names terminating with the vowels *e*, *i*, *o*, *u*, or *y* take the ending *us*, *a* or *um*. Examples *Blainvillea*, *Wyvillea*, *Carolina*, *Fatioa*, *Bernaya*, *Quoya*, *Schulzeia*

γ Names terminating with *a* take the ending *ia*. Example *Danaria*

δ In generic names formed from patronymics, the particles are omitted if not coalesced with the name, but the articles are retained. Examples *Blainvillea*, *Benedenia*, *Chajaea*, *Lacapedea*, *Dumerilia*

ε With patronymics consisting of two words, only one of these is used in the formation of a generic name. Examples *Selysius*, *Targionia*, *Edwardsia*, *Duthiersia*

ζ The use of proper names in the formation of compound generic names is objectionable. Examples *Eugrammia*, *Buchiceras*, *Heromorpha*, *Mobiusispongia*

ι Names of ships which should be treated the same as mythological names (*Vega*) or as modern patronymics. Examples *Blakea*, *Hirondellea*, *Challengeria*

θ Barbarous names, that is, words of nonclassic origin. Examples *Vanikoro*, *Chilosa*. Such words may receive a Latin termination. Examples *Yetus*, *Fossarius*

k Words formed by an arbitrary combination of letters Examples *Neda*, *Clanculus*, *Sahfa*, *Toriz*

1. Names formed by anagram Examples *Dacelo*, *Verlusia* ? *Linospa*

Article 9 If a genus is divided into subgenera, the name of the typical subgenus must be the same as the name of the genus (see Art 25)

Article 10. When it is desired to cite the name of a subgenus, this name is to be placed in parentheses between the generic and the specific names Examples *Vanessa (Pyramus) cardui*

Specific and subspecific names

Article 11 Specific and subspecific names are subject to the same rules and recommendations, and from a nomenclatural standpoint they are coordinate, that is, they are of the same value

Article 12 A specific name becomes a subspecific name when the species so named becomes a subspecies, and *vice versa*

Article 13 While specific substantive names derived from names of persons may be written with a capital initial letter, all other specific names are to be written with a small initial letter Examples *Rhizostoma Cuvieri* or *Rh cuvieri*, *Francolinus Lucani*, or *F lucani*, *Hypoderma Diana* or *H diana*, *Laophonte Mohammed* or *L mohammed*, *Oestrus ovis*, *Corvus corax*

Article 14 Specific names are

a. Adjectives, which must agree grammatically with the generic name
Example *Felis marmorata*

b. Substantives in the nominative in apposition with the generic name
Example *Felis leo*

c Substantives in the genitive Examples *Rosae*, *sturonis*, *antillarum*, *galhae*, *sancti-pauli*, *sanctae-helenae*

If the name is given as a dedication to one or several persons, the genitive is formed in accordance with the rules of Latin declination in case the name was employed and declined in Latin Examples *Plinii*, *Aristotelis*, *Victoris*, *Antonii*, *Elisabethae*, *Petri* (given name)

If the name is a modern patronymic, the genitive is always formed by adding, to the exact and complete name, an *i* if the person is a man, or an *ae* if the person is a woman, even if the name has a Latin form, it is placed in the plural if the dedication involves several persons of the same name Examples *Cuvieri*, *Möbrius*, *Nunezi*, *Meranae*, *Sarasinorum*, *Bovi* (not *Bovis*), *Salmoni* (not *Salmonis*)

Recommendation The best specific name is a Latin adjective, short, euphonic, and of easy pronunciation Latinized Greek words or barbarous words may, however, be used Examples *gymnocephalus*, *echinococcus*, *ziczac*, *aguti*, *hoaculi*, *urubitinga*

It is well to avoid the introduction of the names *typicus* and *typus* as new names for species or subspecies, since these names are always liable to result in later confusion

Article 15 The use of compound proper names indicating dedication, or of compound words indicating a comparison with a simple object does not form an exception to Article 2 In these cases the two words composing the specific name

are written as one word with or without the hyphen Examples *Sanctae-Catharinae* or *sanctacatherinae*, *jan-mayeni*, or *janmayeni*, *cornu-pastoris* or *cornupastoris*, *cor-anguinum* or *coranguinum*, *cedo-nulli* or *cedonullu*

Expressions like *rudis planusque* are not admissible as specific names

Article 16. Geographic names are to be given as substantives in the genitive, or are to be placed in an adjectival form Examples *sancti-pauli*, *sanctae-helenae*, *edwardiensis*, *dremenensis*, *magellanicus*, *burdigalensis*, *vindobonensis*

Recommendation Geographic names used by the Romans or by Latin writers of the middle ages are to be adopted in preference to more recent forms Words like *bordeausiacus* and *vinnensis* are poor, but are not to be rejected on this account

Article 17. If it is desired to cite the subspecific name, such name is written immediately following the specific name, without the interposition of any mark of punctuation Example *Rana esculenta marmorata* Hallowell, but not *Rana esculenta (marmorata)* or *Rana marmorata* Hallowell

Article 18. The notation of hybrids may be given in several ways, in all cases the name of the male parent precedes that of the female parent, with or without the sexual signs

a The names of the two parents are united by the sign of multiplication (\times)

b Hybrids may also be cited in the form of a fraction, the male parent forming the numerator and the female parent the denominator Example $\frac{\textit{Capra hircus}}{\textit{Ovis aries}}$

This second method is in so far preferable that it permits the citation of the person who first published the hybrid form as such Example

$\frac{\textit{Bernicla canadensis}}{\textit{Anser cygnoides}}$ Rabé

c The fractional form is also preferable in case one of the parents is itself a hybrid Example $\frac{\textit{Tetrao tetrix} \times \textit{Tetrao urogallus}}{\textit{Gallus gallus}}$ In the latter case, however, the parentheses may be used Example $(\textit{Tetrao tetrix} \times \textit{Tetrao urogallus}) \times \textit{Gallus gallus}$

d When the parents of the hybrid are not known as such (parents), the hybrid takes provisionally a specific name, the same as if it were a true species, namely, as if it were not a hybrid, but the generic name is preceded by the sign of multiplication Example $\times \textit{Coregonus dolosus}$ Fatio

Formation, derivation, and orthography of zoological names

Article 19. The original orthography of a name is to be preserved unless an error of transcription, a *lapsus calami*, or a typographical error is evident

Recommendation For scientific names it is advisable to use some other type than that used for the text Example *Rana esculenta* (italics) Linné, 1758, lives in Europe

Article 20. In forming names derived from languages in which the Latin alphabet is used, the exact original spelling, including diacritic marks, is to be retained Examples *Selysius*, *Lamarckia*, *Köllikeria*, *Mülleria*, *Stahia*, *Kröyeria*, *Ibaneza*, *Mobiusi*, *Medaci*, *Czyzeki*, *spitzbergensis*, *islandicus*, *paraguayensis*, *patagonicus*, *barbadensis*, *fardensis*

Recommendations The prefixes *sub* and *pseudo* should be used only with adjectives and substantives, *sub* with Latin words, *pseudo* with Greek words, and they should not be used in combination with proper names. Examples *subviridis*, *subchelatus*, *Pseudacanthus*, *Pseudophis*, *Pseudomys*. Words like *subwilsoni* and *pseudo-grateloupana* are not recommended.

The terminations *oides* and *ides* should be used in combination only with Greek or Latin substantives, they should not be used in combination with proper names.

Geographic and patronymic names from countries which have no recognized orthography or which do not use the Latin alphabet, should be transcribed into Latin according to the rules adopted by the Geographic Society of Paris. (See Appendix G.)

In proposing new names based upon personal names, which are written sometimes with *a*, *o* or *u*, at other times with *ae*, *oe*, and *ue*, it is recommended that authors adopt *ae*, *oe* and *ue*. Example *muelleri* in preference to *mulleri*.

Author's name.

Article 21. The author of a scientific name is that person who first publishes the name in connection with an indication, a definition, or a description, unless it is clear from the contents of the publication that some other person is responsible for said name and its indication, definition, or description.

Article 22 If it is desired to cite the author's name, this should follow the scientific name without interposition of any mark of punctuation, if other citations are desirable (date, *sp. n.*, *emend.*, *sensu stricto*, etc.) these follow after the author's name, but are separated from it by a comma or by parenthesis. Examples *Primates* Linné, 1758, or *Primates* Linné (1758).

Recommendation When the name of the author of a scientific name is abbreviated, the writer will do well to conform to the list of abbreviations published by the Zoological Museum of Berlin.

Article 23 When a species is transferred to another than the original genus or the specific name is combined with any other generic name than that with which it was originally published, the name of the author of the specific name is retained in the notation but placed in parentheses. Examples *Taenia lata* Linné, 1758, and *Dibothriocephalus latus* (Linné, 1758), *Fasciola hepatica* Linné, 1758, and *Distoma hepaticum* (Linné, 1758).

If it is desired to cite the author of the new combination his name follows the parentheses. Example *Limnatis nilotica* (Savigny, 1820) Moquin-Tandon, 1826.

Article 24 When a species is divided, the restricted species to which the original specific name of the primitive species is attributed may receive a notation indicating both the name of the original author and the name of the reviser. Example *Taenia solium* Linné, partim, Goeze.

The law of priority

Article 25 The valid name of a genus or species can be only that name under which it was first designated on the condition

a. That this name was published and accompanied by an indication, or a definition, or a description, and

b. That the author has applied the principles of binary nomenclature.

Application of the law of priority

Article 26. The Law of Priority obtains and consequently the oldest available name is retained

- a. When any part of an animal is named before the animal itself
- b. When any stage in the life history is named before the adult
- c. When the two sexes of an animal have been considered as distinct species or even as belonging to distinct genera
- d. When an animal represents a regular succession of dissimilar generations which have been considered as belonging to different species or even to different genera

Article 28 A genus formed by the union of two or more genera or subgenera takes the oldest valid generic or subgeneric name of its components. If the names are of the same date, that selected by the first reviser shall stand.

The same rule obtains when two or more species or subspecies are united to form a single species or subspecies.

Recommendation In absence of any previous revision, the establishment of precedence by the following method is recommended.

- a. A generic name accompanied by specification of a type has precedence over a name without such specifications. If all or none of the genera have types specified, that generic name takes precedence the diagnosis of which is most pertinent.
- b. A specific name accompanied by both description and figure stands in preference to one accompanied only by a diagnosis or only by a figure.
- c. Other things being equal, that name is to be preferred which stands first in the publication (page precedence).

Article 29 If a genus is divided into two or more restricted genera, its valid name must be retained for one of the restricted genera. If a type was originally established for said genus, the generic name is retained for the restricted genus containing said type.

Recommendation To facilitate reference, it is recommended that when an older species is taken as type of a new genus its name should be actually combined with the new generic name in addition to citing it with the old generic name. Example *Gilbertella* Eigenmann, 1903, Smithsonian Misc. Coll. v 45, p 147, type *Gilbertella alata* (Steindachner) = *Anacyrtus alatus* Steindachner.

Article 30 The designation of type species of genera shall be governed by the following rules (a-g), applied in the following order of precedence.

I Cases in which the generic type is accepted *solely* upon the basis of the original publications.

- a. When in the original publication of a genus, one of the species is definitely designated as type, this species shall be accepted as type, regardless of any other considerations. (Type by original designation.)

- b. If in the original publication of a genus, *typicus* or *typus* is used as a *new* specific name for one of the species, such use shall be construed as "type by original designation."

- c. A genus proposed with a single original species takes that species as its type (Monotypical genera.)

- d. If a genus, without originally designated (see a) or indicated (see b) type, contains among its original species one possessing the generic name as its specific

or subspecific name, either as valid name or synonym, that species or subspecies becomes *ipso facto* type of the genus (Type by absolute tautonymy)

II Cases in which the generic type is accepted not solely upon basis of the original publication

e The following species are excluded from consideration in determining the types of genera

α Species which were not included under the generic name at the time of its original publication

β Species which were *species inquirendae* from the standpoint of the author of the generic name at the time of its publication

γ Species which the author of the genus doubtfully referred to it

f. In case a generic name without originally designated type is proposed as a substitute for another generic name, with or without type, the type of either, when established, becomes *ipso facto* type of other

g If an author, in publishing a genus with more than one valid species, fails to designate (see a) or to indicate (see b, d) its type, any subsequent author may select the type, and such designation is not subject to change (Type by subsequent designation)

The meaning of the expression "select the type" is to be rigidly construed. Mention of a species as an illustration or example of a genus does not constitute a selection of a type

III Recommendations In selecting types by subsequent designation, authors will do well to govern themselves by the following recommendations

h In case of Linnaean genera, select as type the most common or the medicinal species (Linnaean rule, 1751)

i If a genus, without designated type, contains among its original species one possessing as a specific or subspecific name, either as valid name or synonym, a name which is virtually the same as the generic name, or of the same origin or same meaning, preference should be shown to that species in designating the type, unless such preference is strongly contraindicated by other factors (Type by virtual tautonymy) Examples *Bos taurus*, *Equus caballus*, *Ovis aries*, *Scomber scombrus*, *Sphaerostoma globosum*, contraindicated in *Dipetalonema* (compare species *Filaria dipetala*, of which only one sex was described, based upon one specimen and not studied in detail)

j If the genus contains both exotic and nonexotic species from the standpoint of the original author, the type should be selected from the nonexotic species

k If some of the original species have later been classified in other genera, preference should be shown to the species still remaining in the original genus (Type by elimination)

l Species based upon sexually mature specimens should take precedence over species based upon larval or immature forms

m Show preference to species bearing the name *communis*, *vulgaris*, *medicinalis*, or *officinalis*

n Show preference to the best described, best figured, best known, or most easily obtainable species, or to one of which a type specimen can be obtained

o Show preference to a species which belongs to a group containing as large a number of the species as possible (De Candolle's rule)

p. In parasitic genera, select, if possible, a species which occurs in the man or some food animal or in some very common and widespread host species

q. All other things being equal, show preference to a species which the author of the genus actually studied at or before the time he proposed the genus

r. In case of writers who habitually placed a certain leading or typical species first as "chef de file," the others being described by comparative reference to this, this fact should be considered in the choice of the type species

s. In case of those authors who have adopted the "first species rule" in fixing generic types, the first species named by them should be taken as types of their genera

t All other things being equal, page precedence should obtain in selecting a type

Article 32 A generic or a specific name, once published, can not be rejected, even by its author, because of inappropriateness. Examples Names like *Polyodon*, *Apus*, *albus*, etc., when once published, are not to be rejected because of a claim that they indicate characters contradictory to those possessed by the animals in question

Article 33 A name is not to be rejected because of tautonymy, that is, because the specific or the specific and subspecific names are identical with the generic name. Examples *Trutia trutia*, *Apus apus apus*

Article 34 A generic name is to be rejected as a homonym when it has previously been used for some other genus of animals. Example *Trichina* Owen, 1835, nematode, is rejected as homonym (1) of *Trichina* Meigen, 1830, insect

Article 35. A specific name is to be rejected as a homonym (1) when it has previously been used for some other species of the same genus. Examples *Taenia ovilla* Rivolta, 1878 (n. sp.) is rejected as homonym of *T. ovilla* Gmelin, 1790

When in consequence of the union of two genera, two different animals having the same specific or subspecific name are brought into one genus, the more recent specific or subspecific name is to be rejected as a homonym

Specific names of the same origin and meaning shall be considered homonyms if they are distinguished from each other only by the following differences

a The use of *ae*, *oz*, and *e*, as *caeruleus*, *coeruleus*, *ceruleus*, *ei*, *z*, and *y* as *chiropus*, *cheiropus*, *c* and *k* as *microdon*, *mikrodon*

b. The aspiration or non-aspiration of a consonant, as *oxyrynus*, *ocyrrhynchus*

c The presence or absence of a *c* before *t*, as *autumnalis*, *auctumnalis*

d By a single or double consonant, *litoralis*, *littoralis*

e By the endings *ensis* and *iensis* to a geographical name, as *timorensis*, *timoriensis*

Article 36 Rejected homonyms (1) can never be again used. Rejected synonyms can again be used in case of the restoration of erroneously suppressed groups. Example *Taenia Giardi* Moniez, 1870, was suppressed as a synonym of *Taenia ovilla* Rivolta, 1878, later it was discovered that *Taenia ovilla* was pre-occupied (*Taenia ovilla* Gmelin, 1790). *Taenia ovilla*, 1878, is suppressed as a homonym, and can never again be used, it was stillborn and cannot be brought to life, even when the species is placed in another genus (*Thysanosoma*). *Taenia Giardi*, 1879, which was suppressed as a synonym, becomes valid upon the suppression of the homonym *Taenia ovilla* Rivolta

Recommendations. It is well to avoid the introduction of new generic names which differ from generic names already in use only in termination or in a slight

variation in spelling which might lead to confusion. But when once introduced, such names are not to be rejected on this account. Examples *Picus*, *Pica*, *Polyodus*, *Polyodon*, *Polyodonta*, *Polyodontas*, *Polyodontus*.

The same recommendation applies to new specific names in any given genus. Examples *necator*, *necatrix*, *furcigera*, *furcifera*, *rhopalocéphala*, *rhopaliocephala*.

If from the radical of a geographic name two or more adjectives are derived, it is not advisable to use more than one of them as specific name in the same genus, but if once introduced they are not to be rejected on this account. Examples *hispanus*, *hispanicus*, *moluccanus*, *molluccensis*, *sinensis*, *sincus*, *chinensis*, *ceylonicus*, *zeylanicus*.

The same recommendation applies also to other words derived from the same radical and differing from each other only in termination or by a simple change in spelling.

D. TYPE BASIS CODE SUGGESTED BY A COMMITTEE ON BOTANICAL NOMENCLATURE OF THE BOTANICAL SOCIETY OF AMERICA

For a number of years the botanists of America have been divided into two groups, those who use the Vienna Code and those who use the so-called "American Code". In 1918 a Committee on Botanical Nomenclature was appointed by the Botanical Society of America to consider this subject and report. In 1920 the committee gave a majority report signed by seven of its members and a minority report signed by two. The majority report outlined a Type Basis Code of Botanical Nomenclature. The report is of so much importance that it is given below in full.

The Committee on Nomenclature of the Botanical Society of America was appointed at the Baltimore meeting of the Society in 1918. A preliminary survey of the situation showed that the taxonomists of the Society were about equally divided in their support of the Vienna Code and the American Code. An attempt to commit the Society to one or the other of these codes would inevitably result in conflict detrimental to progress in nomenclatural matters. The policy was adopted of working on nomenclature along parallel lines. Two subcommittees were appointed, one on the Vienna Code, with J. M. Greenman as chairman, and one on the American Code, with A. S. Hitchcock as chairman.

The subcommittee on the American Code has prepared a plan which is now submitted to the Society. It is not asked that the Society commit itself to this plan by adopting the report but it is hoped that the Society will receive the report without prejudice. The committee will then continue to study the plan and modify it if necessary to meet the wishes of its supporters. A part of the plan is the establishment of a Nomenclature Commission to pass upon exceptions to the Code. The Committee asks authority to organize this Commission.

We wish especially to call attention of the Society to the fact that the new code which we call the Type-basis Code is not antagonistic to the Vienna Code but is based upon the same principles with certain additions. Both codes are based upon the principle of priority. Both admit exceptions to the application of the

rules The Type-basis Code provides for a Commission to pass upon exceptions The fundamental difference is the acceptance by the Type-basis Code of the principle of Nomenclatural Types This principle is not included in the Vienna Code but is not antagonistic to it The Committee offers the Type-basis Code of Botanical Nomenclature as a new code international in its scope with the hope that its merits will be so evident that it will receive support both in this country and abroad and that it may be combined with the Vienna Code at the next international Congress The chief differences between the two codes are (1) the concept of types, and (2) the idea of a commission to validate exceptions

A type-basis code of botanical nomenclature. Principles.

1 The primary object of formal nomenclature in systematic biology is to secure stability, uniformity, and convenience in the designation of plants and animals

2 Botanical nomenclature is treated as beginning with the general applications of binomial names to plants (Linnaeus' *Species Plantarum*, 1753)

3 Priority of publication is a fundamental principle of botanical nomenclature Two groups of the same category cannot bear the same name

Note a This principle applies primarily to genera and species

Note b Previous use of a name in zoology does not preclude its use in botany, but the proposal of such a name should be avoided

4 The application of names is determined by means of nomenclatural types

Note A generic name is always so applied as to include its type species, a specific name is always so applied as to include its type specimen

Rules and recommendations

Section 1 Publication of names

Article 1 A specific name is published when it has been printed and distributed with a description, or with a reference to a previously published description

Note A recognizable figure may be the equivalent of a description in the literature of paleobotany and diatoms

a In the transfer of a species from one genus to another, the original specific name is retained, unless the resulting binomial has been previously published

Recommendations Botanists will do well, in publishing

1 In describing parasitic fungi to indicate the host and to designate the name of the host by its scientific Latin name

2 To give the etymology of all new generic names

Article 2. A generic name is published when it has been printed and distributed

a. With a generic or specific description (or a recognizable figure, see Article 1, note) and a binomial specific name

b. With a generic and specific name and the citation of a previously published description.

c. With a definite reference to at least one previously published binomial

Note a A name is not published by its citation in synonymy, nor by incidental mention. Such a name may be taken up but not to replace one already properly published

Note b Of names published in the same work and at the same time, those having precedence of position are to be regarded as having priority

Recommendation. Botanists will do well, in publishing, to give the etymology of specific names when their meaning is not obvious

Section 2 Application of names

Article 3 The nomenclatural type of a species is the specimen or the most important of the specimens upon which its original published description was based

a. If only one specimen is cited, that is the type

b. If one specimen is designated as the type, that specimen shall be so accepted, unless an error can be demonstrated

c. A species transferred without change of name from one genus to another retains the original type even though the description under the new genus was drawn from a different species

d. The publication of a new specific name as an avowed substitute for an earlier one does not change the type of the species

e. When more than one specimen was originally cited and no type was designated the type should be selected in the following manner

1. The type specimen interprets the description and fixes the application of the name, hence, primarily the description controls the selection of the type

2. The type may be indicated by the specific name, this being sometimes derived from the collector, locality, or host

3. If one specimen is figured in connection with the original description this may usually be regarded as the type

4. Specimens that are mentioned by the author as being exceptional or unusual, or those which definitely disagree with the description (provided others agree) may usually be excluded from consideration in selecting the type

5. An examination of the actual sheets of specimens studied by the author may aid in determining or selecting the type. He may have written the name or left notes or drawings upon one of the sheets

Note Specimens known to have been received by the author subsequent to the study resulting in the original publication should be excluded from consideration

6. If an author, in publishing a new species, gives a description of his own, this takes precedence over synonymy or cited descriptions, in determining the type specimen

Article 4 The nomenclatural type species of a genus is the species or one of the species included when the genus was originally published

a. If a genus includes but one species when originally published this species is the type

b. When more than one species is included in the original publication of the genus, the type is determined by the following rules (these rules are Articles 3 to 6 of the Report of the Committee on Generic Types published in (1) *Science* N S 49 334-336 1919, (2) *Bot Soc Amer Publ* 73, 70-71 1919)

Recommendations In the future it is recommended that authors of generic names definitely designate type species, and that in the selection of types of genera previously published, but of which the type would not be indicated by the preceding rules, the following points be taken into consideration (This includes Article 7, a to g, of the Report on Generic Types published in *Science* and in *Bot Soc Amer Publ* 73, loc cit)

Section 3 Rejection of Names

Article 5 A name is rejected

a. When preoccupied (homonym)

1. A specific name is a homonym when it has been published for another species under the same generic name

2. A generic name is a homonym when previously published for another genus

3. Similar names are to be treated as homonyms only when they are mere variations in the spelling of the same word, or in the case of specific names, when they differ only in adjective or genitive termination

b. When there is an older valid name based on another member of the same group (metonym)

c. When there is an older valid name based on the same type (typonym)

d. When it has not been effectively published according to the provisions of Section 1 of these rules (hyponym)

Article 6. There may be exceptions to the application of the principles and rules of this Code in cases where a rigid application would lead to great confusion. Such exceptions become valid when approved by the Nomenclature Commission

E SUGGESTIONS AND RECOMMENDATIONS OF THE SOCIETY OF AMERICAN BACTERIOLOGISTS

The preliminary report of the Committee of the Society of American Bacteriologists on Characterization and Classification of Bacterial Types (Winslow, Broadhurst, Buchanan, Krumwiede, Rogers and Smith 1917) suggested the following to the Society

We recommend to the Society of American Bacteriologists

a. That the International Rules for Botanical Nomenclature be accepted by the Society as governing bacterial terminology, with the exception that French, English or German may be substituted for Latin in the diagnosis

b. That the date of publication of the third edition of Zopf's *Spaltpilze* be considered as the date for the beginning of bacteriological nomenclature for the purpose of determining priority, with the exception of a list of genera *conservanda* to be adopted by the Society at its 1918 meeting

c. That the Society take steps to present these recommendations to the next International Botanical Congress, and if possible to secure favorable action thereon by that body

In the final report (1920) the following statement is made

The following names are recommended for adoption as approved genera

<i>Acetobacter</i> Fuhrmann	<i>Leuconostoc</i> Van Tieghem
<i>Actinomyces</i> Harz	<i>Micrococcus</i> Cohn
<i>Bacillus</i> Cohn	<i>Rhizobium</i> Frank
<i>Bacterium</i> Ehrenberg	<i>Sarcina</i> Goodsir
<i>Chromobacterium</i> Bergonzini	<i>Spirillum</i> Ehrenberg
<i>Clostridium</i> Prazmowski	<i>Staphylococcus</i> Rosenbach
<i>Erythrobacillus</i> Fortineau	<i>Streptococcus</i> Rosenbach
<i>Leptotrichia</i> Trevisan	<i>Vibrio</i> Mueller

Its work so far as possible being completed, we recommend that the Committee on Characterization and Classification of Bacterial Types be discharged and that a new Committee on Bacterial Taxonomy be appointed (1) to study and report to the Society from time to time in regard to problems of nomenclature, including such revisions of the nomenclature in the present report as may seem necessary, and (2) to take the proper steps to secure action at the next International Botanical Congress leading to the general ends contemplated in the 1916 recommendations of the Society

a. That French, English or German may be substituted for Latin in the diagnosis of bacterial species

b. That the date of publication of the third edition of Zopf's Spaltpilze (1883) be considered the beginning of bacterial nomenclature for the purpose of determining priority, with the exception of a definite list of genera conservanda

c. That such of the approved generic names specified above as may be found to require such action be recognized as genera conservanda in bacterial taxonomy

STATEMENT AS TO NOMENCLATURAL PRACTICE IN PRESENT VOLUME

It would seem evident from the preceding discussion that the fundamental basis for nomenclature in bacteriology must be the International Botanical or Vienna-Brussels Code. However, it is quite apparent that there are certain rules which have not been and cannot well be recognized by bacteriologists, particularly the rule requiring that the specific diagnoses be published in Latin in order to be valid. It is also evident that stability in bacteriological nomenclature must come through the application of a type basis. Apparently the best that has been formulated for botanists is that contained in the majority report of the Committee on Botanical Nomenclature given above. We shall therefore utilize the latter as a basis for naming the bacteria described. It should be remembered that acceptance of the code of this report does not mean abandonment of the International Code.

GENERAL SYSTEMATIC BACTERIOLOGY

Occasional reference will be made to the Zoological Code where organisms have a doubtful status in the plant kingdom. It will also be used to determine good usage where the Botanical Code is silent or ambiguous.

CHAPTER III

NOMENCLATURAL STATUS OF NAMES WHICH HAVE BEEN APPLIED TO GROUPS OF BACTERIA OF HIGHER RANK THAN SPECIES

On the following pages is given alphabetically arranged a list of the names which have been proposed for subgenera, genera, subtribes, tribes, subfamilies, families, orders, and classes of bacteria, including casual or vernacular terms which have had more or less extensive recognition and have been confused with scientific names. In general an attempt has been to give the original diagnosis of the term, to discuss its usage by bacteriologists, and to point out whether its use apparently contravenes good usage or the rules of the code. In many cases an opinion is given as to probable validity.

Acetimonas. A generic name for the vinegar bacteria proposed by S. Orla-Jensen (1909, p. 312). This genus is listed as the fourth of the family *Oxydobacteriaceae*, order *Cephalotrichinae*. The genus is first described as follows:

Die Essigsäure bakterien sind, soweit bekannt, ebenso wie die vorhergehenden Bakterien monotrich oder unbewegliche Stäbchen, die sich auf der Oberfläche der Nährflüssigkeiten zu einer Haut ausbreiten. Ich habe ihnen daher den Gattungsnamen *Acetimonas* gegeben. Von den Schnellwüchsigen bis zu den Essigbildnern des Weines und des Bieres steigen die Aussprüche auf organische Nahrung. Bekanntlich gedeihen die Schnellwüchsigen (z. B. *Schützenbachii*) mit Äthylalkohol als einzige Kohlenstoffquelle. Es ist jedoch kaum wahrscheinlich, dass Alkohol die erste Kohlenstoff- und Energiequelle war, die den Essigsäurebakterien hier auf der Erde zur Verfügung stand, dies sind eher Kohlenhydrate gewesen, welche sie ja auch in höherem oder geringerem Grade zu oxydieren im Stande sind. Nach meinen Untersuchungen können die meisten Essigsäurebakterien sich mit Salpeter als Stickstoffquelle begnügen. Einzelne Arten (am stärksten *B. aceti*) reduzieren nach längerer Zeit kleinere Mengen Salpeter bis zum Ammoniak, eine Eigenschaft die auch andere obligat aerobe Bakterien (z. B. *Azotobacter* und gewisse Heubacillen) besitzen, und die zeigt, dass solche Bakterien ausser dem freien Sauerstoff auch gebundenen verwenden können.

Later the same author (1909, p. 477) gave the following description: "Bedürfnis in der Regel nicht organischen N-Quellen. Sauerstoffbedürfnis positiv. Oxydiert Alkohol zu Essigsäure."

It will be noted that no type species is designated. Of the two species mentioned "*B. Schützenbachii*" is given first, and may be regarded as the type.

Other names which have been applied in a more or less strict generic sense to the group of acetic bacteria are *Mycoderma* (Persoon 1822, p. 96), *Ulvina* (Kützing 1837 ?), *Umbina* (Nageli 1857, p. 760), *Acetobacter* (Beijerinck 1898 ?), *Acetobacterium* (Hoyer 1898, p. 870).

The organisms belonging to this group have commonly been placed in the genera *Bacillus* and *Bacterium* by various authors.

The name *Acetimonas* was regarded as a synonym of *Mycoderma* Persoon by Buchanan (1918, p. 45) and Winslow *et al.* (1917, p. 551), and finally of *Acetobacter* Furhmann by Winslow *et al.* (1920, p. 201).

It is here included as a synonym of *Acetobacter* Beijerinck, *q. v.*

Acetobacter. A generic name proposed by Beijerinck (?) for the group of acetic acid bacteria.

The exact time and method of publication is difficult to determine.

According to Enlows (1920, p. 11), the name is first found in Kral's *Sammlung v. Mikroorg.* (Prague 1898, pp. 7 and 8) where *Acetobacter pastorianus* (Hansen ?), Beijerinck and *A. aceti* are listed as synonyms of *Bacterium Pasteurianum* and *Bact. aceti* respectively. In the 1900 and 1904 editions these are likewise listed as synonyms of *Bacterium Pasteurianum* and *Bact. aceti* respectively.

Beijerinck (1901, p. 219), in an article in which he describes several genera of lactic bacteria at length, mentions *Acetobacter* as producing catalase, in contrast to *Lactobacillus* which produces none.

It is of interest to note that in a translation (published in 1898) of a Dutch paper (published in 1897) on the acetic bacteria, Beijerinck uses the name *Bacterium* consistently for the species of this group.

Ludwig (1898, p. 870) in a review of Hoyer's (1898) dissertation on the acetic bacteria uses the name *Acetobacterium xylinum*. Examination of the original is necessary to show whether this is a misprint.

The first adequate description of an organism placed definitely in this genus thus far found is *Acetobacter phicatum* Furhmann (1905, p. 8). It was this that led the Committee of the Society of American Bacteriologists to ascribe the genus to Furhmann. If this is done, however, his species *A. phicatum* would constitute the type.

Acetobacter melanoginum was described later by Beijerinck (1911, p. 171).

The generic name *Acetobacter* was accepted by Winslow *et al.* (1920, p. 201), for the acetic bacteria, displacing *Mycoderma*. They give the following description:

Cells rod shaped, frequently in chains, non-motile Cells grow usually on the surface of alcoholic solutions as obligate aerobes, securing growth energy by the oxidation of alcohol to acetic acid Also capable of utilizing certain other carbonaceous compounds, as sugar and acetic acid Elongated, filamentous, club shaped, swollen and even branched cells may occur as involution forms

The type species designated by Winslow *et al* was *Acetobacter aceti* (Thompson 1852) Committee

It is probable that the correct designation of the type species would be *Acetobacter aceti* Beijerinck

Other generic names which have applied more or less strictly to this group are *Mycoderma* (Persoon 1822, p 96), *Ulvina* (Kutzing 1837 ?), *Umbina* (Nageli 1857), *Acetobacterium* (Ludwig ? 1898) and *Acetimonas* (Orla-Jensen 1909) *q v*

Buchanan (1918, p 45) listed *Acetobacter* as a synonym of *Mycoderma* Bergey *et al* include this genus as the sixth of the tribe *Nitrobactereae*

In this treatise *Acetobacter* is recognized as a valid generic name with *Acetobacter aceti* Beijerinck as the type

Acetobacterium. A generic name apparently first used by Hoyer (1898) as a designation for acetic bacteria The original thesis is not available, but the name *Acetobacter xylinum* is used by Ludwig (1898, p 870) in an extended review of Hoyer's paper He also in the same year apparently ascribed the name to Beijerinck in the following statement

Nach den Beobachtungen Beyerinck's wurde der *Leuconostoc Lagerheimi* mit dem *Bacterium xylinum* Brown zusammen in ein und dieselbe Gruppe der Acetobacterien geboren Beyerinck bezeichnet ihn in lit als *Acetobacterium xylinum* var *Lagerheimi*

The only reference to such a conception thus far found in the writings of Beijerinck is a footnote (1898, p 211) to a brief description of *Bacterium xylinum* in which he says, "Hauptmasse des Schleims beim Schleimflusse lebender Eichen bildet." There is no use of the name *Acetobacterium*

It is probable that *Acetobacterium* should be regarded as a synonym of *Acetobacter*

Achromatiaceae A family of the order *Thiobacterales* proposed by Buchanan (1918, p 462) to include the three genera *Achromatium*, *Thiophysa* and *Hillhousia*

The diagnosis given is "Unicellular, large, motile (by means of flagella ?) cells containing granules of sulphur (or in one form possibly

oxalate), but no bacteriopurpurin" The family is recognized by Bergey *et al* (1923, p 410) in this sense

The generic type is *Achromatium oxalisferum* Schewiakoff (1893, p. 1)

Achromatium. A genus of schizophytes proposed by Schewiakoff (1893, p 1) and tentatively placed by him with the bacteria

The species described, *Achromatium oxalisferum*, was found in river slime, by R Lauterborn, in considerable numbers In form the organism varies from nearly spherical in newly divided cells to ellipsoidal In size the organism is larger than any other form previously described among bacteria, cells about to divide are 15 to 43 μ in length, diameter 9 to 22 μ Occasionally longer cells are found. The cells are closely packed with large granules which were at first interpreted as composed of sulfur. The use of various solvents and microchemical tests indicated them to be, however, of calcium oxalate When the granules are dissolved the cell is found to contain protoplasm of a coarse alveolar structure The cells are motile, but flagella were not demonstrated Cell division is described as the result of constriction, as among flagellates, rather than by the fission characteristic of most bacteria

Migula (1900, p 1038) tentatively added *Achromatium Mulleri* (Warming) Migula to this genus The organism was originally described by Warming (1876) as *Monas Mulleri* The descriptions make it probable that this organism is misplaced in *Achromatium*, and belongs with the sulfur forms Migula gives the following description of the genus

Zellen gross, eiförmig cylindrisch mit grobkörnigem Inhalt, der jedoch nicht aus Schwefel, sondern aus Oxalsäure und einem Calciumsalz besteht Bewegung vorhanden, doch ist eine Geissel als Bewegungsorgan nicht nachgewiesen Teilung von der Bakterien abweichend, durch Einschnürung, wie bei den Flagellaten

Achromatium oxalisferum Schewiakoff was subsequently described independently by Frenzel (1897, p 801) as *Modderula hartwigii* (from the name "Modder" given by North German fishermen to the slimy river bottoms) Lauterborn (1898, p 95) called attention to the previous description of *Achromatium*, and agreed with Schewiakoff in placing the organism with the bacteria Frenzel found this form to vary from 12 to 50 μ in length He was able to find no adequate explanation of the motility The cell membrane he describes as firm, almost doubly contoured, clear, colorless and structureless He interpreted the granules contained within the cell as sulfur The organism was not cultivated in artificial media, although it continued to

multiply for a time in the laboratory. He concluded that the organism is most closely related to the bacteria, but that it constitutes a very distinct type.

It would appear that the genus *Achromatium* Schewiakoff is valid, with the type species *A. oxalisferum* Schewiakoff. It is listed by Smith (1905, p. 174) as a doubtful bacterial genus and in another list as a rejected name. Migula (1904, p. 128) states that he appends the genus *Achromatium* to the bacteria, although (because of the cell dimensions) he considers the relationship distant. A careful rechecking of the work of Schewiakoff on the question of the composition of the granules would be helpful in determining the affinities of this organism.

Buchanan (1918, p. 462) included this genus as the type of the family *Achromatiaceae* with the following description:

Cells large, nearly spherical in newly divided cells to ellipsoidal, 15 to 43 by 9 to 22 μ . Cells closely packed with large granules, at first interpreted as sulphur, but later interpreted as calcium oxalate. When granules are dissolved cells show coarse alveolar structure. Cells are motile, flagella not demonstrated. Cell division resembles the constriction of flagellates rather than the fission characteristic of bacteria.

This generic diagnosis was also used by Bergey *et al.* (1923, p. 410).

In this treatise *Achromatium* is accepted as a valid bacterial genus.

Achromobacter. A generic name proposed by Bergey *et al.* (Committee Soc. Am. Bact., 1923, p. 132) for the single genus of the tribe *Achromobacteriae*. The generic description is:

Non-pigment forming (at most no pigment formed on agar or gelatin) rods, occurring in water and soil. Motile or non-motile. Gram-negative.

The type species is *Achromobacter liquefaciens* (Frankland).

In all some 51 species are included in the genus.

Achromobacteriae. A name used by Bergey *et al.* (1923, p. 132) for the second tribe of the family *Bacteriaceae*. The tribal description is:

Rods, small to medium in size, occurring principally in water and soil. Form no pigment on agar or gelatin but may produce a brownish growth on potato. Cultural characters variable. Motile or non-motile. Gram-negative.

One genus only is included, *Achromobacter*.

Acidobacteriaceae. The first family of the order *Peritrichinae* of Orla-Jensen (1909, p. 343). It includes seven genera: *Denitrobacterium*, *Bacterium*, *Propionibacterium*, *Caseobacterium*, *Streptococcus*, *Micrococcus* and *Sarcina*. It includes organisms which usually cause acid

formation from carbohydrates, which when motile, have peritrichous flagella, which are not obligate anaerobes and which do not typically form alkalies by splitting of proteins. The genus *Denitrobacterium*, characterized by large denitrifying powers, does not strictly belong in this family, but is intermediate between the other genera and the family *Luminobacteriaceae* of the preceding order *Cephalotrichinae*.

The genera of the family *Acidobacteriaceae* as recognized by Orla-Jensen may be differentiated by the following key

- | | | | |
|---|--------------------------------|-------------|--------------------------|
| A | Primarily denitrifying rods | Genus 1 | <i>Denitrobacterium</i> |
| B | Not primarily denitrifiers | | |
| | I Cells rod shaped | | |
| | a Producing succinic acid | Colon group | |
| | | Genus 2 | <i>Bacterium</i> |
| | b Producing lactic acid | Genus 4 | <i>Caseobacterium</i> . |
| | II Cells spherical. | | |
| | a Producing propionic acid | | |
| | | Genus 3 | <i>Propionibacterium</i> |
| | b Not producing propionic acid | | |
| | 1 Cells in chains | Genus 5 | <i>Streptococcus</i> . |
| | 2 Cells irregularly arranged | | |
| | | Genus 6 | <i>Micrococcus</i> |
| | 3 Cells in packets | Genus 7 | <i>Sarcina</i> |

Orla-Jensen (p 340) suggests that the additional genera *Sporosarcina* for sarcina forms that develop spores and *Peptonococcus* for the lactic acid bacteria that form peptones from proteins should be added

The name apparently has not been used by subsequent writers

It may be noted that the family name *Acidobacteriaceae* is not derived from any of the component genera, and is therefore incorrect in form, and may be regarded as invalid

Actinobacille. A casual name given by Lignières and Spitz (1904, p 454) to a group of bacteria related to *Actinomyces*. The French designation *actinobacille* was used apparently as a translation of the original Spanish term *actinobacilo* (q v) by Lignières and Spitz (1902, p 169). This is the form later designated by the generic name *Actinobacillus* (q v) by Brumpt (1910, p 849).

The description given by Lignières and Spitz is as follows

L'agent spécifique de l'actinophytose à actinobacille se présente, dans les cultures sous l'aspect d'un microbe parfois à peine plus gros que celui du choléra des poules. Dans les premières cultures, il est nettement bacillaire, plus tard, surtout sur gélose, il affecte la forme du cocco-bacille ou de diplocoque, dans les cultures en bouillon sérum, on observe l'aspect streptobacillaire. Enfin,

dans les vieilles cultures, principalement en bouillon additioné de sérum agglutinant, on trouve souvent des formes d'invololution les plus bizarres. Jamais on ne trouve de filaments dichotomisés comme les autres actinophytoses déjà signalés 101

Enlows (1920, p 11) gives the following characterization

Pleomorphic Sometimes rodlike, sometimes coccuslike, in pairs, also streptobacillary forms occur 0.15 μ to 1.25 μ in length by 0.4 μ , non-motile no spores, gram-negative, bipolar staining The final stage of growth gives rise to little masses, in which the organisms are pressed closely together, giving the ray like aspect These masses consist of a central germinative zone and an outer or vegetative zone Cause of actinobacillosis in cattle

This is to be regarded as a casual name for the genus *Actinobacillus* *Actinobacillus*. A generic name given by Brumpt (1910, p 849) to the organism causing "Actinobaclosis" and termed "Actinobacilo" by Lignières and Spitz (1902, p 169) The organism was accorded no specific name by Lignières, it was named by Brumpt in his honor, *Actinobacillus Lignieresii* This species therefore constitutes the generic type

The organism is the cause of an infection in cattle and sheep termed actinobacillosis The lesions resemble closely those of actinomycosis In culture of the organism on solid media small, transparent colonies appear in twenty-four hours They consist of small diplococcus-like bacilli or of streptobacilli, resembling the fowl cholera organism when grown in broth in their occurrence in long chains Dextrose and lactose are not fermented Indol reaction is weak The pus in the lesions contains granules the size of a millet seed Microscopically these show at their peripheries swollen, finger shaped, branched forms whose central ends are pointed The filaments are Gram-negative The central portion does not contain a branched mycelium Isolated threads are not found in the pus

Enlows (1920) states "Type species (monotypy) *A. lignieresii* Streptobacillary, bacillary or coccoidal in form Gram-negative Never filamentous"

Merrill and Wade (1919, p 64) state that this name was given to a supposed subtype of *Discomyces*, but that the distinction has not been recognized, and by most authors the name is considered a synonym

Buchanan (1918, p 404), Winslow *et al* (1920, p 198) and Bergey *et al* (1923, p 338) recognize the genus with the following diagnosis.

Filament formation, resembling streptobacilli In lesions no mycelium is formed, but at peripheries finger shaped branched cells are visible Gram negative Not acid fast

It is included in the family *Actinomycetaceae*

Bejerinck (1914, p 96) renamed his *Bacillus oligocarophilus* as *Actinobacillus oligocarophilus* and his *Actinomyces (Streptothrix) paulotrophus* as *A. paulotrophus*. No indication is given of the previous use of this generic name by Brumpt. The organisms described are probably sufficiently unlike those causing actinobacillosis in cattle to prevent their inclusion in the same genus. Probably Bejerinck's *Actinobacillus* is to be regarded as an invalid homonym. His description of the genus and the two species follows

Drittens, die Gattung *Actinobacillus*, welche zu der Familie der Actinomyceten gehört und deshalb typisch unbeweglich ist. Unterscheidet sich von der Neumann und Lehmann aufgestellten Gattung *Mycobacterium* durch das vollständige Fehlen der Verzweigung, sodass man nur Stäbchen oder Fäden findet. Erzeugt die charakteristische treibende "Kahmhaut" auf der Oberfläche der nitrifizierenden Flüssigkeiten. Darin finden sich zwei Arten nämlich

a *Actinobacillus oligocarophilus* welche sich ernähren kann von den Kohlenstoffverbindungen der atmosphärischen Luft und dann das "Kahmpilzmerkmal" zeigt, andererseits auf den verschiedensten organischen Nährboden wächst ohne das "Kahmpilzmerkmal". Letzterer Zustand auf Kieselplatten zurückgeimpft zeigt das "Kahmpilzmerkmal". Doch kann das Merkmal verloren gehen durch lange fortgesetzte saprophytische Lebensweise. Verflüssigt Nährgelatine nicht.

b *Actinobacillus paulotrophus*. Erzeugt auf den nitrifizierenden Platten schimmelartige Kolonien mit "Lufthyphen," besteht jedoch mikroskopisch anscheinend aus gleichartigen Stäbchen und Fäden. Wächst durchaus nicht bei Gegenwart organischer Substanz.

Viertens, die Gattung *Actinomyces* wovon verschiedene Arten, in geringer Anzahl in den "Kahmhauten" von *Actinobacillus* vorkommen können, daraus jedoch bei den Ueberimpfungen bisweilen gänzlich verschwinden.

If generic rank is to be accorded organisms of this type, the name *Actinobacillus* Brumpt would appear to be valid, otherwise it should be suppressed as a synonym. Organisms of this type have probably been included by various writers under the generic names *Actinomyces*, *Bacterium* and *Bacillus*.

It is possible that *Actinobacterium* Haass should be regarded as an invalid synonym.

In the present volume the generic name *Actinobacillus* is accepted in the sense of Brumpt.

Actinobacilo. A Spanish casual name first used by Lignières and Spitz (1902, p 169) for the organisms later named *Actinobacillus* (q v).

Actinobacter. A generic name for certain milk bacteria proposed by Duclaux (1882, p 110).

The first species described (the type species) is *Actinobacter polymorphum* Duclaux. In milk the organism occurs as rods 2 to 3μ in length, each surrounded by a hyaline capsule 5 to 6μ in length. At first the milk retains its normal appearance, and gradually becomes gelatinous and at last viscous. The opacity is in part lost, the transparency increases, the coagulum settles out and disappears. Finally the transparent liquid shows little more viscosity than the normal milk. In the later stages the organisms are 2 to 3μ long by 1μ in diameter and without capsule. In Liebig's bouillon there are no capsules, the cells may attain a length of 8 to 10μ . In a glycerine medium the capsules are present. The capsule does not disappear as in milk, "sans doute parce que le liquide reste acide." In sucrose medium the growth and appearance is much as in glycerine. Gas (carbon dioxide and hydrogen) were formed from sugar, but not from glycerine.

Maggi (1886, p. 174) includes *Actinobacter* as a valid genus. It appears evident from the characterization of this organism that it is a member of the colon group of bacteria, probably closely related to organisms of the type of *Bacterium lactis aerogenes*. The description is too indefinite however to allow of accurate diagnosis at the present time. It is included by Lohnis (1911) with the *Bacterium pneumoniae* group. It is rejected as invalid by E. F. Smith (1905, p. 174).

Enlows (1920, p. 12) characterizes the genus as follows

Type species (monotypy) *A. polymorphus*. Polymorphic. Non-motile, very thin small rods 2 to 3μ long, surrounded by a hyaline, oval or round, gelatinous envelope 5 or 6μ long. The organism gradually passes over into very short cylinders not more than 1μ long, which are not capsulated. No capsule ever forms in Liebig's bouillon, where the organism may reach 10μ in length. Aerobic. Multiplication by transverse division. Transforms the casein of milk into a water-soluble albuminoid, and milk-sugar into alcohol and acetic acid. Found in milk.

If the group of colon bacilli or colon-typhoid bacilli are to be accorded generic rank the name *Actinobacter* must be considered as one which might be used in its designation. It is questionable however whether the type species can ever be identified from the description, and probably it is unwise to use as a genus name one in which the type species is so uncertain. Other bacteria have apparently not been placed in the genus by other authors. It is evident therefore that without emendation this name can scarcely be regarded as valid. It may be listed as a synonym of *Bacterium* (*q. v.*).

Actinobacterium. A generic name proposed by Haass (1906, p 180) to apply to two strains of organisms termed by him *Actinomyces* Klinik I and II which were found to differ from all of eleven other types studied. These were isolated from pus from cases of human actinomycosis. They were similar to the forms described by Wolff and Israel, by Silberschmidt and by Wright as anaerobes. The organisms had short cells, somewhat branched, and were Gram-positive. Haass notes that Lignières and Spitz previously used the generic name *Actinobacillus* for a similar type. He believes the genus to be intermediate between *Actinomyces* and *Corynebacterium*. Sampietro (1908, p 331) described an *Actinobacterium israeli* var. *spitzi*.

No species of the genus was named by Haass, if valid therefore it should be credited to Sampietro. It may perhaps be regarded as a synonym of *Actinomyces*.

Reitz (1906, p 731) apparently confused this name with *Actinobacter* (*q v*) and refers to *Actinobacterium lactis viscosum*.

Actinocladothrix. A generic name ascribed to Affanassieff by Levy (1899, p 2), by Schlegel (1913, p 303), and by others. It seems probable that they are in error in giving the name generic rank. Affanassieff (1888, p 79) himself gave the name *Bacterium actinocladothrix* to an *Actinomyces*. De Toni and Trevisan (1889, p 928) also give the synonym *Bacterium actinocladothrix*. Afanassjew 1888 to *Nocardia Actinomyces* Trevisan (*Actinomyces bovis* Harz). Brumpt (1910, p 844) gives the name the same specific form. The use of the name *Actinocladothrix* as a generic designation is therefore based upon false premises. The correct generic designation would seem to be *Actinomyces*.

Actinococcus. A name proposed by Kützing for a genus of algae, included by Pfeiffer (1870, p 11) in the tribe *Chaetophorae*.

Beijerinck (1914, p 196) suggested the generic name *Actinococcus* to apply to a bacterial species *A. cyaneus*. Beijerinck isolated this organism from garden soil on a medium particularly adapted to the culture of *Actinomyces*, containing glucose, calcium malate, ammonium sulphite, dipotassium phosphate and agar. Gelatin was liquefied slowly, and agar made alkaline by the formation of ammonium carbonate. In a medium with peptone the color is rose, with glucose and nitrate of potassium or ammonium it remains violet. Ammonium acetate is very favorable for the production of the violet color. The pigment is blue with alkalis and red with acids. The organism he believes belongs with the genera *Corynebacterium*, *Mycobacterium* and *Actinomyces*. This relationship is indicated by the dry colonies, the disposition of the cell

in radiating rays, and the branched appearance of the rays. Later (1914, p. 99) he mentions the creation of the generic name.

It would seem that the prior use of *Actinococcus* for a genus of algae would invalidate its use for a genus of bacteria.

Actinomyce. A generic name proposed by Meyen (1827, p. 442) for a fungus species *Actinomyce Horkelii*. The genus was placed in the new group *Hydrotremellinae*. The generic description follows:

Sporodochia, cellulis hyalinis simplicibus, enormiter et multipliciter ramificantibus sporis impletis, substantiae uniformi gelatinosa hyalina induta.

This generic name is of importance in bacteriology only because it was given prior to the use of *Actinomyces* by Harz for an entirely distinct organism. In many references the spelling is made identical, but it will be noted that there is a difference. For example Streinz (1862, p. 3) in his "Nomenclator fungorum" writes "*Actinomyce* Meyen (Stiahlenpilz) eine neue Pilzgattung."

Perroncito (1879, p. 35), Musgrave, Clegg and Polk (1908, p. 458), Merrill and Wade (1919, p. 65), Chalmers and Christopherson (1916, p. 227) and others have urged that this use precludes the validity of *Actinomyces* Harz, *q v*.

It is probable that *Actinomyce* Meyen should not render invalid *Actinomyces* for the following reasons:

1. The spellings are different. The botanical code expressly provides that names differing only by a single letter may both be valid if both do not come within the same family, and are not likely to give rise to confusion.

2. No other author apparently has used *Actinomyce* Meyen, and it may be said to be "universally regarded as invalid."

Actinomyces. The generic name *Actinomyce* was used by Meyen (1827, p. 442) for a genus of fungi belonging to the *Hydrotremellinae* Meyen, with one species *A. Horkelii* Meyen. For reasons why this should not be regarded as invalidating the subsequent use of *Actinomyces*, see *Actinomyce*.

In 1877 the name *Actinomyces* was applied to a genus of thread bacteria by Harz (1877, p. 125). The material containing the organism was sent to Harz by Bollinger. It consisted of granules secured from pus from cattle affected by "lumpy jaw." The granules were found to consist of slender filaments, irregularly branched, radiating from the center, and with the ends of the filaments in the form of refractive swellings. The name *Actinomyces bovis* was given it.

In the following year, Rivolta (1878, p. 201) used the generic designation *Discomyces* (q v). The name *Actinomyces bovis* Harz, however, was recognized by Rivolta in 1884. At this time he differentiated between *A. bovis* and two other species which he termed *Discomyces equi* and *D. pleuriticus*.

Actinomyces was used as a generic name by several authors during the next decade. Perroncito (1879, p. 35) noted that the name *Actinomyces* should be changed because of the previous use by Meyen of *Actinomyce* for one of the *Coniomyces*. Hertwig (1886, p. 365) described an *Actinomyces musculorum*, which was further studied by Duncker (1887, p. 224). Schroeter (1886, p. 174) recognized the genus *Actinomyces* Harz, regarding *Discomyces* Rivolta as a synonym. He gives the following generic description:

Faden sehr dünn, etwa bis 0.5 mm lang, um einen festen Kern, welcher in den Faden durchgezogen wird, strahlig ausgebreitet, mit unregelmässig abgehenden Aesten, an den Enden in keulenförmige, stark lichtbrechende Auschwüelungen übergehend, welche Gliederung zeigen.

Trevisan (1889, p. 927) recognized the fact that an organism named by Cohn (1875, p. 186) *Streptothrix Foersteri* was closely related to the *Actinomyces bovis* Harz. He concluded that neither *Actinomyces* Harz, nor *Streptothrix* Cohn were tenable, the former because of prior use by Meyen (l. c.) and the latter by Corda (1839, p. 23). He accordingly named the genus *Nocardia*, with *Nocardia Actinomyces* Trevisan as one of the five species. Sauvageau and Radais (1892, p. 242) suggested as a result of comparing the genus *Streptothrix* of Cohn with other known *Hyphomycetes*, that this genus should be merged with the *Oospora* of Wallroth (1837, p. 182). Several authors have used this generic name. Lindau (1910, p. 720) as recently as 1910 reaches the conclusion that this form is most closely related to *Oospora*.

The name *Actinomyces* has been used as a generic designation by many authors. Ludwig (1892, p. 3) defined the genus as having branched filaments, the branches swollen at the tips. Terni (1894, p. 79) noted the previous use by Corda of *Streptothrix* and accepted *Actinomyces*. Gasperini (1895, p. 82) concluded that it is valid and should replace *Streptothrix* Cohn, *Discomyces* and *Pleurimices* Rivolta, *Bacillus* Nocard, *Micromyces* Gruber and *Oospora* Sauvageau and Radais. Lachner-Sandoval (1898) made a careful study of the nomenclature of the genus and concluded the right generic designation to be *Actinomyces*. Levy (1899, p. 1) confirmed his conclusions. Berestnew (1899, p. 390) concluded that on botanical grounds the name is tenable,

and that it is a most appropriate designation both because it indicates a most important characteristic of the colonies on artificial media and because it shows relationship to the fungi. Lehmann and Neumann (1896) in the first edition of their *Grundriss* use the name *Oospora*, but in later editions they have followed Lachner-Sandoval in the use of *Actinomyces*.

Geddoelst (1902) concludes *Actinomyces* to be invalid, and substitutes for it Rivolta's term, *Discomyces*. This point of view is also taken by Moore and Wade (1919, p. 55).

Wright (1904, p. 349) concludes that the name *Actinomyces* should be strictly limited to the parasitic forms which produce rays in tissues, and do not form chains of conidia or spores on artificial media. For the forms which do not show the clubbing in tissues and which develop chains of conidia he used the name *Nocardia* Trevisan as emended by Blanchard (1896, p. 853). Jordan (1910, p. 411) used the same basis of differentiation. The genus *Actinomyces* has also been recognized by Orla-Jensen (1909, p. 334) and by Schlegel (1913, p. 302). The latter author emphasized the formation of a true branched mycelium, reproduction by conidial chains, and by the presence of the rayed form in the body tissues. Petruschky (1913, p. 270) gives a very confused discussion of this nomenclature. He concludes that *Actinomyces* should be strictly limited to the clubbed or ray forms in tissues, and that others should be termed *Streptothrix*. Pinoy (1911) introduced the new generic name *Cohnistreptothrix* for organisms of the type of Cohn's *Streptothrix foersteri*. In a later article (1913, p. 929) he reviews the use of the various generic names proposed, and concludes that *Streptothrix* must be reserved for the hyphomycete of Corda, and that *Actinomyces* is equally untenable. *Discomyces* he states was not proposed originally as a strict generic designation, but as a common name and furthermore that it resembles in form too closely the group name *Discomycetes* of the *Pezizales*. He quotes the action of the botanical section of the first International Congress of Comparative Pathology, which decided to adopt the generic name *Nocardia* of Trevisan. He divides the old genus *Actinomyces* into two, *Nocardia* with *N. bovis* as the type, and *Cohnistreptothrix* with *C. foersteri* and *C. israeli* as the best known species. *Nocardia* includes those forms which are aerobic, grow readily on artificial media and produce arthrospores. *Cohnistreptothrix* on the other hand, includes those forms which are anaerobic or at least microaerophilic, are relatively more difficult of cultivation and do not produce arthrospores. This author comes to the conclusion

that the presence or absence of "clubs" when growing in tissues is not diagnostic. These are the result of certain interreactions between parasite and host, and may be produced in either genus under the right conditions.

Buchanan (1918, p. 405) characterized the genus as follows:

Branched filaments, resembling mycelium, breaking up into segments which may function as conidia. Usually parasitic. Clubbed ends conspicuous in lesions. Not producing aerial hyphae or conidia.

The type species is *Actinomyces bovis* Harz, the cause of bovine actinomycosis.

Castellani and Chalmers (1919, p. 1041) reject *Actinomyces* in favor of *Nocardia*. Waksman (1919) in his monographic treatment of the group accepts the generic name *Actinomyces* as does also Lieske (1921, p. 5).

The description used by the "Committee" of the American Society of Bacteriologists (1920, p. 198) and by Bergey *et al.* (1923, p. 339) follows:

Organism growing in form of a much-branched mycelium, which may break up into segments that function as conidia. Sometimes parasitic, with clubbed ends of radiating threads conspicuous in lesions in animal body. Some species are microaerophilic or anaerobic. Non-motile.

The type species is *Actinomyces bovis* Harz.

Merrill and Wade (1919, p. 63) insist upon the non-validity of *Actinomyces* and the validity of *Discomyces*.

The claim of legitimacy for *Actinomyces* as a generic name for the organism causing lumpy jaw and other similar affections is menaced by two facts. First, the name *Actinomyces* was used by Meyen in 1827 for an entirely distinct fungus. Second, the generic name *Streptothrix* Cohn has priority. Probably the best analysis of the situation is that which has been furnished by Breed and Conn (1919, p. 585, 1920, p. 489). They say (1919, p. 594):

Two attempts have already been made to establish the validity of a generic name for this group of organisms through legislative action both of which must be regarded as abortive. The first was taken by a Committee of the English Society of Pathologists who, as reported by Foulerton (1912, p. 304), approved the term *Streptothrix* Cohn in ignorance of the general and wide usage of *Streptothrix* Corda. The second was taken by the Botanical Section of the First International Congress of Comparative Pathology according to Pinoy (1913, p. 933) and confirmed by him in a letter dated September 9, 1918. The name ac-

cepted was *Nocardia*. The official record of the Congress however contains no reference to this action which is stated to have been taken during the discussion of a paper by Potron (1912). As this action ignores the stronger claims of *Actinomyces* and *Discomyces*, and does not appear in the official record, it cannot be regarded as final.

And further

Because of confusion between *Streptothrix* Corda 1839 and *Streptothrix* Cohn 1875 and the general use of the former term by mycologists, the latter term should be generally disregarded. According to the International Rules of Botanical Nomenclature, the limited use of the term *Actinomyce* by Meyen in 1828 and 1832 is not sufficient to invalidate the generally used *Actinomyces* Harz 1877 provided the latter is accepted as a *genus conservandum* by an International Botanical Congress. The continued use of the latter term is therefore recommended. The type species of the genus is *A. bovis* Harz.

Later they note that both botanical and zoological codes provide for the retention of generic names which differ only by a single letter (International Botanical Code, Chapter III, Article 57). They note further etymological differences between *Actinomyce* and *Actinomyces*, (1920, p. 490).

Although the stem words from which *Actinomyce* and *Actinomyces* are derived have identical meaning in the original Greek, "myce" is derived from the less commonly used feminine word, *μυκη*, while "myces" comes from the masculine noun, *μυκης*. Thus the two generic terms in question ought not to be regarded as homonyms as is done by Merrill and Wade. This view we find has already been expressed by Gussow (1914) in a paper which we had overlooked, and is confirmed by those authorities with whom we have consulted. Thus being the case legislative action by an International Congress is unnecessary. *Actinomyces* Harz is valid without such action and should be retained rather than *Discomyces* Rivolta.

The generic name *Streptothrix* Cohn which has been held by some writers to have priority over *Actinomyces* is clearly invalid. Corda's genus *Streptothrix* is well characterized both by description and figure, it has had additional species described, and is generally recognized in classifications of the *Hyphomycetes*. The genus *Oospora* Wallroth includes forms which are quite unrelated to *Actinomyces*, these forms should therefore not be ascribed to it.

A real difficulty also presents itself when an attempt is made to pick the type of the genus. It would seem certainly that to it the name *Actinomyces bovis* should be given. Harz did not publish cultural characters. It is impossible to determine with certainty just what his

species was Bostroem cultivated an aerobic type which produced spores Wolf and Israel described an anaerobic asporogenous type. Wright and Pinoy have both shown the latter to be far more common. The probabilities are that it was this form that was observed by Harz. It would seem logical therefore to apply to it the name *Actinomyces bovis*. Pinoy (who accepts *Nocardia* instead of *Actinomyces*) insists that the organism described by Bostroem, i.e., the aerobic spore producing form be taken as the type. Wright states, "I think that the generic name *Actinomyces* should be restricted to microorganisms with the biological characters described in this paper (i.e., the Wolf-Israel type)" Certainly to call the organism most frequently associated with lumpy jaw in cattle anything other than *A. bovis* would prove to be confusing.

The difficulty is even more apparent and its solution more important if the genus is to be split into two. Wright uses *Actinomyces* for the anaerobic sporeless forms, and for the aerobic conidia formers the name *Nocardia*, following the emendation of the genus by Blanchard (1896, p. 856). Pinoy, on the other hand, uses the designation *Cohnistreptothrix* for the forms termed *Actinomyces* by Wright, using the name *Nocardia* in Wright's sense.

It would appear, therefore, that the demands of priority would be satisfied by Wright's use of the terms. If all are to be included in a single genus, it should be *Actinomyces*, if two genera they should be *Actinomyces* and *Nocardia*. *Cohnistreptothrix* is to be regarded as a synonym of *Actinomyces* in the narrow sense. The generic names *Streptothrix* and *Oospora* should not be used for this group.

Actinomycetaceae. A family name suggested by Buchanan (1918, p. 403) as the single family of the order *Actinomycetales*. The Committee of the Society of American Bacteriologists (1920, p. 198) have suggested its use as the name of one of the two families of the *Actinomycetales*, the other being *Mycobacteriaceae*. The Committee description is "Filamentous forms often branched and sometimes formed mycelia. Conidia sometimes present. Some species parasitic." This description is also used by Bergey *et al.* (1923, p. 338).

In the present volume the recommendation of the Committee is followed.

Actinomycetales. A designation suggested by Buchanan (1917, p. 162) for the order of the mold bacteria. The original description follows.

Mold-like organisms, not typically water forms, saprophytic or parasitic. Sheath not impregnated with iron, true hyphae with branching often evident,

conidia may be developed, but never endospores Without granules of free sulphur and without bacteriopurpurin Never producing a pseudoplasmodium Always non-motile

The description was emended by the Committee of the Society of American Bacteriologists to read as follows

Cells usually elongated, frequently filamentous and with a decided tendency to the development of branches, in some genera giving rise to the formation of a definite branched mycelium Cells frequently show swellings, clubbed or irregular shapes No pseudoplasmodium No deposits of free sulphur or iron No bacteriopurpurin Endospores not produced, but conidia developed in some genera Usually Gram-positive Non-motile Some species are parasitic in animals or plants Not water forms Complex proteins frequently required As a rule strongly aerobic, (except for some species of *Actinomyces* and the genera *Fusiformis* and *Leptotricha*) and oxidative Growth on culture media often slow, some genera show mold-like colonies

This description is also used by Bergey *et al* (1923, p 337).

The committee recognized two families as belonging to this order, *Actinomycetaceae* and *Mycobacteriaceae* The committee's suggestion is followed in the present volume

Actinomycetes. This family designation of the thread fungi was probably first proposed by Balbiani (1886, p 542) Later Lachner-Sandoval (1898) included in it the single genus *Actinomyces* Harz Lehmann and Neumann (1901, p 127) include within the family three genera, *Corynebacterium* L and N, *Mycobacterium* L and N, and *Actinomyces* Harz They give the following description of the family

Delicate thread organisms, without chlorophyll, with true branching, in some cases with an abundant branched mycelium and the formation of conidia Young cultures often show only unbranched rods resembling bacteria which can in no way be differentiated from the ordinary fission fungi According to many authors there is a tendency to the formation of clubs or knobs at the ends of the threads

The three genera may be differentiated by the following key

- A Cultures showing ordinary bacterial characters, flat or medium, microscopically rods with swollen ends *Corynebacterium* L & M
- B Cultures on solid media more or less folded
 - I Usually short slender rods only, seldom short branched filaments, without aerial mycelium and without aerial conidia Acid fast—*Mycobacterium* L & N
 - II Mycelial filaments long, often bent, without sheath, with true branching Many species produce conidia or aerial hyphae Not acid fast—*Actinomyces* Harz

Orla-Jensen (1909) gives *Actinomycetes* as the name of the second family of the bacterial order *Cephalotrichinae*. He includes in it four genera, *Rhizomonas* Jensen, *Corynemonas* Jensen, *Mycomonas* Jensen, and *Actinomyces* Harz. The designation has been extensively used in the literature as a group name.

It would seem that the form *Actinomycetaceae* would better fit nomenclatural usage in accordance with Article 21 of the botanical code, and in this form would be a valid family name.

Aerobacter. A generic name proposed by Beijerinck (1900, p. 198) for certain gas producing bacteria of the so-called "colon" group. In sugar solutions all are facultative anaerobic, all ferment dextrose and levulose with the formation of levo-rotatory lactic acid and usually with production of gas as well. The gas is a mixture of carbon dioxide and hydrogen. Sulfates are never reduced, while nitrates are readily reduced with formation of nitrites, but never of ammonia. No spores are formed. The organisms are easily killed by heating to 65°C. The cells are frequently motile, either peritrichous or monotrichous. The temperature optimum is about 28°C. The following are the best known species which he lists:

- Aërobacter aërogenes* (*Bacillus lactis aërogenes* Escherich)
- A. coli* (with many varieties, best known, *commune*)
- A. viscosum*
- A. liquefaciens* (Digests gelatine, monotrichous)

Buchanan (1918, p. 53) used this as a subgeneric designation for the colon group of the genus *Bacterium*, with the following diagnosis: "Fermenting both glucose and lactose with formation of both acid and gas. Pathogenicity slight." The type species was designated as *Bacterium* (*Aerobacter*) *coli*. This is probably to be regarded as an error. A reading of the characterization of the genus by its author emphasizes that *Bacterium* (*Aerobacter*) *aerogenes* is, instead, the type for in the list and description of species the latter organism is first given.

Beijerinck (p. 197) emphasizes that this group is to be regarded as a true genus and not merely a physiological genus such as his *Photobacter*.

Aerobacter is rejected by Smith (1905, p. 174).

An *Aerobacter tartarivorum* was named by Nijdam (1907).

Enlows (1920, p. 13) gives the following characterization of the genus:

Facultative anaerobic organisms which give the "white lead test" with production of sulfides, and certain related ferment organisms. No spores. Very resistant to drying. Ferment dextrose and levulose with production of gas and usually lactic acid. Sulfates not reduced. Nitrates reduced to nitrites, but not to NH_3 .

Species *Bacillus coli communis* Escherich. Includes here also *B. liquefaciens* Tataroff, and *Bact. lactis aerogenes* Escherich, the latter becoming *Aerobacter aerogenes*.

Bergey (1923, p. 205) recognizes the genus with the following description:

Motile or non-motile rods, commonly occurring in the intestinal canal of normal animals. Produce acetyl-methyl-carbinol.

The type species is *Aerobacter aerogenes* (Escherich) Castellani and Chalmers.

If the colon typhoid group is to be subdivided, it would seem that *Aerobacter* should be regarded as a valid generic name with *Aerobacter aerogenes* (Escherich) Beijerinck as the type.

Aerogenesbacterium. A generic name suggested by Orla-Jensen (1921, p. 272). His statement follows:

The genus *Bacterium* will undoubtedly dissolve into several genera, of which I may especially mention the *Colibacterium*, and *Aerogenesbacterium*. The reason why I am now inclined, in contrast to my earlier opinion, to consider the coli- and aerogenesbacteria as two different genera, is because they differ not only in morphological, but, as later researches have shown, also in biological respects. I have myself proved (1914), that the *Aerogenesbacteria* completely oxidize the carbohydrates when the nutrient matter offers a sufficient buffer effect, and they thus correspond with their name in forming more gas than do other bacteria, and Rogers, Clark and Davis (1914) have shown that in the gas developed by the *Colibacteria* there is proportionately more hydrogen than in that developed by the *Aerogenesbacteria*.

This generic designation is apparently a synonym of *Aerobacter* Beijerinck. It is also questionable whether the indirect reference is sufficient to establish a species. It is probable that *Aerogenesbacterium aerogenes* would be considered the type species, although this combination was not used by Orla-Jensen.

Aethyl-Bacillus. A name used by Fitz (1878, p. 48) as a pseudo-generic or casual designation for a rod shaped organism capable of producing ethyl alcohol from glycerin. It is rejected by Smith (1905, p. 174) and is invalid as a generic name.

Agonium. A generic name used by Oersted (1844, p. 44) to designate a marine organism, probably an alga (See De Toni. Sylloge

Algarum 5, 684, 1907) It is placed questionably among the bacteria by De Toni and Trevisan (1889, p 938) with the following description

Filamenta cylindrica, articulata, simplicia, basi ab apice superiore distincta, e puncto centrali commune radiatum exorientia, caespites formantia Sporae (endosporae) maximae, ovaes, singulae in unoquoque articulo obvientes

The only species (type) is *Agonnum centrale* Oersted

It is doubtful whether this genus should be included with the bacteria.

Actinomyces. A German variant in the spelling of the generic name *Actinomyces*, *qv* Apparently the first record of its use is by Johne (1881, p 143) It is probably invalid.

Albococcus. A genus of cocci proposed by Winslow and Rogers (1906, p 541) with *Micrococcus pyogenes* (Ros) Mig. as the type The characteristics of the genus are given as follows

Parasites Cells in groups and short chains (never in packets) Generally stain by Gram Growth on agar streak abundant and porcelain white in color Sugars fermented with production of slight amount of acid Gelatin liquefaction and nitrate reduction may or may not occur

Four species are noted *Albococcus pyogenes* (Ros) W and R, *A rhenanus* (Mig) W and R, *A candidans* (Flügge) W and R and *A canescens* (Mig) W and R

Later the Winslows (1908, p 192) revised their list of species giving *Albococcus pyogenes* (Rosenbach) Winslow, *Alb epidermidis* (Gordon) Winslow, *Alb candidus* (Cohn) Winslow, and *Alb tetragenus* (Gaffky) Winslow

Kligler (1913, p 432) has added one species, *Alb ureae* (Cohn, Flügge) Kligler

The genus *Albococcus* is one of the genera included by the Winslows in the subfamily *Paracoccaceae* (*qv*)

To organisms of this group the generic name *Micrococcus* (*qv*) was given by Cohn The forms occurring in irregular masses or grape like clusters, particularly those occurring in pus, were named *Staphylococcus* (*qv*) by Rosenbach (1884, p 19) The species described were the golden and the white pus cocci, or *Staphylococcus pyogenes aureus* and *Staphylococcus pyogenes albus* On a later page, in a main heading, the phrase "*Staphylococcus aureus* (mit oder ohne *albus*)" is used The trinomial forms of the names first given would probably bar the use of the generic designation *Staphylococcus* unless a species

Staphylococcus pyogenes with two varieties *aureus* and *albus* could be conceived. However, the use by Rosenbach himself of the binomial forms *Staphylococcus aureus* and *Staphylococcus albus* answers all nomenclatural requirements. Migula (1894, p. 237) concluded this genus should be merged with the older genus *Micrococcus* Cohn, the species becoming *M. aureus* (Ros.) Migula and *M. pyogenes* (Ros.) Mig.

The Winslows concluded that these species should constitute the types of the genera *Aurococcus* and *Albococcus* respectively. They discarded the name *Staphylococcus* (*qv*) apparently without sufficient justification. The rules of botanical nomenclature state explicitly that when a genus is split into two genera the original generic name must be retained for one genus. (See Art. 45, International Rules for Botanical Nomenclature.) Inasmuch as *Staphylococcus aureus* was the first to be described, the generic name *Staphylococcus* might appropriately be retained for the genus of which this is the type. Accordingly the generic designation *Albococcus* for the white form would appear consistent. It should be noted that the original specific designation of *albus* should be retained, the type of the genus then being *Albococcus albus* (Rosenbach) and not *Albococcus pyogenes* (Rosenbach) Winslow.

In the Preliminary Report of the Committee of the Society of American Bacteriologists (Winslow *et al.*, 1917, p. 559), the genus *Albococcus* is recognized with the following diagnosis:

Differs from *Staphylococcus* in forming more abundant surface growth of porcelain white color, and in fact that liquefaction of gelatin when present is less vigorous.

Buchanan (1917, p. 612) included *Albococcus* as a synonym of *Staphylococcus*.

In the Final Report (Winslow *et al.*, 1920) the genus *Albococcus* is merged with *Staphylococcus*.

Winslow characterizes the genus as follows:

Parasites. Cells in groups and short chains (never in packets). Generally stain by Gram. Growth on agar streak abundant and porcelain white in color. Sugars fermented with production of a slight amount of acid. Gelatin liquefaction and nitrate reduction may or may not occur.

Type species (original designation) *A. pyogenes* (*M. pyogenes* (Rosenbach) Migula). Also include here *A. rhenanus* (Migula), *A. candicans* (Flügge), and *A. canescens* (Migula).

This genus has been considered invalid by Winslow, Rothberg, and Parsons (J Bact , v 5, no 2, 1920, p 161) The type species *A. pyogenes* together with *Staphylococcus epidermidis albus* (Welch) Gordon, and *Albococcus epidermidis* the Winslows, becomes *Staphylococcus epidermidis*

If the white pus cocci and related forms are to be grouped together as a genus, the generic name *Albococcus* Winslow with the type species *Albococcus albus* (Ros) would appear to be valid. If they are not sufficiently distinct to warrant generic recognition, they should be grouped with the golden forms in the genus *Staphylococcus* Rosenbach

Alcaligenes. A generic name proposed by Castellani and Chalmers The first reference is 1919, p. 936. It is here cited, however, as 1918 The diagnosis given is as follows:

Definition Ebertheae which do not ferment glucose or lactose, and are characterized by their general lack of fermentative power and by actually increasing the alkalinity of the media Milk is not clotted, and is rendered alkaline.

Type. *Alcaligenes faecalis* (Petruschky, 1896), *emendavit* Castellani and Chalmers, 1918.

Here also comes *Alcaligenes vivax* (Archibald, 1918), which was obtained from the blood of a case of enteroides in the Anglo-Egyptian Sudan, produced acidity in galactose and mannitol and was characterized by its marked motility.

The genus was also later discussed (1920, p 604) with two species described

Bergy *et al* (1923, p 233) included this as the sixth genus of the tribe *Bactereae* with the description "Motile or non-motile rods, generally occurring in the intestinal canal of normal animals Do not form acetyl-methyl carbinol Do not ferment any of the carbohydrates "

Alkalibacteriaceae. A family of bacteria named by Orla-Jensen (1909, p 313) belonging to the order *Peritrichinae* Orla-Jensen It contains three genera, *Liquidobacterium* Orla-Jensen, *Bacillus* Cohn, *Urobacillus* Miguel It includes those bacterial genera characterized by the production of a decided alkaline reaction in media as a result largely of the development of ammonia The bacteria are neither obligate anaerobes or microaerophiles The name of this family does not conform to Article 21 of the botanical code because it is not designated "by the name of one of its genera or ancient generic names with the ending *aceae* " It would appear therefore not to be tenable

Alcaligenes. A variant spelling of *Alcaligenes* (*qv*) It was used by Castellani and Chalmers (1919, pp. 1405, 1408, 1409) in referring to their genus *Alcaligenes* described earlier in the same volume. It is also used by Evans (1923, p 1945)

Allococcaceae. A subfamily of bacteria belonging to the family *Coccaceae* described by Fischer (1897) (1903, p 59) The subfamily is regarded as made up of two genera, *Micrococcus* Cohn and *Planococcus* Migula. It contains those cocci in which the planes of fission have no definite sequence, and there are no pronounced colonies or growth forms The cells are isolated, in short chains or irregular clusters

The ending of the name of this subfamily, if used, should be *ordeae* rather than *aceae* in conformity with Article 23 of the International Code The code specifically states "Names of subfamilies (*subfamiliae*) are taken from the name of one of the genera in the group with the ending *ordeae*" As a subfamily name *Allococcaceae* would therefore appear not to be tenable for this reason and because not derived from the name of one of the constituent genera

Amoebobacter. A genus of the sulphur bacteria described by Winogradsky (1888, p 71) He says,

findet man eine Familie von so dicht zusammengepressten Zellen, dass man die einzelnen Zellen fast gar nicht unterscheidet, und die ganze Gruppe fast homogen erscheint Die Färbung ist zart rosenroth hier und da finden sich wenige Schwefelkörnchen Um die Zellenmassen sieht man eine Hülle, welche aus zwei Schichten besteht, eine innere schwach lichtbrechende und eine äussere stark lichtbrechende deutlich doppelt contourirte Schicht Stellt man eine solch encystierte Familie ein, so sieht man etwa nach 4 Tagen dass die Cyste gesprengt wird und ihr nunmehr deutlich aus Coccen bestehender Inhalt ausserst langsam herauszukriechen beginnt Der Austritt dauert manchmal 1-2 Tage Es bleibt eine leere Cyst zurück, welche sich noch lange unverändert erhält Der Gestaltwechsel der Amoebobacterfamilien hat seinen Grund darin, dass sowohl die einzelnen Zellen innerhalb der Familie, als die Familien als ganze beweglich sind

Three species are described, *Amoebobacter roseus*, *A granula* and *A bacillosus* In the first the cells are spherical 2.8 to 3.4 μ in diameter, or elongate to 6 μ In the second the cells are very minute, scarcely 0.5 μ , and in the last the cells are rod shaped, 2 to 4 μ long and 1.7 μ thick

The genus was next recognized by De Toni and Trevisan (1889, p 1043) as the only genus of the tribe *Amoebobacterieae* Trevisan They give the following definition

Cocci globos vel ovoideo-ellipsoidei, densissime in familias primitus adparenter fere homogenes, amoeboidemobiles cumulati cysti debus bistratosi circumcincti interiori stratu parum refringente, exteriori valde refrigente cito delabentibus Cocci demum liberi mobiles Coccorum divisio in unam directionem

Winogradsky's three species are recognized

An adequate study of this group since the monograph of Winogradsky does not seem to have been made, although the genus *Amoebobacter* Winogradsky has been accepted by many subsequent authors. This genus was included by Migula (1900, p. 1045) in the subfamily *Amoebobacteriaceae*, together with the genera *Thiodictyon*, *Thiothece* and *Thiopolycoccus*. Migula (1904, p. 146) gave the following generic diagnosis: "Zellen zu Familien vereinigt, nach einer Richtung des Raumes sich teilend, Familien amoeboid beweglich, Zellen durch Plasmafaden verbunden."

Erwin F. Smith (1905, p. 163) likewise places the genus in the family *Amoebobacteriaceae*, and with the following description.

Cells connected by plasma threads. Families amoeboid motile. The cell families slowly change form, the cells drawing together into a heap or spreading out widely, thus bringing about a change in the shape of the whole family. In a resting condition a common gelatin is extruded, the surface of which becomes a firm membrane.

In several descriptions, the connecting threads are designated as protoplasmic, apparently without sufficient justification. For example, Frost (1911, p. 61) in a tabulation of Migula's classification states:

The cells are united into colonies and after division in one direction of space remain attached together by threads of protoplasm. The colonies possess amoeboid motility. The cells change form by contraction and the spreading out of the protoplasm.

Buchanan (1918, p. 409) designated the first species described by Winogradsky, *Amoebobacter roseus*, as the type. His description follows:

Cells connected by plasma threads. Families amoeboid motile. The cell families slowly change form, the cells drawing together into a heap or spreading out widely, thus bringing about a change in the shape of the whole family. In a resting condition a common gelatin is extruded, the surface becomes a firm membrane.

Enlows (1920, p. 14) gives the following characterization:

Cells divide in one direction of space, usually round, and united into families by means of plasma threads. Families have amoeboid motion. In the resting state the extruded gelatin becomes stiffened, forming a firm 2-layered membrane. Sulfur granules here and there. Cell masses a delicate rose red.

Type species (subsequent designation by Buchanan (J. Bact., v. 3, no. 5, 1918, p. 469)) *Amoebobacter roseus* Winogradsky. Cells spherical, 2.8 to 3.4 μ in diameter. Winogradsky also includes here *A. bacillosus* (rodlike cells 2 to

4 μ long by 1.7 μ), and *A. granula* (cells spherical, exceedingly small—scarcely 0.5 μ in diameter)

Bergey *et al* (1923, p 398) followed Buchanan

Orla-Jensen (1909, p 334) proposed that *Amoebobacter* be replaced by a new term *Amoebomonas*

Apparently no author has studied the organisms of this genus since Winogradsky's original description

Amoebobacteriaceae. A variant spelling of *Amoebobacteriaceae* used by Bergey *et al* (1923, p 398) for the fourth tribe of the subfamily *Chromatiodeae* with the description "Sulphur bacteria in which the cells are united into families Cell division occurring only in one direction of space"

One genus only, *Amoebobacter* is included

Amoebobacteriaceae. A subfamily of the family *Rhodobacteriaceae* of the sulphur bacteria established by Migula (1900, p 1045) to contain the genera *Amoebobacter*, *Thiodictyon*, *Thiothece*, and *Thioplycoccus*

This subfamily is characterized by having the cells united into families, cell division occurring in one direction of space It has generally been recognized by subsequent authors dealing with this group, as Migula (1904, p 146), Erwin F Smith (1905, p 163), Frost (1911, p 61)

Article 23 of the International Rules for Botanical Nomenclature would require the change of the subfamily name to *Amoebobacterioidae*

Amoebobacteriaceae. A subtribe of the tribe *Ascococceae* Trevisan (1889, p 1043) established by Trevisan, to include the single genus *Amoebobacter* The diagnosis given is "*Cocci in muco matricali in series filamentosas laze conjuncti Cystides universales tenuissimae, cito dilabentes Cystides speciales nullae*"

The International Rules of Botanical Nomenclature (Article 23) would require the change of this name if used for designation of a subtribe to *Amoebobacterinae*

The same name was used by Buchanan (1918, p 469) to designate a tribe with the following diagnosis "Sulphur bacteria in which the cells are united into families Cell division occurring only in one direction of space" Four genera were included *Amoebobacter*, *Thiodictyon*, *Thiothece* and *Thioplycoccus* Bergey *et al* (1923, p 398) used the spelling *Amoebobacteriaceae* *g v*

Amoebomonas A name proposed by Orla-Jensen (1909, p 334) to replace the generic name *Amoebobacter*, Winogradsky of the *Rhodo-*

bacteriaceae It is a synonym, and probably invalid as a generic designation. No species were designated.

Amphitricha. A designation suggested by Messea according to Kruse for those motile bacteria with a flagellum at each end of the cell. It is the first group of the *Trichobacteria* q v.

Amylobacter. A casual name used by Trecul (1865, p. 435) for certain rods found between the cells of decaying plant tissues. He said

Malgré la variété des formes de ces petits végétaux, au plutôt à cause de cette variété même puisqu'il on passe d'une forme à l'autre par toutes les gradations, en raison aussi de l'amidon qu'ils contiennent, et pour rappeler la ressemblance des formes cylindriques avec les Bactéries, je crois utile de les réunir sous le nom d'*Amylobacter*. Si on voulait les diviser d'après les formes les plus disparates, on pourrait établir trois genres, 1, '*Urocephalum* qui comprendrait les formes en têtard, 2, l'*Amylobacter* vrai, auquel seraient attribuées les formes cylindriques, 3, le *Clostridium*, qui refermerait les formes en fuseau.'

This author gave no specific names. What was later regarded as the same organism was named *Bacillus amylobacter* by Van Tieghem (1879, p. 25).

Duclaux (1895, p. 813) revived the name, describing two species, *Amylobacter butylicus*, and *Amylobacter ethylicus*.

Ce sont partout des bacilles dont la largeur et la longueur varient avec le milieu de culture, cylindriques lorsqu'ils sont jeunes, se renflant plus ou moins quand ils vieillissent, en un point où apparaît la spore. Chez la plupart d'entre eux, de même du reste qu'avec des bacilles qui ne consomment ni amidon ni cellulose, la formation de la spore est précédée d'une période où une partie du protoplasme du bacille se colore par l'iode.

He notes that Beijerinck had previously suggested *Granulobacter* for this type of organism.

The genus should be ascribed to Duclaux. *A. butylicus* is the type. *Amylobacter* is listed as a synonym of *Clostridium* by Prazmowski (1880, p. 23) and by De Toni and Trevisan (1889, p. 1002). It is rejected by Erwin F. Smith (1905, p. 174).

The use by Trecul of the name *Amylobacter* does not entitle it to generic recognition because it was not coupled with the name of any particular species in the form of a binomial. It existed therefore, simply as a casual name until its use by Duclaux. In the meantime, however, the genus *Clostridium* (q v) had been created for such types of bacteria. As a generic designation it should probably be written

Amylobacter Duclaux It is a synonym of *Clostridium*, and *Granulobacter* at least in part

Amylobakter. A German spelling of *Amylobacter* A casual name It has been used among others by Schattenfroh and Grassberger (1899, p 700) It is not a valid spelling of the generic name

Aphaneroglia. A generic name proposed for the single genus of the first order *Protobacteries* by Maggi (1886, p 81) One species, *A. apuiae potabilis* is listed, but without description

Aplanobacter. A generic name proposed for non-motile rods by Erwin F Smith (1905, p 171) The diagnosis given by Smith is as follows

An unattached, non-motile, rod-shaped organism, destitute of chlorophyll and multiplying by fission, sometimes forming threads of considerable length The type of the genus, in the family *Bacteraceae*, is that organism causing anthrax and most commonly known in literature as *Bacillus anthracis* Cohn

The generic names *Bacillus* Cohn and *Bacteridium* Davaine have both been applied to this type species The type of the genus *Bacillus* Cohn is generally accepted as *Bacillus subtilis* Cohn The essential morphological difference between *B subtilis* and *B anthracis* is the possession of peritrichous flagella by the former, and their absence in the latter Smith contends that lack of motility is a characteristic which warrants the separation of the anthrax organism from *Bacillus* Evidence is continually accumulating to show that generic separations among the bacteria solely on the basis of motility lead to a violent disarrangement of natural groupings Granting, however, that another generic name should be found, Davaine's genus *Bacteridium* would appear to be tenable It was founded upon the *B anthracis* and created because of the lack of motility of this form Smith argues that the name *Bacteridium* (q v) is preoccupied in botany However, the names previously used (see discussion of *Bacteridium*) have been spelled *Bactridium* Smith regards this as the same word, making *Bacteridium* untenable As noted elsewhere, a strict interpretation of the rules of the International Code would indicate that the difference in spelling is sufficient to make the generic name *Bacteridium* valid It should also be noted that the generic name *Pollendera* Trevisan (1885) has priority over *Aplanobacter*

Migula has defined the genus *Bacterium* (q v) to include the non-motile rods As noted elsewhere, this use of the term *Bacterium* is scarcely justified, it probably should not be used in this group.

It would seem that the generic name *Aplanobacter* Smith must be regarded as a synonym of *Bacteridium* Davaine, and hence invalid unless it can be shown that for some reason the latter name as well as *Pollendera* are not tenable.

Arloingillus. A genus of anaerobic spore bearing bacteria proposed by Heller (1922, p 21) Her description follows:

Clostridiaceae that attack sugars with considerable energy but have a somewhat restricted action on proteins Liquefy gelatin but do not produce H_2S demonstrable by a lead-acetate-paper test in blood broth In meat medium produce gas and a pink coloration that soon fades Autoagglutinate readily Clot milk if blood is present Do not digest serum or egg Vegetative forms are small Gram-negative rods with even staining forms about to sporulate are uneven in staining reaction, often far larger than vegetative rods citron or spindle shaped, orgonts (see Heller), show marked tendency to store up granulose Spores oval, vary greatly in length Bacilli do not form long chains in the liver of animals Colonies in deep agar lenticular, sometimes showing concentric formation, or compound lenticular Colonies vary considerably according to species Typically toxic tissue invaders which produce marked haemolysis Pathogenic for guinea-pigs cattle and sheep

The species *A. Chauv.* *Bacterium Chauv.* Arloing, Corneval and Thomas as described in a future paper

This genus contains several species which will be discussed These organisms show some similarity to those of the genus *Clostridium*

The *B. enteritidis-sporogenes* Klein as described by von Hübner 1903 should probably be included in the genus The only character which is markedly different from that of the genus is the energetic fermentation of milk shown by von Hübner's bacillus IV

Arthrobacillus A genus of bacteria proposed by Fischer (1897 p 136) to include the non-motile rod-shaped organisms which produce arthrospores No type was designated

Arthrobacter. A generic name proposed by Fischer (1895 p 141) to include all non-flagellate rod-shaped bacteria which produce arthrospores No species are named, hence the genus is invalid He says "Vorläufig kann keine Art mit Sicherheit hierhergestellt werden vielleicht wurde das *Bacterium Ureae* hierhergehören " He apparently ascribes the name to De Bary as he writes *Arthrobacter* De Bary Fischer (1903) himself later abandoned the designation It is listed as invalid by Erwin F Smith (1905, p 174)

Arthrobacteriaceae. A "Gattung" proposed by Hueppe (1886, p 145) to include the rod shaped bacteria which produce arthrospores or at least do not form endospores The "Untergattungen" *Arthrobacterium* (*Bacterium* s str) and *Spirulina* (*Proteus*) are included Evidently this should be regarded as a casual family name

Arthrobacterium. A name proposed by De Bary (1884) to include rod-shaped bacteria which produce arthrospores. No species were described. The genus was also mentioned by Hueppe (1886) and was included in his classification of bacteria. De Toni and Trevisan (1889, p. 1054) give *Arthrobacterium* De Bary (1887) as a synonym of *Streptococcus*. As stated by Vuillemin (1913, p. 520) this genus has not been preserved or used because it was founded upon the false assumption of arthrospore formation.

Enlows (1920, p. 15) states

According to Comp. Morph. and Biol. Fungi, Mycetozoa and Bact., Auth. Eng. rev. trans. by Garnsey, Oxford, 1887, pp. 454 and 468. This genus is proposed for that group of the genus *Bacterium* in which no endogenous spore-formation occurs. "To denote the species which constitute the genus *Bacterium* of authors, I use partly the generic name *Bacillus* . . . and partly the name *Arthrobacterium*. Single members may simply separate from their connections with others, and under suitable conditions become the initial members of new combinations, they have therefore claim to the name spore. In other respects there is no general characteristic distinction between them and the purely vegetative members.

Species: *Bacterium zopfii* Kurth, *Bact. merismopoeidioides*, *Bact. aceti*, and *Bact. pastorianum* Hansen

Maggi (1886, p. 84) gives *Arthrobacterium aceti* De Bary as a synonym of *Bacterium aceti* Zopf.

It is doubtful whether the rather casual use by De Bary is sufficient to validate the genus. Apparently no type has been designated.

Arthrobactridium. A genus proposed by Fischer (1895, p. 140) to include those rod-shaped bacteria with diffuse flagella which produce arthrospores. The genus was later abandoned by Fischer (1903) as based upon a false interpretation of morphology. It is also rejected by Erwin F. Smith (1905, p. 174). The genus is not tenable because no species have been definitely assigned to it.

Arthrobactrillum. A generic name proposed by Fischer (1895, p. 139) to include those rod-shaped bacteria which are motile by means of a tuft of polar flagella, and which reproduce by means of arthrospores. The name is invalid. It rests upon a mistaken interpretation of morphology and has no described species. It was later abandoned by Fischer (1903) himself as untenable. It is also rejected by Erwin F. Smith (1905, p. 174).

Arthrobactrinium. A generic name proposed by Fischer (1895, p. 139) to include rod-shaped bacteria having one polar flagellum and producing arthrospores. This generic name has never come into

general use It is founded on a misconception of bacterial morphology, has no named species, and has been abandoned by its author (1903). It should be considered as invalid It is rejected by Erwin F Smith (1905, p 174).

Arthrobakteriaceen. A variant of *Arthrobacteriaceen* A name used by Hueppe, (1886, p 145) for a "Gattung" including the rod-shaped bacteria which produce arthrospores or at least which do not produce endospores The "Untergattungen" included are *Arthrobakterium* (*Bakterium* s str.) and *Spirulina* (*Proteus*). The name is not Latin in form It must be regarded as untenable

Arthrobakterium. A spelling of *Arthrobacterium* De Bary (q v) used by Hueppe (1886, p 145) to designate an "Untergattung" of the "Gattung" "*Arthrobacteriaceen* " It is also given as *Bakterium sensu strictu*. It includes those straight rod shaped organisms which produce arthrospores, or at least do not produce endospores

Arthrokokkaceen. A name used by Hueppe (1886, p 144) for a "Gattung" of bacteria with spherical cells which produce arthrospores or whose method of spore production is unknown The following "Untergattungen" are included *Arthro-Streptokokkus*, *Leuconostoc*, *Merista*, *Sarcina*, *Mikrokokkus*, *Askokokkus* It may be regarded as invalid

Arthromitus A generic name proposed by Leidy (1849, p 227) for two organisms found in the rectum of *Julus* These he named *Arthromitus cristatus* and *A nitidus*

The following is the description given by Leidy to the genus and to the first species described

Filaments always simple, cylindric articulated, without ramuli, attached by means of a nuclear body, and with no evidence of interior structure

Arthromitus cristatus Filaments delicate straight or inflected, growing in tufts usually of moderate density, from minute, attached yellowish rounded or oval nuclear bodies Articuli short, cylindric, uniform measuring $\frac{1}{100}$ inch in length by $\frac{1}{1500}$ in breadth, with no traces of interior structure

Length $\frac{1}{35}$ to $\frac{1}{40}$ inch breadth $\frac{1}{1500}$ inch

Habitat Same as *Cladophytum comatum*, but rarely growing in such dense tufts

Later (1852, p 35) he described *Arthromitus nitidus* This species is described as follows

Filaments very long, hyaline, grows usually in twos or fours, pointed at the origin, rounded at the termination Articuli very distinct length equal to the breadth of the filament Sporuli formed within the articuli solitary, usually oblique, oval, amorphous

Length 1 line by $\frac{1}{5000}$ inch broad Spores $\frac{1}{111}$ inch long, by $\frac{1}{12,500}$ inch broad.

Habitat Grows in considerable quantity with a profusion of young of *Entero-bryus elegans* from the mucous membrane of the posterior portion of the rectum of *Julus marginatus*

Remarks Since I established the genus *Arthromitus* I have observed the formation of its sporuli. These originate in the amorphous matter of the articuli, apparently by a very gradual aggregation and condensation of the contents. They are always single, and usually lie oblique, and frequently alternate with each other in this position in the different articuli. When they first appear they are larger than when fully formed, are frequently bent, or clavate in form, and very indistinct, but as they ripen they become more regular, oval, distinct, and quite refractile of light. Usually they are observed at the extremity of the filaments only, but frequently they are found existing in the whole length of the latter.

A species of *Arthromitus*, and also of *Cladophytum* is found in the intestine of *Polydesmus virginensis*

Robin (1853, p 359) gives the following generic and specific description

Filaments toujours simples, cylindriques, articulés, sans ramifications, fixés au moyen d'un corps nucléaire, sans structure bien définie

Arthromitus cristatus, Leidy Filaments fins, croissant en touffes médiocrement épaisses, sur de petits corps arrondis, fixés et jaunâtres, articles courts, cylindriques, uniformes, ayant 0^{mm},002 ($\frac{1}{5000}$ de pouce) de long et 0^{mm},001 ($\frac{1}{12,500}$ de pouce) de large, sans structure intérieure

Ce végétal offre une longueur qui varie entre 0^{mm},065 et 0^{mm},543 ($\frac{1}{375}$ et $\frac{1}{45}$ de pouce), et une largeur de 0^{mm},001 ($\frac{1}{5000}$ de pouce). Il se trouve dans les memes points que le *Cladophytum comatum*, mais rarement en touffes aussi épaisses

Arthromitus nitidus, Leidy Filaments très longs, hyalins croissant au nombre de deux ou quatre, amincis à l'origine, arrondis vers leur terminaison articles très distincts, à longueur égale à la largeur du filament, et renfermant des sporules généralement ovales, amorphes et isolées

Ce végétal a environ 2^{mm},13 (1 ligne anglaise de long), sur 0^{mm},005 ($\frac{1}{5000}$ de pouce) de largeur. Spores, 0^{mm},003 ($\frac{1}{111}$ de pouce) de long, ou 0^{mm},002 ($\frac{1}{12,500}$ de pouce) de largeur

Il croit en grande quantité avec l'*Entero-bryus elegans* dans la muqueuse de l'extrémité du rectum du *Julus marginatus*. Une espèce d'*Arthromitus* se trouve aussi dans l'intestin du *Polydesmus virginensis*

Les sporules des *Arthromitus* se forment par condensation de la substance amorphe qui remplit les articles. Ils sont toujours isolés, ordinairement placés obliquement. Quand ils commencent à se former, ils sont plus grands que lorsqu'ils ont acquis tout leur développement, un peu recourbés et très peu distincts, mais en murissant ils deviennent plus réguliers, ovales, très nets, et réfractant bien la lumière. Quelquefois on ne les trouve qu'à l'extrémité des filaments, mais souvent il en existe dans toute leur longueur

Collin (1913, p. 63) revived the generic name as a substitution for an organism from the alimentary tract of the frog *Arthromitus batrachorum* is described as follows

Son thalle filamenteux, non rameux, à croissance indéfinie, à contenu granuleux partagé en articles d'environ 3,5 μ de large sur 2 à 3 μ de hauteur, circulaires en coupe transverse, rappelle d'assez près le port de certaines Oscillariées mais il est incolore. De place en place, et sur certains thalles seulement, des cellules plus épaissies, à contenu très colorable font quelque peu saillie sur l'alignement des autres après la fixation, on voit souvent, de chaque côté, des cellules claires dégénérées. Je pense qu'il s'agit la peut-être d'éléments vieillis ou morts, tels que ceux qu'on observe chez beaucoup de Cyanophycées (les *Scytonema* par exemple), éléments qui pourraient se trouver libérés par la rupture du thalle. La sporulation a lieu par voie endogène, avec une spore pour chaque cellule et d'une manière toute sporadique pour les cellules d'un thalle donné, la spore est arrondie, circulaire en coupe transverse, mais très nettement ellipsoïde quand on l'aperçoit de profil. Elle est située à l'intérieur de la cellule avec une forte obliquité. Les ébauches sporales, d'abord sphériques et très petites, puis plus volumineuses, se colorent intensément par les méthodes à froid (hématoxyline ferrique par exemple), tandis qu'à maturité complète, elles ne sont plus pénétrables qu'à chaud, comme chez les autres Bactéries. Rectum des têtards d'Alyte en compagnie de *Bacillus campyospora* et de *Spirillum proclerum*, trouvé aussi (ou une forme très voisine) dans l'intestin postérieur des larves de *Bufo calamita* Laur. à Viarmes (Seine-et-Oise).

Chatton and Perard (1913, p. 1160) conclude that the *Hygrocrocis intestinalis* of Valentin and the *Arthromitus cristatus* of Leidy belong to the same genus and on grounds of priority they select *Hygrocrocis*. They say

Dans des préparations de contenu intestinal de territes nous avons retrouvé les *Arthromitus* de Leidy, et M. A. Borrel nous a montré chez des Blattes des schizophytes que nous identifions à l'*Hygrocrocis intestinalis* de Valentin. Il nous semble d'ailleurs que ces formes sont génériquement identiques.

Ce sont de longs filaments à compartiments subgauls cubiques à croissance indéfinie, immobiles et qui au surplus sont d'après Leidy, fixés à la paroi intestinale par bouquets sur une sorte de pied commun globuleux. Il y a dans les régions sporulées de ces filaments une endospore par cellule et ces cellules sporogènes ne sont point hypertrophiées. (See *Hygrocrocis*.)

Apparently *Arthromitus* is a valid generic designation. *Arthrospirobacterium*. Enlows (1920, p. 16) states

This genus has been ascribed to Hueppe. In his "Die Formen d. Bakt.," Wiesbaden, 1885, p. 146, where he classifies the bacteria, he gives as "Gattung" III, Arthro-Spirobakteriaceen. The vegetative cells are screw-like rods. No endogenous spores. Arthrospores. He places *Spirochaeta* as a subgenus of this.

The ascription noted by Enlows has not been found.

Arthro-Spirobacterium. Enlows (1920, p 88) states

Arthro-Spirobacterium Klebs, 1887

Die Allg Path, Jena, 1887, p 88 In a résumé of Hueppe's classification, Klebs gives this spelling, and cites the 1881 edition of Hueppe's *Die Formen d. Bakt*

Arthrospirobakteriaceen. A name used by Hueppe (1896, p 146) to designate a "Gattung" of bacteria in which the cells are spiral, and which produce arthrospores, or at least do not produce endospores One "Untergattung" only, *Spirochaeta*, is included. The name is evidently invalid, or at most, a casual family designation

Arthrostreptococcus. A variant (in proper form) of *Arthro-Streptokokkus* q v.

Arthro-Streptokokkus. A sub-generic name proposed by Hueppe (1886, p. 144) to include those bacteria with spherical cells, occurring in chains in indefinite zoogloea, and without endospores It is based upon a misconception of morphology, it is not spelled as a Latin word, no species are definitely assigned to it, and it is essentially synonymous with *Streptococcus*

Ascobacillus. Apparently first introduced as a casual name by Edington (1887, p 1265) for an organism found in the blood of a patient with scarlet fever De Toni and Trevisan (1889, p 1034) under "species quod genus dubia" give Edington's "ascobacillus of Scarlet Fever" as a synonym of *Klebsiella Edingtoni* Trevisan The following description is given

Baculi cylindracei, tenuibus, 0 8 μ longis, 0 2 μ latis Sporae pluries centum in capsulis, quam baculi multo amplioribus, lucanicaeformibus contentae Hab in squamis et sanguine hominum scarlatina laborantium In culturis gelatinamo lente liquefacit, coloniae flavae

Enlows (1920, p 16) says

Small rods, 0 8 μ long by 0 2 μ broad, often dumb-bell shaped, made up of long ovoid spheres Spores are contained in large sausage-shaped capsules many hundred times larger than the bacilli themselves Found in the blood of a patient dying of scarlet fever

As a generic designation *Ascobacillus* was first used by Unna and Tommasoli (1889) The name first appeared (p 15) as *Ascobacillus citreus* in a list of organisms isolated from the skin Later (p 60) the organism is more adequately described

GENERAL SYSTEMATIC BACTERIOLOGY

The organism is a straight or somewhat bent rod, $1\frac{3}{4}\mu$ by $0\frac{3}{4}\mu$, single or pairs or bundles, motile, spores unknown. On gelatin the growth is very slow, in two weeks the superficial colonies are opaque yellow points, the deep colonies scarcely yellow. Under low magnification the latter are gray yellow, opaque. The organism is aerobic, does not produce gas and forms a citron yellow pigment.

Migula (1900, p. 842) termed this organism *Bacillus citreus*.

Moreno (1901, p. 112) described an *Ascobacillus aquatilis*. Macé (1913, p. 586) regards *Ascobacillus* as a synonym of *Ascobacterium* q.v. making *Ascobacillus citreus* a synonym of *Ascobacterium luteum* Babes. Macé also includes as a synonym the *Ascobacillus sacchari* Greig-Smith (1903).

Enlows states (1920, p. 16 for the Unna type):

Straight or bent bacilli, 1 to 3μ by $0\frac{3}{4}\mu$, single or in twos, grouped or in bundles. Masses taken from agar present the appearance of *Ascococcus billrothii* Cohn. In the interior of these masses the bacilli are nonmotile, but at the periphery a whirling motion may be observed.

Type species (monotypy) *A. citreus*. Produces a citron yellow color on media. Liquefies gelatin.

If the organism described by Unna and Tommasoli is worthy of recognition as a generic type, the name *Ascobacillus* would probably be valid. It is possible that it might be used as a generic name for yellow pigment forming rods if these should be grouped together, as has been done in *Flavobacterium* q.v. Otherwise *Ascobacillus* may be regarded as a synonym and not tenable.

Ascobacteria. A name (form genus) suggested by Billroth (1874) but with no designation of species.

It was proposed as a generic name by Van Tieghem (1880, p. 151). Under the grouping of colonies provided with a membrane, he states that the types with cylindrical cells may be combined into the genus *Ascobacteria*, which he regarded as an encapsulated *Polybacteria*. One species was described, *Ascobacteria ulvina* Van Tieghem. It developed on the surface of water in which leguminous seeds (lupine) were rotting. It occurred in granular, polyhedral masses, each enveloped in a thick cartilaginous membrane, in a layer like the alga *Ulva*. Cells numerous small rods embedded in a common gelatin. After the mass reaches a certain size, it splits, while the gelatinous membrane is continued over the two daughter masses.

Later Van Tieghem (1884, p 1114) evidently regarded this as a form genus. It was also recognized by Billet (1880, p 24)

Apparently the genus was validly described, with type species *Ascobacteria ulvina* Van Tieghem. However, the species is so poorly characterized that its identification is questionable.

Ascobacterium. A generic name used by Babes (1890, p 155) in his description of *Ascobacterium luteum*. This organism is mentioned as common in the air of the laboratory at Buda Pesth and in the water of the Dumbivotza. It forms very pronounced transparent, yellowish colonies on gelatin. The cells are rod-shaped, united in greater or less numbers by a large gelatinous or mucous capsule.

Macé (1897, p 962) believes that there may be a relationship existing between this form and that described under the name *Ascococcus*.

Macé (1913, p 586) states

Les éléments en bâtonnets, véritables Bacilles, sont réunis, en nombre plus ou moins considérable, dans une grande capsule ovale, gélatineuse ou muqueuse. Le nom d'*Ascobacillus* doit être considéré comme synonyme et serait certainement à préférer, si l'on était tout à fait fixé sur l'individualité de ces types microbiens, on peut conserver le terme *Ascobacterium* comme provisoire. Il y a peut-être des rapports à établir entre ce type et ceux décrits sous le nom d'*Ascococcus*.

He describes *Ascobacterium luteum* Babes and *Ascobacterium aquatile* (Moreno) Macé (*Ascobacillus aquatilis* Moreno).

It would seem that this genus has been validly described, with type species *Ascobacterium luteum* Babes. It is possible that it should be regarded as a synonym of *Ascobacillus*, q v.

Ascococceae. A name of a tribe of cocci of the subfamily *Coccogae* Trevisan proposed by Trevisan (1889, p 1037). It contains the sub-tribe *Euascococceae* Trev. with the genera *Lamprocystis* Schroeter, *Ascococcus* Cohn, *Bollingeria* Trevisan, *Cenomesia* Trevisan, *Throcystis* Winogradsky, *Thiothece* Winogradsky, *Leucocystis* Schroeter, the sub-tribe *Gaffkyaee* Trevisan with the genera *Chlamydatomus* Trevisan and *Gaffkya* Trevisan and the subtribe *Amoebobacterieae* Trevisan, with the single genus *Amoebobacter* Winogradsky.

It is characterized as follows: "Cocci in familias tegumentis vesicaeformibus gelatinosis (cystidibus) obductae consociati."

The name has apparently been used by no other author.

Ascococcus. A form genus proposed by Billroth (1874, p 13). It was intended to include those growth forms of the polymorphic

Coccobacteria septica (q v) in which the cells are spherical and embedded in relatively dense zoogloeal masses, the masses being well limited, usually oval and distinct. To one particular form the binomial *Ascococcus parvus* (p 98) was given. This is insufficiently described and cannot now be recognized.

The generic designation *Ascococcus* Billroth should be regarded as invalid for the following reasons

1. The name *Ascococcus* uses the Greek termination *os* instead of the Latin *us*. Article 7 of the Botanical Code specifically states

Scientific names are in Latin for all groups. When taken from another language, a Latin termination is given them, except in cases sanctioned by custom.

2. The name was never proposed by Billroth as a valid generic designation but as a growth form only.

3. The only species (*A. parvus*) assigned to the genus was inadequately described.

This name was changed to *Ascococcus* by Cohn (1875, p 147) who writes "Billroth schreibt *Ascococcus*, wie *Micrococcus*, etc. der Gebrauch sanctionirt jedoch nur die lateinischen Endungen selbst bei Namen griechischen Stammes (Lois de la nomenclature botanique Article 63)." The generic name therefore, in spite of Cohn's statement '*Ascococcus* Billroth char. emend.' should read *Ascococcus* Cohn' (q v).

Ascococcus. The form genus *Ascococcus* (q v) was created by Billroth (1874, p 13) for a growth form of his *Coccobacteria septica* in which the spherical cells are embedded in gelatin or slime. This is not tenable as a generic designation. Cohn (1875, p 154) published the genus name *Ascococcus* Billroth char. emend. This should be written *Ascococcus* Cohn.

The generic description given by Cohn is

Cellulae achromaticae globosae densissime consociatae in familiis tuberculosas globosas vel ovaes irregulariter lobatas, lobis in lobules minores sectis capsula globosa vel ovali gelatinoso-cartilaginea crassissima circumdatas, in membranam mollem facili secendentem floccosam aggregatas.

One species was described, *Ascococcus Billrothii*, from a zoogloea mass developing spontaneously in a culture medium of ammonium tartrate. The specific diagnosis is

Familiae tuberculosae 20-160 μ capsula ad 15 μ crassae. In solutione ammonii tartarici acidi aëro lavata vel butyrico praeditam formantem observavi Mart.

1874 Haud scio citrum eandem an affinem speciem illi Billroth in aqua carnis foetida detexerit

Cohn believed that the organism which he described was the same as that of Billroth, hence the choice of name. He also concluded that the *Ascococcus parvus* of Billroth was not a true *Ascococcus* in the sense of Billroth's original description.

Cienkowski (1878, p. 12) described an organism responsible for a gummy or viscous fermentation of syrups in sugar factories. He regarded it as conforming to Cohn's conception of *Ascococcus* and named it *A. mesenteroides*. A study of the same organism was made by Van Tieghem (1878, p. 198). He concluded that this form was distinct from *Ascococcus* Cohn and made it the type of the genus *Leuconostoc*. He called attention to the points of difference between the two genera. In *Ascococcus* the cells are spherical, very small, and grouped in great numbers to form globular or avoid families which are lobed more or less irregularly. The cells are closely united, separated only by a small amount of gelatinous material, each family is surrounded by a cartilaginous envelope. In *Leuconostoc* the cells are arranged in bent chains separated from each other by a large amount of gelatinous material, the gelatin on the exterior not being thicker than that between the chains. The *Ascococcus* of Cohn grew in ammonium tartrate solution, the *Leuconostoc* in sugar and rendered the solution decidedly acid. He places the genus *Leuconostoc* among the *Nematogenae* in Cohn's classification, while *Ascococcus* is with the *Glaeogaeae*.

Trevisan (1879, p. 137) reworded Cohn's description as follows: "Cellulae globosae, inordinate in colonias conglobatas pluristratas densissime consociatae, tegumentis propriis nullis. Coloniae tegumento communi gelatinoso-cartilagineo crassissimo obvolutae." He included it in the subtribe *Sarcineae*.

Ascococcus Billroth was recognized by Magnin (1880, p. 96) with the species *A. billrothii* Cohn. Winter (1880, p. 47) likewise recognized the genus as *Ascococcus* Cohn. He questions, however, the value of Cohn's genus and also whether the genera of Cohn and Billroth are based upon the same form. He believes it probable that *Ascococcus* Cohn is only a developmental stage of *Micrococcus* Cohn.

Van Tieghem (1880, p. 150) recognized the genus, and added to it the species *Ascococcus vibrans* from the surface of water in which *Beggiatoa* was putrefying.

Grove (1884, p. 15) used practically the same words as Winter in his discussion of the genus. Van Tieghem regarded *Ascococcus* as a subdivision of *Micrococcus* including those forms with cells embedded in gelatin.

Zopf (1885, p. 51) in the third edition of his *Spaltpilze*, though not in the preceding, recognizes *Ascococcus* Billroth as a *Micrococcus* with intensive gelatin production Hueppe (1885) states that in the genus *Ascococcus* the cells are spherical, not in chains, occurring in irregular masses in spherically segmented zoogloea Schroeter (1886, p. 153) recognizes "*Ascococcus* Billroth 1874 (in der Umgrenzung von Cohn 1875)" According to Flugge (1886, p. 140) *Ascococcus* includes spherical bacteria united into definite slime families, colonies solid, filled throughout with cells, cells in large but indefinite numbers united into irregular colonies The species *Ascococcus Billrothii* Cohn is discussed Hansgirk (1888, p. 266) recognized *Ascococcus* as the second genus of his subfamily *Cystococcaceae*. The species *A. Billrothii* and a variety *thermophilus* are described by Hansgirk (188, p. 88). This variety was found in warm waters, cells 1μ thick, colorless, spherical or almost egg shaped, united into families, from 6 to 60μ , colorless or yellowish The same species and variety were included by De Toni and Trevisan (1889, p. 1038) They note that the divisions of the cocci occur in three directions, emphasizing thereby that the cells do not occur in chains The genus was included by Billet (1890, p. 24) and by Cornil and Babes (1890 p. 148) Hansgirk (1890, p. 24) named an *Ascococcus cellaris* He believes the *Thioplococcus* of Winogradsky should be regarded as a synonym His description of *A. cellaris* follows

Zellen kugelig oder fast kugelig, sehr viele zu kugeligen oder rundlichen, soliden, 6 bis 20μ breiten Familien, mehr weniger dicht gehaucht, seltener fest zusammengepresst, mit einer gemeinsamen, nicht geschichtlichen Gallert Hülle umgeben, 0.7 bis 1μ breit farblos

Sternberg (1892, p. 17) recognizes the genus *Ascococcus*, and describes *Ascococcus Billrothii* giving as its origin "Found by Billroth in putrefying flesh infusion" Miller (1892, p. 75) described a new species, *Ascococcus buccalis* from the mouth He says

Ein in etwas unregelmässigen Kokken resp. Diplokokken einzeln oder in Ketten vorkommendes Bacterium Dasselbe bildet kleine erhabene glänzende, in alteren Culturen wie winzig kleine Glasperlen aussehende Colonien welche sich besonders da durch auszeichnen dass sie eine knorpelartige consistenz besitzen und sich mit der Nadel nicht heben lassen, sondern vor derselben auf der Oberfläche der Gelatine herumgleiten

Macé (1897, p. 335) designates the genus as *Ascococcus* Billroth Migula (1897, p. 18) likewise concluded that *Ascococcus* is synonymous

with *Micrococcus*, the species becoming *Micrococcus Billrothii* (Cohn) Migula. The species *Ascococcus cantrabridgensis* described by Hankin from the mouth of a student at Cambridge is noted by Lehmann and Neumann (1901, p. 179). Winslow and Rogers (1905, p. 669) revived Cienkowski's name, *Ascococcus mesenteroides* as the type of their emended genus *Ascococcus* (Cohn). Winslow and Rogers. This was characterized later at greater length (1908, p. 136). The Winslows discuss the name *Ascococcus Billrothii* of Cohn. They conclude that because of the cheesy odor and frequent confusion by Cohn of cocci and bacilli, that in all probability Cohn studied a rod-shaped organism. It would seem that they are in error in this matter, for the illustrations accompanying Cohn's description are very distinctive. They also note Cienkowski's use of the term *Ascococcus* and state

Van Tieghem (1878) a little later worked on the same form and substituted for *Ascococcus* the generic name *Leuconostoc* in order to emphasize the resemblance between the zoogloea-forming coccus and the blue-green alga, *Nostoc*.

This would seem to be scarcely a full statement of the case. Van Tieghem concluded that the organism of Cienkowski differed so markedly from Cohn's description of *Ascococcus* that a new generic designation was required. The Winslows emend the diagnosis of *Ascococcus* to conform to modern conceptions. It would seem that they are misled, perhaps, by the apparent appropriateness of the name *Ascococcus*. There seems to be little question but that Cohn's *Ascococcus* was a form entirely distinct from *A. mesenteroides*, and the generic designation should be reserved for Cohn's species. If Van Tieghem was in error in believing that Cienkowski's organism deserved generic separation from *Ascococcus*, then the name used by the latter may be revived. It should be noted that Winslow's designation *Ascococcus* (Cohn) Winslow and Rogers, is incorrect. Article 41 of the International Rules reads "An alteration of the constituent characters or of the circumscription of a group does not warrant the quotation of another author than the one who first published the name or combination of names." The name *Ascococcus* was rejected by Erwin F. Smith (1905, p. 174). Vuillemin (1913, p. 520) lists *Ascococcus Billrothii* 1874 as one of his "*Formgenera conservanda*."

Macé (1913, p. 433, 435, 646) is almost alone among recent writers in recognition of this generic name. He gives an extended description of *A. Billrothii* Cohn and *A. equi* Macé.

Buchanan (1915, p 5) discussed at length the use of the name by Winslow and Rogers. It is probable that Migula's conclusion that Cohn's *Ascococcus* is a growth form of a *Micrococcus* is correct. The species *A. Billrothii* Cohn has never been certainly recognized since described. The status of the genus is quite uncertain. Its type is *A. Billrothii* Cohn.

Ascokokkus. A variant of *Ascococcus*.

Askokokkus. A variant spelling of *Ascococcus* Cohn used by Hueppe (1886, p 145). He includes this as an "Untergattung" of his "Arthrokoccaceen."

This spelling has been used not infrequently by German writers.

Astasia. This name was first applied to a protozoan genus by Ehrenberg. In this sense it is also given by Perty (1852, p 167). More recently the same word has been used as a generic name by Meyer (1897, p 185) for a new species of rod-shaped bacteria, *Astasia asterospora*. Two characteristics of the organism as described by Meyer differentiated it from other genera. The spore is barrel-shaped, ribbed longitudinally, and with a well differentiated extine and intine, the flagella are produced in bunches, laterally. The original description was later modified (1898, p 49). Migula (1900, p 528) regards *Astasia* as a synonym of *Bacillus* and writes *Bacillus asterosporus* (Meyer) Mig. The organism is described by Migula as a motile rod usually single, seldom in short chains, 1 to 1.3 μ thick, 3 to 6 μ long, becoming motionless before spore formation. Spores ovoid, with longitudinal striae, star-shaped in cross section, with polar germination. Flagella distributed over the entire body. On sliced beet slices it develops as a gray, glassy, gelatinous layer, which spreads so that after five days the slice is covered with a thin gelatinous layer, finally showing gas bubbles. The middle lamellae of the beet are dissolved so that the medium becomes soft. Spores are found after five days, in addition to resting and swarming rods. In dextrose gelatin rapid liquefaction occurs, with some gas bubbles. It grows in nutrient solution, causing a decided clouding. Migula groups this organism with *Bacillus subtilis*, *B. ramosus*, etc. The cultural characteristics with the exception of gas production seem to relate it quite definitely to these forms.

Buchanan (1918, p 38) used this name as a subgeneric designation under the genus *Bacillus*, with the description "Motile rods, spores ovoid with longitudinal stripes, star shaped in cross section." The type is designated as *Bacillus (Astasia) asterospora* (Meyer) Migula.

Asterococcus. A generic name proposed by Borrel, Dujardin-Beaumetz, Jeantet and Jouan (1910, p 179) for the organism causing

bovine peripneumonia. The single species was named *Asterococcus mycoides*. The genus may be characterized as follows. Pleomorphic cells, appearing at different stages as cocci isolated and in chains, and as rods and filaments variously branched and swollen, apparently as involution forms resembling frequently the bacteroids of legumes, cells extremely minute, almost ultramicroscopic. Non-motile. No spores. Stain with difficulty, best with Giemsa. Growth in cultures only in presence of serum or hemoglobin. Apparently the organism has not again been seen or recorded.

The genus was recognized by Buchanan (1918, p. 44) and included in his subtribe *Hemophilinae*.

Astrobacter. A generic name given by Jennings (1896, p. 312) to an organism found in stagnant water near Rubingen by Coppen-Jones. Slides stained by Loeffler's method to demonstrate the flagella of *Spirillum undula* when examined were found to show the organism in question. The original material had been lost, so that the entire description is based upon observation of the mounted slides.

A simple rod-like bacterium. Initial cell division is marked by a bifurcation of the end of the rod, producing a Y-shaped form. The split deepens, and others occur, making tri-radiate types with the rays spreading at an angle of 120° . Four rayed forms with various angles occur, as do also forms with as many as eight rays. The rays do not appear to all lie in a single plane. In a few cases transverse constrictions of some of the rays were noted. Whether this genus has been based upon any form which actually occurs in nature has not been proved. It is entirely possible that the so-called organisms are artefacts of some type. Until there is a demonstration of the living organisms the generic name *Astrobacter* Jennings may well be held in abeyance. It has apparently never been recognized or used by other investigators. It is definitely rejected by Erwin F. Smith (1905).

Athiorhodaceae. A family of the order *Rhodobacteria* proposed by Molisch (1907, p. 28) to include those forms that do not possess free sulfur granules in the cell contents. The following genera, described in the same work, make up this family: *Rhodobacillus*, *Rhodobacterium*, *Rhodocapsa*, *Rhodotheca*, *Rhodococcus*, *Rhodovibrio*, *Rhodocystis*, *Rhodonostoc*, and *Rhodospirillum*. The organisms all develop in the presence of light. They possess two pigments called bacteriopurpurin and bacteriochlorin.

The family name does not conform to the botanical code in that it is not derived from the name of any one of the component genera. It is made a synonym of *Rhodobacterioidae* by Buchanan (1918, p. 128).

Aurococcus. A generic name proposed by Winslow and Rogers (1906, p. 540) to include the orange cocci. The generic diagnosis given is:

Parasites. Cells in groups and short chains, very rarely in packets. Generally stain by Gram. On agar streak growth of orange color. Sugars fermented with formation of small amount of acid. Gelatin often liquefied very actively. May or may not reduce nitrates. Includes *A. aureus* (Rosenbach).

The same genus was later discussed by the Winslows (1908, p. 255). They include three species, *Aurococcus aureus* (Rosenbach) Winslow, *Aur. aurantiacus* (Schroter, Cohn) Winslow, and *Aur. mollis* (Dyar) Winslow.

For the generic name *Aurococcus* Winslow and Rogers to be established as valid, it must be shown that the generic name *Staphylococcus* which is displaced is invalid. The latter name was proposed by Ogston (1883, p. 27), but first used as a generic name by Rosenbach (1884, p. 12) who described a *Staphylococcus pyogenes aureus* and a *Staphylococcus pyogenes albus*. The former is the type chosen for *Aurococcus* Winslow and Rogers. On a later page, Rosenbach uses the combination *Staphylococcus aureus* for the same organism. The genus *Staphylococcus* Rosenbach is split by Winslows into the two genera *Aurococcus* and *Albococcus*, and the original generic name discarded. Article 45 of the International Rules of Botanical Nomenclature states,

When a genus is divided into two or more genera, the name must be kept and given to one of the principal divisions. If the genus contains a section or some other division which, judging by its name or its species, is the type or the origin of the group, the name is reserved for that part of it.

It would seem that the Winslows have shown no adequate nomenclatural reason for abandoning the generic name *Staphylococcus*. It should, therefore, be retained for one of their genera. Inasmuch as *Staphylococcus aureus* was described first, and is in a sense the type of the group, the name *Aurococcus* should be reduced to a synonym of *Staphylococcus*. This genus can be emended, if desired, to exclude the white forms, thus making the genus *Albococcus* valid.

These facts were pointed out by Buchanan (1915, p. 8) and agreed to by the Committee of the Society of American Bacteriologists (1917, 1920) and by Winslow, Rothberg and Parsons (1920, p. 161).

Azotobacter. A genus of bacteria created by Beijerinck (1901, p. 561) to include certain non-symbiotic nitrogen fixing organisms of the soil. Beijerinck gives the following diagnosis of the genus:

Dicke in jungem Zustande meist als grosse Diplokokken oder als Kurzstäbchen vorkommende Bakterien mit hyalinem, oft eine Vakuole fuhren dem Inhalte und schleimiger Wand von sehr verschiedenen Dicke Jungere Zustände mehr oder weniger beweglich vermittelt einzelner polarer oder in polaren vier—bis zehnzahligen Bündeln angeordneten, kurzen Cilien, welche umgefah so lang sind, wie die Bakterien selbst Sporen fehlen, das heisst, wachstumfähig in Nahrlosungen mit geeigneten Kohlenstoff quellen, welche sehr arm sind an Stickstoffverbindungen assimiliert den atmosphaerischen Stickstoff, und ist dadurch in Bezug auf die ubrige Mikrowelt konkurrenzfähig Optimum temperature 28°

The type species named was *Azotobacter chroococcum*

Beijerinck and van Delden (1902, p 3) again described the genus. The genus was rejected by Erwin F Smith (1915) It was accepted by Stockhausen (1907, p 88) The genus is quite generally recognized by writers on agricultural bacteriology Orla-Jensen (1909, p 328) changes the name of this genus to *Azotomonas* and places it in his family *Oxydobacteriaceae*

Lohnis and Hanzawa (1914, p 1) claim to have demonstrated that the organisms of this genus may at certain stages in their life history produce endospores They conclude that *Azotobacter* is therefore an invalid synonym of *Bacillus* The type species is renamed *Bacillus azotobacter* It may be noted that even though the genus should be *Bacillus*, the species should be designated *B chroococcus* in accordance with the botanical rules

Winslow *et al* (Committee 1917, p 552) and Bergey *et al* (1923, p 38) accept the genus with the following diagnosis

Relatively large rods, or even cocci, sometimes almost yeast-like in appearance, dependent primarily for growth energy upon the oxidation of carbohydrates Motile or non-motile, when motile, with tuft of polar flagella Obligate aerobes usually growing in a film upon the surface of the culture medium Capable of fixing atmospheric nitrogen when grown in solutions containing carbohydrates and deficient in combined nitrogen The best-known free-living nitrogen-fixing bacteria of the soil

The same description is used by the Committee in 1920 (p 203) Buchanan (1918, p 46) describes the genus as follows

Relatively large rods, or even cocci, sometimes almost yeast-like in appearance, dependent primarily for growth energy upon the oxidation of carbohydrates, obligate aerobes, usually growing in a film upon the surface of the culture media Capable of fixing atmospheric nitrogen in considerable amounts when grown in solutions deficient in combined nitrogen Motile or non-motile, if the latter, with polar flagella

It is probable that *Azotobacter* Beijerinck may be regarded as a valid genus.

Azotobacteraceae. A tribe of the *Nitrobacteriaceae* suggested by Winslow *et al.* (Committee Soc. Am. Bact., 1920, p. 203), to include the "nitrogen fixing bacteria." The genera *Rhizobium* and *Azotobacter* were included. Bergey *et al.* (1923, p. 38) include this as the second tribe of the family *Nitrobacteriaceae*. Probably the spelling should be emended to *Azotobacteriaceae*.

Azotomonas. This generic name was proposed by Orla-Jensen (1909, p. 484) to replace the older name *Azotobacter* Beijerinck (1901, p. 561) to conform to his rule that organisms with polar flagella should bear generic names terminating in *-monas*. It should be regarded as invalid because of the priority of *Azotobacter*.

Babesia. A genus of bacteria created by Trevisan (1889, p. 29) to include two species of cocci, one associated with yellow fever and described by Babes (1883,) and the other with *erysipelas chronica*. It was included as the fourth genus of his tribe *Streptococceae*. The diagnosis given by De Toni and Trevisan (1889, p. 1054) is as follows.

Cocci ellipsoidei, longitudinaliter binati seriati (diplococci longitudinales) in filamenta moniliformia, pseudodichotoma nuda (i. e., nec capsulis nec vaginis obducta) concatenati. Arthrospora macrosomae in apice filamentorum obvenientes

The genus is differentiated from *Streptococcus* by the pseudodichotomous branching of the filaments and by the production of arthrospores at the ends of the filaments. The generic description corresponds with that of no organism at present known. The two species, *B. xanthopyretica* Trev. and *B. erysipeloides* Trev. were so inadequately described as not to be recognizable. The genus name has never come into use. It may be regarded as invalid or possibly as a synonym of *Streptococcus* (Erwin F. Smith 1905, p. 174).

Babesia xanthopyretica Trev. (*Streptococcus xanthopyreticus* Trev.) may be regarded as the type.

Babesia is also used as a generic name in the pathogenic protozoa.

Bacillaceae. A family proposed by Fischer (1895, p. 139) with the spelling *Bacillacei* and with the following diagnosis

Vegetationskörper einzellig, gerade mit ausgesprochener Längsachse, bald kurz ellipsoidisch, bald gestreckt stabchenförmig. Theilung immer in derselben Richtung, senkrecht zur Längsachse mit oder ohne Kettenwuchs und Bewegung. Sporen theils endospor, theils arthrospor.

In the translation of the Vorlesungen by Coppen Jones (1897, p 32) this is spelled *Bacillaceae*. In the second edition (1903, p 60) the diagnosis given is

Vegetationskörper cylindrisch, ellipsoidisch, eiförmig, gerade, bei den kurzen, fast kugelförmigen Formen und die Trennung von Kokken schwer, Teilung immer senkrecht zur Längsachse, als Wuchsform nur unverzweigte Ketten

In the original description four subfamilies are listed *Bacillaei*, *Bacterineae*, *Bacillulaceae* and *Bacillulidae*. Later (1903) the three subfamilies recognized were *Bacillaceae* (with four genera) *Clostridiaceae* (two genera) and *Plectridiaceae* (two genera)

Schmidt and Weis (1902, p 91) likewise used this family designation, including the three genera *Bacterium*, *Bacillus* and *Pseudomonas* with the following description

Die Zellen sind cylindrisch, kürzer oder länger stabchenförmig, gerade. Die Zellteilungen geschehen nur nach einer Richtung des Raumes. Vor der Teilung strecken sich die Zellen in einer senkrechten Richtung zu der später gebildeten Scheidewand. Endosporbildung bei mehreren Arten bekannt

The name has also been used as a group designation by Flugge (1908)

This family name is also used by Benecke (1912, p 188) to include the rod-shaped bacteria without sulphur or bacteriopurpurin, unbranched and not forming a pseudoplasmodium

Winslow *et al* (Committee Soc Am Bact, 1917, p 561) used this as a designation for their seventh family of the *Eubacteriales*. The diagnosis is "Rods producing endospores, usually Gram-positive. Flagella when present peritrichic. Actively decompose protein media through the agency of enzymes"

Winslow *et al* (Committee Am Soc Bact, 1920, p 212) in their final report used the same diagnosis

Castellani and Chalmers (1920, p 601) state

Cette famille peut être définie Eubacteriales avec cellules longues ou courtes, flagellées ou non flagellées, sporogènes ou non sporogènes, mais toujours cylindriques et droites et se divisant seulement dans une direction

Le genre type est *Bacillus* Cohn 1872, mais ce genre et l'autre genre ancien, *Bacterium*, renferment un si grand nombre d'espèces et de variétés que nous avons essayé de simplifier leur diagnose en définissant les tribus et les genres suivant

They divide the family into 10 tribes

Bergey *et al* (1923, p 272) included this as the fifth family of the order *Eubacteriales* with the description "Rods producing endospores, usually

Gram-positive Flagella, when present, peritrichous Often decompose protein media actively through the agency of enzymes "

Two genera, *Bacillus* and *Clostridium* are included

Bacillacei. A family name proposed by Fischer (1895, p 139) For discussion, see *Bacillaceae* (the corrected form)

Bacilleae. A tribe established by De Toni and Trevisan (1889, p 941) with the following diagnosis "*Baculi cocci nunquam capsulis membranaceogelatinosis involuti* " It is the first tribe of the subfamily *Baculogenae* Trev. It includes the sub-tribes *Pasteureae* (one genus) *Thiodictyeae* (one genus), *Eubacilleae* (six genera) and *Spirilleae* (two genera)

With the spelling *Bacillei* this name was also used by Fischer (1895, p 140) as a subfamily of the family *Bacillacei* to include all rod-shaped organisms "Unbeweglich, ohne Geisseln " Four genera were included, *Bacillus*, *Bactrinum*, *Bactrillum* and *Bactridium*.

The name was used by Buchanan (1918, p 33) as a tribe of the family *Bacteriaceae* with the following description

Cells rod shaped, never spiral or strictly filamentous, single or in chains, usually motile by means of peritrichous flagella, producing endospores under suitable conditions of growth, usually Gram-positive

The four genera *Bacillus* *Plectridium*, *Clostridium* and *Metabacterium* were included

Castellani and Chalmers (1919, pp 933, 959) propose this as a tribe of the family *Bacillaceae* Apparently they are in error in ascribing the name to Castellani and Chalmers 1918 They state "*Bacillaceae* growing well on ordinary laboratory media and producing endospores One genus only is included *Bacillus*, the type genus " They again describe it in 1920 (p 601)

Bacillee A name given by Trevisan (1889, p 12) to the third tribe of the sub-order *Baculogene* It is ascribed to Trevisan (1887) The description given is

"*Baculi, filamenti e cocci non mai racchiusi in capsule membranaceo-gelatinose.*" Twelve genera are included *Pasteuria*, *Winogradskyia*, *Thiodictyon*, *Mantegazzaea*, *Bacillus*, *Pasteurella*, *Cornelia*, *Clostridium*, *Vibrio*, *Pacinia*, *Spirillum* and *Spiromonas*

Bacillei A subfamily name incorrectly spelled used by Fischer (1895, p. 140) For discussion, see *Bacillaceae*

Bacilococcus. A casual name used by Frankland (1890, p 122) for a nitrifying soil organism

Bacillopsis. A generic name suggested by Petschenko (1908, p. 359) for an organism questionably bacterial Enlows (1920, p. 19) states

Type species (monotypy) B stylopygae Found in the digestive tube of *Blatta orientalis* (*Stylopyga orientalis*) Length 10μ , by 2.5μ wide A slightly curved rod, with one end slightly pointed, the other obtuse Nucleus is present Also highly refractive "corpuscles" in the transparent protoplasm, which are probably nutritive substances Reproduction by a sort of budding, in which the very small daughter cell remains attached to the mother cell by a delicate filament until it has attained the size of mother cell After this stage of active growth there is a stage in which filiform prolongations appear, and the "corpuscles" unite into 1 or 2, rarely 3 large round bodies Vacuoles observed In doubt as to position of organism, but does not think it belongs with the bacteria (Has been included by other authors among the bacteria)

Bacillus. A generic name established by Cohn (1872, p. 174) to include three species of rod-shaped organisms, *Bacillus subtilis*, *B. ulna* and *B. anthracis* He included in this genus those rod-shaped organisms that grow in filaments The type of the genus was *B. subtilis* A discussion by Cohn of spore production in *Bacillus subtilis* embodied the first accurate description of endospores His inclusion of the anthrax organism indicates that in his conception of the genus motility was an unimportant characteristic In a later paper, Cohn (1875, p. 141) included the genus *Bacillus* under the tribe *Nematogenes* and characterized it as follows

Cells disposed in filaments, filaments not branched, free or interlaced, cylindrical, colorless, articulations not very distinct, very slender and short

It will be noted that in the preceding description emphasis is placed upon the occurrence of the rods in short chains or filaments Sporulation and motility were not included in the generic diagnosis Within the genus were listed both motile and non-motile forms Sporulation was discussed in certain of these species, but not rated definitely as a generic character These facts are emphasized because subsequent generic designations frequently stress other characteristics

Winter (1880, p. 38) uses the genus practically in the sense of Cohn He states

Zellen verlängert cylindrische, fast stets in geraden, stielrunden (nicht oder wenig eingeschnurten) Reihen oder Faden zusammenhängend, durch Quertheilung sich vermehrend Sie bilden Zoogloeen, kommen aber oft auch in dichten Schwärmen (ohne Gallertausscheidung) vereinigt vor Fortpflanzung durch Sporen

This is the first definite inclusion of the character of spore production in the generic diagnosis, but this should not be emphasized as a differential character, for in the description of *Bacterium* there is the statement "Sporenbildung ähnlich wie bei *Bacillus* "

Luerssen (1879, p. 22) accepted Cohn's classification practically without modification. Emphasis is placed upon the occurrence of the organism in jointed threads, and no mention is made of sporulation. Magnin (1880, p. 87) used the same description.

De Bary (1884) influenced largely by the idea of bacterial pleomorphism, abandoned the old genus *Bacterium* and put all endosporous rods into the genus *Bacillus* and all arthrosporous rods and those for which no spores are known into *Arthrobacterium*. De Bary states (translation by Garnsey and Balfour 1887, p. 460)

The forms included under this term (endosporous bacteria) are chiefly known in the growth-form of single rods consisting of one or a few cells, or of rods joined together and forming long filaments, they may also be collected together into larger gelatinous masses or membranes. In some forms the rods are spirally twisted, and these I name here *Spirillum* of Van Tieghem. Others do not show these curvatures, but are either straight or very slightly bent, all these I include under the term *Bacillus* and place under that genus all the endosporous forms which have been hitherto known either as *Bacillus* or as *Clusridium*, *Bacteridium*, *Vibrio*, or by some other name. All non-endosporous forms bearing these names on account of their growth form are, of course, excluded from the group.

It will be noted that this is a direct abandonment of the generic conception of Cohn although the type species may well be said still to be *B. subtilis*.

Van Tieghem (1884, p. 1110) uses the term *Bacillus* as a form group to include those rods which are not associated but are united into filaments of greater or less length, practically in the sense of Cohn. Crove (1884, p. 26) follows Winter minutely. Trevisan (1885, p. 94) included in this genus those organisms showing three stages of development, as rods, filaments and cocci. In the rod or baculus stage the cells are normally cylindric, or ellipsoid, straight or slightly curved, jointed or not, colorless or colored, cytoplasm equally distributed. Spores single in the rods. Filaments and cocci transitory stages. He included descriptions of thirty-one species, the first one described being *Bacillus anthracis* Cohn. The views of Van Tieghem were in part accepted by Zopf in the third edition of "Die Spaltpilze" (1885, p. 61). In previous editions the term *Bacillus* was regarded merely as the designation of a growth form of a *Bacterium*. His diagnosis in the third edition is

"Coccen und Stabchenformen oder auch nur Stabchen in gewöhnlichen oder gewundenen Faden Sporenbildung vorhanden in Stabchen oder in Coccen auftretend" Although the definition is greatly modified by the current ideas of bacterial pleomorphism the emphasis is placed upon the development of endospores by the members of the genus Huetpe (1885) likewise included in the genus *Bacillus* those rods that developed endospores

Flügge, (1886, p 136) used "Bacillen" to include rods whose length was two to twenty times the diameter, returning in a sense to the conception of Cohn

Maggi (1886, p 173) included this genus, without description, in his fourth order, *Desmobacteries* Eleven species are listed

Schroeter (1886, p 156) considers the genus *Bacillus* to include organisms with long cylindric vegetative cells, daughter cells often remaining united for a time in a chain, often actively motile, with endogenous spores produced singly in cells that do not become swollen when sporulating In some species the cells grow into long filaments just preceding sporulation (*Streptobacter*) In a rather loose fashion a somewhat similar meaning is given by Cornil and Babes (1890, p 165), these authors emphasize sporulation Hansgug (1888, p 264) includes *Bacillus* as the first genus of the subfamily *Microbacteria* He divides the genus into two sections or subgenera, *Eubacillus* and *Chromobacillus* Trevisan (1889, p 12) included *Bacillus* as the fourth genus of the tribe *Bacillie* The generic description is "Baculi cilindrici, cylindracei o ellipsoidi, con protoplasma uniformemente diffuso Spore microsome (cioe che non mai raggingono un diametro maggiore del diametro trasversale normale dei baculi entro cui si formano) provenienti in baculi normali immutati" The generic synonyms are given as "*Metallacter* e *Chromatium* Perty (1852), *Bacteridium* Davaine (1863), *Bacillus* e *Bacterium* F Cohn (1873), *Streptobacteria* Billroth (1874), *Pollendera* Trevisan (1884), *Coccobacillus* Leube (1885), *Proteus* Hauser (1885), *Bacteriopsis* Trevisan (1885), *Coccothrix* Lutz (1886), *Sclerothrix* Metschnikoff (1888) In all 201 species were listed De Toni and Trevisan (1889, p 1889) elaborated the diagnosis of *Bacillus* somewhat to allow of the differentiation from it of numerous related genera Their diagnosis is as follows

Baculi cylindrici vel cylindracei, recti vel leviter curvi, apicibus conformibus rotundatis vel truncatis, plasmate uniformiter diffuso Filamenta vulgarissima e baculis, coccis nullis interjectis conflata Sporae (endosporae) microsomae, in baculis normalibus immutatis obvenientes

A large number (228) of species are described Billet (1890, p 23) described *Bacillus* as "Formes rectilignes, element 5 a 10 fois plus long que large"

Baumgarten (1890) was apparently the first to use the generic name *Bacillus* to include all rod-shaped, "monomorphic" bacteria, practically merging with it the genus *Bacterium* This conception was adopted by Sternberg (1892, p 18) who defined the genus as follows

Rod-shaped and filamentous (not spiral) bacteria in which there is no differentiation between the extremities of the rods, reproduction by binary division in a direction transverse to the long axis of the rods, or by binary division and the formation of endogenous spores, rigid or flexible, motile or non-motile

This use has led to the rapid "vulgarization" of the term *Bacillus* until it is frequently now noted as being used in two senses, a general sense with reference to shape, and as a genus name This has contributed to much confusion in the literature

Ludwig (1892) defines the genus as containing small rods which produce endogenous spores

Clements (1894, p 27) lists four species, with the following generic description

Cells cylindrical, straight or curved, motile filaments present at time of spore formation, falling into small cells as soon as the endogenous spore is formed

Migula (1894, p 237) emended the genus *Bacillus* Cohn, the diagnosis reading "Zellen mit über den ganzen Körper angehefteten Bewegungsorganen, oft mit Endosporenbildung (Z B *Bacillus subtilis* Cohn)" Migula (1895, p 25) later elaborated this conception He based his principal generic designations upon differences in flagellation Accepting *B subtilis* as the generic type, he characterized the genus as made up of straight rods with flagella scattered over the entire cell body Sporulation and filament formation he considered secondary to flagellation

The second emendation in 1895 of *Bacillus* was that of Fischer (1895 p 140) This author rejects Migula's use and states

Unbeweglich, ohne Geisseln, mit Endosporen in unveränderten, nicht spindeligen oder kopfigen Stäbchen, eine schwache allseitige Vergrösserung der Stäbchen, bei der Sporenbildung kommt zuweilen vor Sporen in der Mitte oder am Ende der Stäbchen, *Bacillus anthracis* Cohn *Bacillus Carotinum* A Koch

Later (1897, p 32) he says

The word *Bacillus* might then be used, in memory of Koch's first work for all those species which, like the anthrax parasite, are non-motile and retain their shape during sporulation

This of course constitutes a direct abandonment of *Bacillus subtilis* as the generic type

Lehmann and Neumann (1896, p 279) returned to the differentiation of the genus upon the basis of endospore production Their diagnosis reads "Gerade Stabchen haufig zu Faden auswachsend Dicke oft betrachtlich, selten unter 0.6, meist uber 0.8 μ Endosporen bildend"

Migula later expanded his classification (1897, p 47) and again emphasized the peritrichous flagellation of the cells as the principal diagnostic character Macé (1897, p 484) used as a diagnosis "Elements in the form of rods, short or long, straight or slightly bent" He concludes that the *Bacteridium* of Davaine, the *Clostridium* of Prazmowsky, the *Tyrophium* of Duclaux, the *Proteus* of Hauser are synonyms Hewlett (1898, p 18) includes all rod-shaped bacteria under the genus *Bacillus* Migula (1900, p 515) again defined *Bacillus* as made up of peritrichous rods He says

Kurzere oder langere, stabchenformige bis ovoide gerade Zellen, oft zu ziemlich langen Faden verbunden, beweglich, mit uber den ganzen Korper zerstraut stehenden, wellig gebogenen Geisseln Endosporenbildung ist haufig

Lehmann and Neumann (1901, p 124) consistently adhere to the diagnostic importance of endospores, and include all spore bearing rods in this genus Chester (1901, p 199) follows Migula in basing his diagnosis upon peritrichous flagellation The same description is used by Kendall (1902, p 484) Schmidt and Weis (1902, p 91) follow Migula Matzuschita (1902, p IV) included all rod-shaped organisms Allan J Smith (1902, p 270) gives the generic description "Cells with flagella, peritrichous, endospores present or absent" Fischer in the second edition of his lectures (1903, p 60) again defines *Bacillus* as comprising the non-motile rods Migula (1904, p 145) reaffirmed peritrichic flagellation as important He is followed by Erwin F Smith (1905, p 160), Ellis (1909, p 6), Frost (1911, p 57), Schneider (1912, p 23) and many other authors Orla-Jensen (1909, p 344) used the name *Bacillus* to designate organisms of the type of *B. mycoides*, placing the genus in his family *Alkalibacteriaceae* These organisms are said to require organic nitrogen, free oxygen, produce spores and cause decay Conn (1909,

p 12) apparently introduced still another conception of *Bacillus* when he included in the genus all "Rod-shaped bacteria with flagella and consequently motile"

Heim (1911, p 250) and Lohnis (1913, p 45) included in *Bacillus* the spore-forming rods Vuillemin (1913, p 526) comes to the conclusion that the name *Bacillus* has been so vulgarized by various usages among bacteriologists, that it should be regarded merely as a form genus without nomenclatural status As a substitute for this generic designation he revives the name *Serratia* (q r) and gives as the type *Serratia subtilis* Vuillemin

Winslow *et al* (Committee Soc Am Bact Preliminary Report, 1917, p 562) included the spore bearing rods in this genus with the following diagnosis "Aerobic forms Mostly saprophytes Liquefy gelatin. Often occur in long threads and form rhizoid colonies Form of rod usually not greatly changed at sporulation" *Bacillus subtilis* Cohn was designated as the type

Buchanan (1918, p 33) discussed in detail the generic designation, and defined the genus as follows

Cells rod shaped, straight or at least never spiral motile by diffuse flagella or non-motile Endospores produced under favorable conditions, not usually distorting the cell, usually Gram-positive Growth usually good on laboratory media, commonly liquefying gelatin Aerobic or facultative

Three subgenera were recognized, *Eu-Bacillus*, *Bacterium*, and *Astasia*

Winslow *et al* (Committee Soc Am Bact 1920, p 212) in their final report use the same diagnosis as in 1917 This is also used by Berg *et al* (1923, p 272)

The usages of various authors may be summarized as follows

1 *Bacillus* Rod-shaped organisms occurring in filaments or chains Spore production flagella distribution and motility not emphasized Cohn (1872, and 1875) Magnus (1878) Winter (1879) Lueissen (1879)? Van Tieghem (1884) Grove (1884) Flugge (1886) Schroeter (1886)

2 *Bacillus* Rods producing endospores (some authors recognizing other spore-bearing genera in addition) De Bary (1884 and 1887), Zopf (1885), Hueppe (1885), Cornil and Babes (1885 and 1890), De Toni and Trevisan (1889), Ludwig (1892), Freudenreich (1894), Lehmann and Neumann (1896), Chester (1897), Flugge (1908), Orla-Jensen (1909), Heim (1911), Lohnis (1913), Winslow *et al* (1917, 1920), Buchanan (1918), Bergey (1923)

3 *Bacillus* Rods motile by means of peritrichous flagella Migula (1895, 1897, 1904), Chester (1901), Kendall (1901), A J Smith (1902), E F Smith (1905), Ellis (1909), Frost (1911), Schneider (1912)

4 *Bacillus* Non-motile rods, cylindrical, producing endospores. Fischer (1895, 1903), Lotsy (1907)

5 *Bacillus* Any rod-shaped organism Baumgarten (1890), Sternberg (1892), Macé (1897), Hewlett (1898), Matzuschita (1902)

6 *Bacillus* Any motile rod Conn (1909)

It would seem that the genus name *Bacillus* should be retained for the designation of a limited group of organisms, though probably no harm can come from the use of the term *bacillus* as a casual name to indicate rod-shaped organisms in general. It should be emphasized that when used as a generic designation it is a proper noun, and *must be capitalized*, whenever used. Many bacteriologists have ignored this rule which is followed so carefully by systematists and biologists in treating all other forms of life. Vuillemin's contention that the use of *bacillus* as a vulgar noun leads to confusion is not well premised. The genus *Aster* is universally recognized by botanists as valid. This does not interfere with the use and usefulness of the common name *aster* which includes many species not belonging to the genus *Aster*. By no means all of the plants commonly known as lilies are placed by the botanist in the genus *Lilium*. There is no reason why we cannot continue to speak of the tubercle bacillus even though it does not belong to the bacterial genus *Bacillus*.

In some form or with some definition the genus *Bacillus* should be retained. The type practically always accepted is *B. subtilis*. The definition of Fischer should therefore be abandoned as including only non-motile forms. He would exclude from the genus its first described species. The original description of Cohn is scarcely sufficient, for much stress was laid upon cell grouping and length of cell and not upon other characters. The use of Migula's diagnosis, including in the genus all rods with peritrichous flagella, is the cause of great confusion. It brings into the genus such discordant types as the hay bacillus and the typhoid bacillus. It excludes from the genus the anthrax bacillus so closely related to the hay bacillus. It does not result in bringing forms that are closely related together. Migula's definition should be abandoned as not based upon natural affinities. The definitions which would include all rods in the genus *Bacillus* have the merit of simplicity, but when organisms so diverse in characteristics as the tubercle bacillus, the typhoid bacillus, the tetanus bacillus, and the anthrax bacillus, are

all included in one genus the simplicity is more apparent than real. The existence of such diverse forms has led most recent authors to arrange bacteria into well marked groups. It is the opinion of the author that the larger groups should be recognized as genera. The term *Bacillus* should therefore be restricted. It would seem that it should be defined more nearly in the terms of De Bary, Zopf, Hueppe, etc., who emphasized the importance of spore production as a diagnostic character.

Objection may be raised that a definition of *Bacillus* as a genus made up of endosporous rods would exclude forms which have lost the power of spore formation but are in other respects closely related. It is evidently impracticable to base generic diagnoses upon a single character. Even though an organism be a variant in one or even more characters, the other resemblances would be sufficient to include the organism in question in the correct genus. Illustrations of this fact may be taken from higher plants. The Lombardy poplar is always classified in the genus *Populus*. It never produces fruit, it persists solely as the result of vegetative multiplication. Yet the genus *Populus* is based in part upon certain fruit characters. The other characters are so evidently poplar-like, however, that we do not question the correctness of the assignment of this species to the genus *Populus*.

Bactereae. A name given to the fourth tribe of the *Bacteriaceae* by Winslow *et al* (Committee Soc Am Bact 1910, p 210). It is evidently a typographic error for *Bacteriense* q v. This spelling is also used by Bergey *et al* (1923, p 194).

Bacteria. A group name frequently used by authors as synonymous with *Schizomycetes*. Hansgig (1888, p 299) gave the two as synonyms. Migula used this name with the following diagnosis:

Phycochromfreie Spaltpilzen mit Teilung nach ein- oder dreier Richtungen des Raumes. Viele Arten besetzen Endosporenbildung. Wo Beweglichkeit der Zellen vorhanden ist, wird dieselbe durch geißelartige Bewegungsorgane, seltener durch undulierende Membranen (Uebergang zu den Phycochromaceen) vermittelt.

The name in this general sense has been used by many writers. Heller (1912, p 449) proposes this as the name of a phylum.

Bacteriaceae. Cohn (1872a, p 237) under the description of *Microsphaera* uses the expression "Gruppe der *Bacteriaceae*". Cohn (1872b) says:

Den gemeinschaftliche Charakter der von mir hier als Bacterien zusammengefassten Organismen scheint nur in Folgen dem zu liegen. Die Bacterien sind

chlorophyllose Zellen von kugliger, oblonger oder cylindrischen, mitunter gedrehter oder gekrümmter Gestalt, welche ausschliesslich durch Quertheilung sich vermehren, und entweder isolirt oder in Zellfamilien vegetiren

Apparently the name is used for the entire group of *Schizomycetes*

The next reference to a use of this name in literature is apparently a statement of Lanzi (1876, p 25) who mentions that Prof Caruel in his lectures includes the family "Bacteriacee" among the "Schizofiti" It was used in the next year by MacNab (1877, p 340) and in much the same sense

Trevisan (1879, p 135) used the following description of this family

Plantae fungales, unicellulares vel pluricellulares, liberae aut in muco matricali amorpho nidulantes, nunc motu proprio spontaneo, vel oscillante, vel rotatorio, vel flexuoso-repente, vel spirali praeditae, nunc immobiles, acidum carbonicum non assimilantes, in aquis plus minusque corruptis, dulcibus, thermalibus vel marinis, in liquidis fermentantibus, in substantiis putrescentibus, in corporibus animalium viventes Vegetatio terminalis vel non terminalis Ramificatio nulla aut rarissime spuria Cytoderma flexibile, molle, tenuissimum Cytio-plasma chlorophyllo vel substantia chlorophyllo affini omnino carens, achroum vel coloratum, nucleo destitutum Multiplicatio divisione cellularum vegetativa aut in omnes directiones aut semper ad eandem directionem longitudinalem repetita Propagatio tum sporis perdurantibus sine foecundatione e cytioplasmatis condensati metamorphosi genitis, tum microgoniis, in paucis detectis, e serie cellularum divisione longitudinali et transversa succedentiae multipartitarum ortis Per divisionem formae conservantur, multiplicantur et generationum series extensissimas evolvuntur, per sporas et microgonia morphae non raro parentibus plus minus dissimiles, colonias mucosas per cellularum generationes successivas ortas formantes, nascuntur

This family name with the aberrant spelling *Bacteriaceen* was also used by Zopf (1883, p 45) He included within it all organisms which might show four developmental stages, cocci, short rods (bacteria), long rods (bacilli) and filaments (*Leptothrix* forms) There is no differentiation of base and tip in the rods Typical spiral forms are lacking Two genera were recognized *Bacterium* and *Clostridium* In the third edition of Die Spaltpilze he (1885, p 51) changes his conception of the family essentially He now defines it to include organisms which show spherical, rod or filamentous growth forms, the latter either straight or bent The coccus forms may be lacking In the rod and filamentous forms there is no contrast between base and tip Cell division, so far as is known, is in one direction of space Spores present, lacking, or unknown Six genera are recognized, *Bacterium*, *Spirillum*, *Vibrio*, *Leuconostoc*, *Bacillus* and *Clostridium* Schroter (1886, p 155) uses

the family name *Bacteriaceae* Zopf as the sole family of his order *Eubacteria*. He includes nine genera *Bacterium*, *Chromatium*, *Bacillus*, *Clostridium*, *Spirillum*, *Spirochaete*, *Microspira*, *Myconostoc* and *Cystobacter*. Hansgirk (1888, p. 264) included *Bacteriaceae* as the first family of his order *Eubacteria*, but with no description. It is divided into the two subfamilies *Spirobacteria* and *Microbacteria*.

The classification of Hueppe (1891, p. 30) included two genera only, *Bacterium* and *Bacillus*. Migula (1894, p. 236) says

Bacteriaceae Zopf emend Mig. Zellen langes oder kurzer cylindrisch, gerade, niemals schraubig, gekrümmt. Teilung nur nach einer Richtung des Raumes nach vorausgegangener Langstreckung des Stäbchens.

He (1895, p. 20) included three genera, excluding all spherical, spiral and filamentous forms. The genera recognized were *Bacillus*, *Bacterium* and *Pseudomonas*. Later (1897, p. 46) he gave the same definition.

Clements (1894, p. 25) redefined the family as follows

Minute fungi without mycelium, typically unicellular, or at least the divisions not visible except at the time of the abscission of new cells, cells of very diverse form (round, elliptical, bacillar, or filamentous), generally surrounded by a gelatinous envelope, one or both ends often provided with 1 to 3 flagella, growth peripheral, never apical, sexual reproduction undeveloped, asexual reproduction by simple division or by spores (endo-spores or arthrospores).

Lehmann and Neumann (1893, p. 103) gave the following definition

Zellen mindestens 1½ mal, meist aber 2-6 mal so lang als breit, gerade oder in nur einer Ebene etwas gekrümmt, nie schraubig, zu weilen lange echte oder Scheinfäden bildend. Teilung (ast, stets quer zur die Längsachse nach Streckung des Stäbchens. Mit oder ohne Geisseln. Mit oder ohne Endosporen. Die der Endosporen entbehrenden Arten sollen nach manchen Autoren zu zweien Arthrosporen bilden. Doch ist es nicht möglich diese von vielen Forschern ganz gelegneten "Arthrosporen" diagnostisch zu verwerthen.

Macé (1897, p. 438) recognized the family "*Bacteriaceae*". He says

Ce groupe renferme des Bactéries dont les éléments allongés suivant une direction, ont une longueur qui l'emporte sur la largeur. La forme typique est le batonnet, il est tantôt court et trapu, apparaît en coupe opt que presque comme un carré ou comme un court rectangle, tantôt sa longueur dépasse un certain nombre de fois sa largeur. La première de ces dimensions peut devenir très grande par rapport à l'autre, c'est la forme de filament. Les batonnets ou les filaments sont droits ou courbés. La courbure peut être très simple et ne représenter qu'une faible portion de circonférence, elle peut être compliquée, l'élément décrit une spire à tours plus ou moins nombreux, plus ou moins serrés. Quelles que soient

la forme ou la longueur de ces éléments, on ne leur distingue jamais d'extrémité antérieure et d'extrémité postérieure, de base ni de pointe, les mouvements, lorsqu'ils existent, paraissent toujours se faire également dans les deux sens, et quand des individus se fixent ou plutôt s'accrochent à un support, leur partie fixée ne diffère en rien de celle qui reste libre

The definition of Migula of *Bacteriaceae* has been followed among others by the following authors Chester (1901, p 117), Kendall (1902, p 484), A J Smith (1902, p 270), Klocker (1903, p 332), Migula (1904, p 145), Erwin F Smith (1905, p 160), Ellis (1909, p 3), Frost (1911, p 57), Schneider (1912, p 23)

Winslow *et al* (Committee Soc Am Bact, 1917, p 560) recognized this as the sixth family of the order *Eubacteriales* with the following diagnosis "Rod-shaped cells without endospores Gram-negative Flagella when present peritrichic Metabolism complex, amino-acids being utilized, and generally carbohydrates"

Buchanan (1918, p 27) recognized this as the second family of the *Eubacteriales*, with the following description

Cells rod-shaped, straight, or at least not spherical or spiral Never containing sulphur granules, nor with bacterio-purpurin, many species pigmented May or may not be motile by polar or diffuse flagella Cells may be single or in chains Endospores produced in some genera, not in others A pseudoplasmodium never developed Growth energy not secured by the oxidation of ammonia or nitrites

In 1920 Winslow *et al* (p 208) describe *Bacteriaceae* as the fifth family of the *Eubacteriales* including seven tribes

Bergey *et al* (1923, p 84) recognizes this as the fourth family of the order *Eubacteriales* with the description "Rod-shaped cells without endospores Motile or non-motile Metabolism complex, amino-acids being utilized, and generally carbohydrates Usually Gram-negative" Nine tribes are included

It is evident that while the earlier writers used this name as a designation of the entire group of bacteria, the later writers have uniformly regarded it as a family within the group

Bacteriaceen. An aberrant form of spelling of the family *Bacteriaceae* (*q.v.*) used by Zopf (1884, p 45, and 1885, p 51)

Bactériacées. An aberrant form of spelling of the family *Bacteriaceae* (*q.v.*) used, among others, by Cornil and Babes (1890, p 151)

Bacteriacei An aberrant form of spelling of the family *Bacteriaceae* (*q.v.*) used by Schroter (1885, p 155)

Bacteriales. An ordinal name suggested by Clements (1909, p. 8) His description is as follows "Globose, rod-like or filamentous, single or in colonies, sometimes grouped into a loose mass (zoogloea), but never forming pseudoplasmodia or sporangium-like masses" He recognized five families, *Beggiatoaceae*, *Chlamydobacteriaceae*, *Spirillaceae*, *Bacteriaceae* and *Coccaceae*

Bacteridia The Anglicized equivalent of Davaine's casual name *bactériodie* for the anthrax organism

Bactériodie. A casual name given by Davaine (1863, p. 195) for the anthrax bacillus Used for *Bacteridium q v*

Bacteridiaceae The fifth tribe of the family *Bacillaceae* as described by Castellani and Chalmers (1919, p. 932) One genus only, *Bacteridium* Schroeter, is included The tribal designation is,

Bacillaceae growing well on ordinary laboratory media, without endospores, and either fluorescent or chromogenic

Type Genus *Bacteridium* Schroeter, 1872

Bacteridium A genus proposed by Davaine (1868, p. 21) to include organisms with filiform bodies, more or less distinctly articulated in consequence of imperfect division, always non-motile The name was created to provide for the anthrax bacillus He states,

III Genre Bactériodie *Bacteridium* (Davaine) Corps filiforme, droit ou en flechi, plus ou moins distinctement articulé, par suite d'une division spontanée imparfaite, toujours immobile

Apparently in this article at least he did not give a species name He writes "Bactériodie Charbonneuse (Davaine)" for the anthrax bacillus The latin generic name is definite enough, as is also the species considered The casual designation *bactériodies* was first proposed in 1863 (p. 195), in the following statement

On voit, d'après les considérations qui précèdent, que le nom de bactérie, par lequel j'ai désigné primitivement ces corps, ne peut leur convenir Lorsque l'étude des êtres microscopiques qui jouent un grand rôle dans la fermentation, dans la putrefaction, etc., est à peine commencée, il serait prématuré de vouloir classer d'une manière définitive les corpuscules du sang de rate qui ont avec ces êtres une analogie évidente Je me bornerai donc, pour désigner ces corpuscules, à modifier légèrement le nom que je leur ai primitivement donné, et je les appellerai désormais des *bactériodies*

It is apparent that we have the anomalous situation of a suitable generic name, *Bacteridium*, being published, together with an adequate

description of a species which should be considered the type, but without a scientific name being given to the species. There may be some question under such circumstances of the validity of the generic name, though there would seem to be no statement in the International Rules of Botanical Nomenclature which would definitely make it invalid. It should be noted that while this author repeatedly in later writings used the term *bactérienne*, the generic name *Bacteridium* was not employed with a specific name to designate the anthrax bacillus (1864, p 393), (1865, p 334). It is of interest to note that Davaine earlier (1863, p 222) used the designation *bacterium du sang de rate*.

Schroeter (1872, p 110) used this designation to include certain non-motile pigment-producing bacteria, among them *Bacteridium prodigiosum*. This use of the genus name was discussed and disapproved by Cohn (1872, p 182) because the pigment producers have no marked character in common with the anthrax bacillus other than lack of motility.

Cohn says,

Wegen des Mangels der spontanen Bewegung hat Schroeter in dem voranstehenden Aufsatz die Kugelbakterien, welche Pigmente erzeugen, mit demselben Namen belegt, welchen Davaine für die unbeweglichen Stäbchen des Milzbrandblutes eingeführt hat (*Bacteridium*). Die Milzbrandbakteridien unterscheiden sich jedoch durchaus von den Pigmentbakterien, da sie stäbchen- oder langfadenförmig sind, sie können daher mit den Kugelbakterien nicht in einer Gattung zusammengestellt werden, da der Mangel der Bewegung der einzige beiden gemeinschaftliche Charakter ist.

The name was abandoned by Schroeter (1886). Macé (1897, p 485) makes the following comments:

Davaine avait cru devoir distinguer autrefois un genre *Bacteridium*, caractérisé par l'immobilité des éléments à tout les stades de leur existence. Il l'avait établi pour la Bactérie du charbon et quelques autres observées dans l'intestin et les infusions. Mais l'absence ou la présence de mouvements n'offre jamais de constance suffisante pour en faire un caractère spécifique. Un grand nombre d'espèces, très mobiles à une certaine phase de leur existence, deviennent complètement immobiles à la période suivante ou seulement quand les conditions de vie, tout en restant bonnes, viennent à changer, beaucoup deviennent inertes, par exemple au moment de la sporulation. Les Bactéries mobiles ne diffèrent du reste des immobiles par aucun caractère de valeur.

Erwin F. Smith (1905, p 158) claims that this genus name is invalid because of the previous existence of the genus *Bactridium* Kunze 1817. Whether or not this use invalidates *Bacteridium* Davaine depends upon

the interpretation of Article 57 and Recommendation XXXI of the botanical code. The former states that when the difference between two names, especially two generic names, lies in the termination, these names are to be regarded as distinct even though differing by one letter only. The latter reads

Many names differ by a single letter without risk of confusion. In cases where a close approach to identity is a source of error (ex *Astrotenma* and *Asterosemma* in one and the same family, *Asclepiadaceae*) only one, the older, of the names should be kept in accordance with Article 51, 4°

This last Article reads

Every one should refuse to admit a name in the following cases

4. When the group it designates embraces elements altogether incoherent, or when it becomes a permanent source of confusion or error

Smith suggested the name *Aplanobacter* in its place

Vuillemin (1913, p. 519) states that lack of motility does not justify generic recognition

Buchanan (1918, p. 33) recognized *Bacteridium* as a subgenus of *Bacillus* to include the non-motile species, with *Bacillus* (*Bacteridium*, *anthracis* Cohn as the type

Castellani and Chalmers (1919, p. 933) revive the *Bacteridium* of Schroeter with *Bacteridium prodigiosum* as the type. Their definition (p. 958) is "*Bacteridiae* which are chromogenic." It constitutes the type genus of the tribe *Bacteridiae* *q v* with the definition "*Bacillineae* growing well on ordinary laboratory media, without endospores and either fluorescent or chromogenic."

It would seem that the use of the genus *Bacteridium* in bacteriology and of *Bactridium* in mycology is valid if organisms of which the anthrax bacillus is the type are to be grouped into a separate genus. The previous use of *Bactridium* should no more invalidate *Bacteridium* than *Micrococca* Benth. a valid genus of the *Euphorbiaceae* created in 1849 should invalidate *Micrococcus* Cohn (1872).

Bacteriaceae. A tribe of bacteria created by Trevisan (1879, p. 136) with the following description

Bacteriaceae unicellulares. Individua globulosa vel ellipsoidea vel cylindrica, e cellula unica conflata. Sporae intra cellulam matricalem solitariae. Multiplicatio divisione cellularum vegetativa in unam vel duos vel omnes directiones.

Two subtribes were included, namely, *Eubacterieae* and *Sarcineae*

Buchanan (1918, p 39) recognized this tribe as the second of the family *Bacteriaceae*, with four subtribes *Fusiforminae*, *Hemophilinae*, *Rhizobinae* and *Bacterinae*. The description is "Cells rod-shaped, never spiral nor strictly filamentous, single or in chains, motile or non-motile, never producing endospores, either Gram-positive or -negative."

Bergey *et al* (1923, p 194) used this designation (with the spelling *Bactereae*) for the sixth tribe of the family *Bacteriaceae* with the following description "Gram negative rods generally growing well on artificial media. Generally attack carbohydrates forming acid and often gas composed of CO_2 and H_2 . When motile, the flagella are peritrichous." Six genera are included.

Bacterien. A casual name used by Cohn (1872, p 132) and subsequent German writers for the bacteria.

Bactéries. A casual name used by Davaine (1863) and subsequent French authors for the group bacteria.

Bacterinae. A subtribe of the tribe *Bacterieae* named by Buchanan (1918, p 47) with the following diagnosis:

Cells not fusiform, rod shaped, not hemoglobinophilic, aerobic, facultative, or microaerophilic, not securing growth energy exclusively by the oxidation of carbonaceous compounds. Spores never formed. Motile or non-motile.

It included the following genera: *Pseudomonas*, *Serratia*, *Chromobacterium*, *Pasteurella*, *Bacterium*, *Proteus*, *Pfeifferella*, *Lactobacillus*, *Erysipelothrix*, and *Corynebacterium*.

Bacterina. A group created by Perty (1852, p 179) to include the genera *Bacterium*, *Vibrio* and *Metallacter*. The name has never come into common use.

Bacteriopsis. A generic name proposed by Trevisan (1885, p 103) for a group of bacteria. The organisms were considered as showing three stages of development: 1. Bacilli, 2. Filaments, 3. Cocci. The bacillus form was regarded as most typical. The cells were short cylindrical or ellipsoid, straight, not articulate, colorless or colored, showing both macrobacilli and microbacilli, cytoplasm equally diffused in cells. Spores none or unknown.

The species included in the genus are as follows:

Bacteriopsis Rasmusseni = *Leptothrix* I Rasmussen

Bacteriopsis aceti = *Mycoderma aceti* Pasteur

Bacteriopsis Pasteuriana = *Bact Pasteurianum* E. Chr. Hansen.

Bacteriopsis merismopedioides = *Bact merismopedioides* Zopf

Bacteriopsis ureae = *Micrococcus ureae* Zopf
Bacteriopsis vanthana = *Bact. vanthnum* Zopf
Bacteriopsis synzantha = *Vibrio synzanthus* Ehrenb
Bacteriopsis ovata = *Panhistophyton ovatum* Lebert

The genus has not been used by subsequent authors, and is not even recognized by DeToni and Trevisan (1889).

It is rejected by Erwin F. Smith (1905, p. 174).

Probably the first named species, *Bacteriopsis Rasmusseni*, should be regarded as the type. It was later (1889, p. 930) placed by De Toni and Trevisan in the genus *Rasmussenia* (the fourth species, *R. anceps*). If no older generic name for the mouth leptothrix forms can be found, this would probably be a valid designation for them. It may be regarded as a synonym of *Leptotrichia*.

Bacterium. A generic name proposed by Ehrenberg (1828, p. 8). His diagnosis is as follows:

Bacterium, Novum Genus, Familia Vibrionorum. Character Generis: Corpus polygastricum? anenterum? nudum, oblongum, fusiforme aut filiforme, rectum, monomorphum (contractione nunquam dilatatum), parum flexile (nec aperte undatum), transverse in multas partes sponte dividuum.

The only species described was *Bacterium triloculare*. The description was as follows:

B. triloculare nov. spec., distincte triloculare s. triarticulatum, subfusiformum, hyalinum.

Animalculum 1-300 lineae longum, corpore tereti. Articuli s. septa interna divisionem instantem multiplicem transversam indicare videntur. Mobile sed pigrum animalculum.

In Oasi Jovis Hammonis Siwae observatum, praeterea nullibi.

Bacterii Generis physiologia huiusque obscura. Cibo colorato Ventriculos replere hae formae respuunt ideoque ad Polygastrica non nisi dubitanter et interim collocantur.

While it is not improbable that Ehrenberg actually described one of the organisms now included with the bacteria, it is evident that there is nothing in the description which would make probable a modern identification of the species.

Two years later Ehrenberg (1830, p. 38) included some eleven species in his *Bacterium*. It is to be noted that some were not new species. Several of Mueller's species of infusoria described in 1783 were included, among them being *Bacterium (Monas) termo* (Mueller) Ehrenberg. In the previous (1828) paper, Ehrenberg had recognized *Monas termo*.

Mueller as distinct, not including it in the genus *Bacterium*. Inasmuch as *Bacterium termo* has been much discussed in the literature, the description given by Mueller (1786, p. 1) should be noted

Monas termo Animalculum omnium, quae microscopium simplex offert, minimum simplicissimum, punctulum, gelatinosae, substantiae pisum microscopium compositum eludere videtur, dum ne quidem sub hoc distinctius appareat

Sphaericum, an orbiculare? haud video

Guttula aquae, in qua maceratio facta est, his corpusculis adeo saepe repletur, et ne minimum vacuum distingui liceat, ipsamque aquae substantiam in aliam minus hyalinam, globularem ex punctis consetitissimis, omnem calculum superantibus mutam crederes

In hac massa motus, qualem radii solares in aqua micantes effingere solent, oculis exhibetur, dum animalcula, examinis opum instar, vehementer commoventur

In infusione vegetabilium et animalium Huius guttula jam intra viginti quatuor horas conspicitur quasi massa globularis, aut nullus in ea motus nec odor percipitur, brevi vero motus seu fermentatio cum foetore interabili insequitur, at non in omni

Fig. 1 Guttulam aquae fluvialis *Monade Termone* scetentam valde auctam exhibet

Ehrenberg's description of *Monas termo* Mueller as given in 1828 is also pertinent

M. Termo (Mueller) 1/1664 ad 1/500 lineae crassa, exacte blobosa, hyalina

Specimina, seu potius Myriades, quas Siwae in Oasi libyca vidi et illae, quas infuso Pipere nigro in Arabia accepi, magnitudine 1/1664 lineae fere aequabant Quae e sinaitica valle Wadi Esle depromsi mensuram 1/1660 lineae offerebant Berolini et in Sibiria interdum maiora individua, ab eadem specie vix segreganda, vidi, maxima 1/500 lineae crassa, quae, si divisione spontanea funduntur, in partes 1/1000 lineae crassas, reliquis similes, abeunt

Monadum speciales characteres, ob microscopiorum, quae vocamus, sufficienter eas augendi vim deficientem, praeter corporis formam et mensuram, non habent quo nitantur et firmitate physiologia saepe carent Motus, coloris, mensurae et propagationis ratio cum forma nonnullas quidem species bene distinguere sunt, in aliis characterum gravitas desideratur, sed formarum magnam copiam in aquis vagari unusquisque, qui his animalculis studium impendit, facile sibi persuadet Quin ex Monadibus, Vibrionibus, Bacteriis et Bodonibus us, quae minutie excellunt et cibos colaratos respuunt, novae animalium classis vestigia posteris pre-menda entescere saepe mihi visa sunt

The generic description given for *Bacterium* by Ehrenberg (1830, p. 38) is "*Bacterium* Nov Gen Haec genera, Oscillatorius valde affinia ore nutriri nundum vidi" Eleven species are listed

Two years later (Ehrenberg 1832) differentiated six species of the genus *Bacterium* Two of these, *Bact. articulatum* and *Bact. triloculare*

are definitely included. Four other species *Bact. Enchelys*, *Bact. punctum*, *Bact. tremulans* and *Bact. termo* are questioned.

Six years later in his classic work on the *Infusoria* Ehrenberg (1836, p. 75) characterized the genus *Bacterium* as follows: "Animal e familia Vibrioniorum, divisione spontanea in catenam filiformem rigidulam abiens."

One species only, *Bacterium triloculare* is included. In his drawing of *Bacterium triloculare* a polar flagellum is shown.

A careful perusal of the description certainly reveals nothing that would enable one even with a moderate degree of certainty to identify Ehrenberg's species. Certainly most of them were based on impure or mixed cultures. From different localities and conditions the same organism was frequently described though differing sufficiently in size and appearance to justify the conclusion that the forms were quite distinct.

Inasmuch as some effort has been made to tie up the genus *Bacterium* to *Bacterium termo* as a type species, the following points are worthy of note:

1. *Bacterium termo* (Mueller) Ehrenberg was not included as a member of the Genus *Bacterium* in the paper in which the genus was first proposed, in fact, as *Monas termo* Mueller, it was placed in another genus.

2. Mueller (1786, p. 1) states that *Monas termo* is spherical and Ehrenberg (1828) likewise describes it as globular ("exacte globosa").

3. The organism was found in water and in infusions of animal and vegetable matter.

It should be noted that Ehrenberg includes the genus *Bacterium* as one of five genera in his family *Vibrionia*, the other genera being *Vibrio*, *Spirochaeta*, *Spirillum* and *Spirodiscus*. *Bacterium* is differentiated from *Vibrio* in that it is non-flexuous, while the latter is flexuous.

Dujardin (1841, p. 212) practically adopted Ehrenberg's classification. He designated as *Bacterium* all those forms which are straight, more or less definitely jointed, non-flexile, and which were capable of only a slow wave-like motion. *Vibrio*, on the other hand, contained those organisms which are straight or bent, more or less definitely jointed, and which show sinuous motion like a snake. He placed three species in this genus, *Bact. termo*, *Bact. catenula*, and *Bact. punctum*. His description reads: "Corps filiforme, roide, devenant plus ou moins distinctement articulé par suite d'une division spontanée imparfaite. Mouvement vacillant non ondulatoire." Here again there is nothing which will enable the bacteriologist to identify any one of the species.

It is of interest to note that many subsequent writers ascribe the species *Bacterium termo* to Dujardin

Diesing (1850, p 14) gives the following generic description of *Bacterium*

Corpus nudum, lorica destitutum head mutabile, cylindricum subovatum aut subglobosum, partitione imperfecta uniseriali, multiplici, transversa in syntherium lineare, moniliforme, rigidulum, rectum, hyalinum accrescens Flagellum in articulo primo terminale simplex Ocellus nullus

Perty (1852, p 179) developed a classification of the infusoria, recognizing a family *Vibrionida* with two subfamilies *Spirillina* and *Bacterina* Under the latter the genera *Vibrio*, *Bacterium*, *Metallacter* and *Sporonema* are included *Vibrio* is defined as flexuous, *Bacterium* is non-flexuous The only species of *Bacterium* described is *Bact termo* ascribed to Dujardin It is recorded as occurring "in den aller-verschiedensten faulenden Aufgüssen und Sumpfwässern das ganze Jahr" He notes that it was recorded by Dujardin "im Eiter und andern pathologischen Flüssigkeiten" and even "nach Leeuwenhoek im Zahnschleim" The description is wholly unsatisfactory, and shows a tendency, which became more pronounced during the succeeding two decades, to call all organisms associated with putrefaction and decay *Bacterium termo*

Cohn (1854, p 123) for the first time definitely grouped the form now termed bacteria under the plant kingdom, removing them from the *Infusoria* He studied at length the *Bacterium termo* Dujardin, which he regarded as a synonym of *Vibrio lineola* Ehrenberg He concluded this form to represent merely the swarm cell state of an alga-like fungus He was unable to see the "cilia" He gave it the generic name *Zoogloea* with the following characters

Cellulae minimae, bacilliformes, hyalinae, gelatina hyalina in masses mucosae globosae, uvaeformes, mox membranaceae consociatae, dein singulae elapsae, per aquam vacillantes

The species became *Zoogloea termo* As synonyms he notes "*Palmella infusionum* Ehr, *Micraloa teres* V Flotow, *Bacterium termo* Dujardin and *Vibrio lineola* Ehrenberg

Several authors in the next two decades followed Cohn in the substitution of *Zoogloea* for *Bacterium* For example, Rabenhorst (1865, p 35) in his "*Flora Europaea Algarum*" includes this genus among the algae under the heading "*Formae achroae, hyalinae (mea sententia ad*

fungos referendae”). The generic description given is identical with that of Cohn. The single species *Zoogloea termo* (Muller) Cohn is briefly described as “Z thallo gelatinoso globoso, cellulis liberis mobilibus, rectis, 1/2000–1/700” aequantibus. Hab. in aquis stagnantibus, imprimis corruptis ubique”

A *Zoogloea ramigera* was later described by Itzigsohn (1867, p 414).

The next author of note defining *Bacterium* was Davaine (1864, p 629) who, like Cohn, separated the *Vibriones* from the protozoa, and described the genera *Bacterium*, *Vibrio* and *Spirillum*. His definition of *Bacterium* (given in 1868, p 68) is identical with that of Dujardin. It should be noted that this author emphasized the importance of motility in the generic diagnosis.

An interesting sidelight on the utilization at this time of the term *Bacterium termo* to include the greatest variety of microorganisms is to be found in the writings of Luders (1866, p 36) who concluded that this wide spread organism was a growth stage of a fungus. Hallier (1886) introduced and developed his theory of pleomorphism which reached its culmination in the work of Billroth (1874). This led to general distrust at this time in the existence of true bacterial species and indirectly contributed to the lack of regard for the rules of nomenclature during the next several decades.

Davaine (1868, p 21) used the name in the sense of Dujardin differentiating the non-motile forms such as the anthrax bacillus in the new genus *Bacteridium*. His description of *Bacterium* is as follows:

Genre Bacterie *Bacterium* (Ehrenberg, Dujardin) Corps filiforme, roide, devenant plus ou moins distinctement articule par suite d'une division spontanee imparfaite, mouvement vacillant non ondulatoire (Dujardin).

In summary, to the year 1872, the generic name *Bacterium* was not applied to any species which can now be recognized satisfactorily. The original *Bacterium triloculare* Ehrenberg was no longer recognized in the literature. *Bacterium termo* the organism found commonly in putrefying solutions was the only species.

Cohn (1872, p 167) rather radically emended the previous generic descriptions of *Bacterium* to include those rod-shaped organisms in which the cells are short, never filamentous. He made it the sole representative of his tribe *Microbacteria*. The cells are short cylindric or elliptical, never in chains or filaments, often forming firm zooglocal masses, alternating motile with non-motile stages. His description was largely based on his conception of *Bacterium termo*, which he regarded

as the principal agency of decay ("ohne *Bacterium termo* keine Faulniss") He described the reaction secured by inoculating the bacteria from putrefying beans into his ammonium tartrate solution and noted the development therein of a green fluorescence

Should this description of Cohn be used as fixing the type for *Bacterium* as *Bacterium termo*? The following objections may be raised

1 Unquestionably Cohn was dealing with impure or mixed cultures. The fact that when this mixture was inoculated into Cohn's nutrient solution a green fluorescence was produced by no means fixes this as a characteristic of *Bacterium termo* Bacteria capable of bringing about this change are widely distributed in nature In other words, in the heterogeneous mass of organisms present in infusions, known as *Bacterium termo*, there are included certain fluorescent bacteria capable of growing in Cohn's nutrient solution The International Botanical Code is explicit on this point

Article 51 Every one should refuse to admit a name in the following cases

4 When the group which it designates embraces elements altogether incoherent or when it becomes a permanent source of confusion or error

2 The species *Bacterium termo* was first described as *Monas termo* Mueller, it was tentatively transferred to the genus *Bacterium* by Ehrenberg, but later rejected by him, and first definitely included in the genus by Dujardin It was later made a type of the genus *Zoogloea* by Cohn Inasmuch as this organism was not at first included in the genus *Bacterium*, in fact was definitely excluded by the author of the name, it does not seem advisable to accept it as the type

Schroeter (1872, p 120) put in the genus *Bacterium* three species of bacteria which had previously been described in other genera, the *Vibrio synzanthus*¹ Ehrenberg (1840, p 202) he renamed *Bacterium synzanthum* (the spore bearing rod responsible for yellow milk), the *Vibrio syncyaneus* Ehrenberg² (1840, p 202) he termed *Bacterium syncyaneum* (the polar flagellate organism of blue milk) and named the *Bacterium aeruginosum*, the polar flagellate organism of blue green pus Both of these organisms were later (1886, p 157) transferred by Schroeter to the genus *Bacillus* The names used by Schroeter were accepted by Cohn (1872b, p 173)

Should one of Schroeter's species be accepted as the generic type? It probably would be inadvisable to designate thus either of the species

¹ The *Vibrio anthogenus*, Fuchs (1841, p 193)

² The *Vibrio cyanogenus*, Fuchs (1841, p 190)

transferred to this genus from other genera, for it is quite conceivable, that inasmuch as *Vibrio synchyaneus* Ehrenberg was probably the first species of *Vibrio* sufficiently described so that it can be certainly recognized and identified today, it might be designated as the type of *Vibrio*. In other words, if it is to be accepted as a type, it should be for *Vibrio* and not for *Bacterium*. The form from yellow milk quite certainly should be placed in the genus *Bacillus*. *Bacterium aeruginosum* is so closely related to *Bact. synchyaneum* that they probably should be placed in the same genus. It would certainly be unwise to choose the former as the type, as there might well be question as to whether it did not belong likewise in the genus *Vibrio*. It would seem that stability of nomenclature would not be conserved by the choice of any one of these as a type species.

If *Bacterium* is to be retained as a bacterial genus, it seems necessary to pick the type species from those described subsequently to 1872.

Lister (1873, p. 408) gave the name *Bacterium lactis* to an organism responsible for lactic acid fermentation. While his description may not be sufficiently accurate to enable us to determine with certainty whether he described the lactic streptococcus or one of the aerogenic lactic bacilli, (such as *Bacterium aerogenes*) now generally regarded as belonging to the colon-typhoid group, it is probable that the former should be the interpretation. *Bacterium lactis* became *Streptococcus lactis* and is not available as a type.

Lankester (1873, p. 13) described a "peach coloured *Bacterium*" from water under the name *Bacterium rubescens*. The description makes it evident that he was working with organisms belonging to the group of forms containing bacteriopurpurin, that is to the group of sulfur bacteria. The description further makes it evident that he was not dealing with a pure culture. He dilates upon its variability, the appearance of rods and spirals, and motile and non-motile cells etc. Certain of the forms were quite certainly previously described under other names. Cohn wrote Lankester (1874, p. 399) that his organism was the *Monas Okeni* Ehrenberg, the *Clathrocystis roseopersicina* of Cohn. While Lankester combated Cohn's statements, modern knowledge of the group would seem to justify them. Certainly the recognition of *Bacterium rubescens* as a type species would be unwise.

Warming (1875) described several species of *Bacterium* from the water on the coast of Denmark. These were *Bacterium griseum*, *Bact. litoreum*, and *Bact. fusiforme*. They were not grown in pure cultures and apparently are not identifiable with certainty.

Dallinger (1878, p 159) studied *Bacterium termo* microscopically, demonstrating motility to be due to a terminal flagellum It is defined in the sense of Cohn

In 1875 Cohn included the genus in the tribe *Glaeogenae* It was defined as having cylindrical cells, without phycochrome, very small, cells united in glairy families, amorphous when in repose, or entirely free, never disposed in filaments In this revision he accredited the genus to Dujardin rather than to Ehrenberg Magnin (1878, p 80) follows Cohn exactly The diagnosis reads

Bacterium, Duj emend Cells cylindrical or elliptical, free or united in pairs during their division, rarely in fours, never in chains, sometimes in *zoogloae*, having spontaneous movements, oscillatory and very active, especially in media rich in alimentary material and in presence of oxygen

The species are divided into three groups

1 The bacteria of putrefaction (*Bact termo* Ehr and *Bact lineola* Cohn)

2 The bacteria of lactic and acetic fermentations, and

3 The chromogenic bacteria (including *Bact xanthinum* Schroter, *Bact syncyaneum* Schroter, *Bact aeruginosum* Schroter, and *Bact brunneum* Schroter)

Of the species, the chromogenic forms can for the most part still be readily recognized

Lanzi (1876, p 256) designated the organism responsible for acetic fermentation *Bacterium aceti*

Trevisan (1879, p 136) gives the following synonymy and description

Bacterium Ehrenb (1830 non 1828) emend Cohn (Beitr I Heft 2, p 177), (*Chromatium* Perty 1852, *Zoogloea* Cohn 1853, *Rhabdomonas*, Cohn 1875) *Cel-lululae breviter cylindricae, vel ellipsoideae, vel fusiformes Multiplicatio divisione cellularum vegetativa semper ad eandem directionem longitudinalem*

The species recognized are *Bact aceti* Lanzi 1876, *Bact termo* Ehrenb, *Bact lineola* Cohn, *Bact tremulans* Trevisan, *Bact catenula* Dujard, *Bact. punctum* Ehrenb, *Bact synxanthinum* Schroeter, *Bact syncyaneum* Schroeter and *Bact aeruginosum* Cohn

The generic description of *Bacterium* by Luerssen (1879, p 20) does not assist in fixing a type species

Van Tieghem (1879) described a *Bacterium lucens* from the surface of liquids

Sternberg (1880, p 80) in his translation of Magnin's (1878) text on the bacteria gives the following characterization of the genus *Bacterium*

Cells cylindrical or elliptical, free or united in pairs during their division, rarely in fours, never in chains (*Leptothrix* or *Torula*), sometimes in zoogloea (differing from the zoogloea of spherical bacteria by a more abundant and firmer intercellular substance), having spontaneous movements, oscillatory and very active, especially in media rich in alimentary material and in presence of oxygen

He includes descriptions only of species which have already been noted. They are separated into three groups, the bacteria of putrefaction, those causing lactic and acetic fermentation, and the chromogenic forms

Thin (1880) described a *Bacterium foetidum* from malodorous perspiration. The culture methods used did not insure pure cultures.

Winter (1880, p. 51) characterizes *Bacterium* (which he ascribes to Cohn) as follows

Zellen kurz cylindrisch, lang elliptisch oder spindelförmig, durch Quertheilung sich vermehrend, spontan beweglich. Die Tochterzellen trennen sich entweder bald nach der Theilung von einander oder bleiben zu 2 bis mehr kettenartig vereinigt. Auch Zoogloea-Bildung ist häufig. Sporenbildung ähnlich wie bei *Bacillus*.

He lists and describes eight species, none new. It will be noted that he includes for the first time the possibility of spore formation as a character of the genus.

Van Tieghem (1881) described a green organism from rain water which had collected in the pileus of a *Polyporus*, giving it the name *Bacterium viride*.

Thin (1881) also described a *Bacterium decalvans* from *alopecia areata*.

Zopf (1883, p. 48) materially modified the definition of the genus following the conception of pleomorphism, to include those organisms showing four growth forms, cocci, short rods, long rods and filaments, the latter never with differentiation of base and tip. The following year (1884, p. 45) he used the same definition. He included in the genus some seventeen species, most of which are still recognized. Organisms previously described by other authors as in the genus *Bacillus* are included, such as *Bacterium tuberculosis*, *Bact. typhosum*, *Bact. subtile* and *Bact. anthracis*. His *Bact. ianthinum* was the *Bacteridium violaceum* of Schroeter and the *Chromobacterium violaceum* of Bergonzini. The only organism apparently newly named and placed in this genus was *Bacterium acidi-lactici*. The description is scarcely adequate to identify this organism as the one now generally recognized under this name. The specific name *acidi-lactici* has been used in so many senses

by subsequent authors as to make identification difficult. However, *Bact. acidilactici* may be regarded as a possible type species.

Engelmann (1882, p. 94) described an organism from water as *Bacterium photometricum*. It was insufficiently differentiated.

Kurth (1883, p. 97) isolated and described from the alimentary tract of the domestic fowl an organism which he named *Bacterium Zopfii*. This organism is adequately known at the present time.

Lanzi (1883, abs. 1884, p. 786) described an organism as *Bacterium moribilli* but not adequately.

Luerssen (1879, p. 20) used the generic diagnosis of Cohn as did also Grove (1884, p. 22). This latter author lists five colorless species (including *Bact. termo*) and four chromogenic species (including *Bact. synanthum* Schroter, *Bact. syncyanum* Schroter, *Bact. aeruginosum* Schroter and *Bact. violaceum* (Bergonzini) Grove). The chromogenic species may all be recognized.

Van Tieghem (1884, p. 1110) defined *Bacterium* as containing rod-shaped cells, separating as soon as formed, never in filaments, never spiral. He regarded the name more as a form genus than a natural genus.

Escheich (1885) described two organisms, *Bacterium coli commune* and *Bacterium lactis aerogenes*, from infant stools. The description was greatly expanded the next year (1886, pp. 57 and 63). These organisms have been consistently known by these names to the present time.

In 1885 Trelease described several new species of pigmented bacteria under the names *Bacterium candidum*, *Bact. luteum*, *Bact. aurantiacum*, *Bact. chlorinum* and *Bact. incarnatum*. Zukal (1885) studied the morphology of a *Bacterium tortuosum* from water.

Zopf in 1885 (p. 61) completely revised his classification. *Bacterium* is defined as made up of those species which exist as cocci and rods, or the latter only, united into filaments, spore formation lacking or unknown. It will be noted that here the emphasis is changed to absence of spores. Hueppe (1885) used practically the same diagnosis. Maggi (1886, p. 84) listed five species of this genus.

Flügge (1886) outlined a classification of bacteria which has been extensively followed by medical bacteriologists. His genus *Bacterium* is defined to include cylindric forms, short, single or loosely united or in amorphous slimy masses.

Schroter (1886, p. 155) states that the genus *Bacterium* is not well characterized and possibly should be united with the genus *Bacillus* from which it is separated with difficulty. He accepts absence of spores.

as a generic character, and states that the vegetative cells are short rods, little longer than broad, in the rounded form either solitary or embedded in slime (*Zoogloea*), from which a motile form emerges. Three species only are listed *Bact. termo* Ehr, *Bact. septichaemiae* and *Bact. cholerae gallinarum*. The chromogenic species previously listed here by Cohn and Schroter are transferred to the genus *Bacillus*. Hansgurg (1888, p. 265) listed *Bacterium* as the second genus of the subfamily *Microbacteria*, but with no diagnosis. De Toni and Trevisan (1889, p. 1020) give the following generic diagnosis

Bacterium Ehrenb. (1830) emend. Trev. Bacterii et Bacilli species Auctorum. Baculi vulgo brevis, interdum brevissimum, ut plurimum ellipsoidei raro cylindracei, apicibus obtusis, recti. Sporae (arthrospora) globosae, e coccorum metamorphosi ortae.

The genus is included in the subfamily *Baculogenae* and the tribe *Bacilleae*. In all, thirty-four species are described. Of these *Bact. acetii* (Kuetz) Lanzi, *Bact. Pasteurianum* (Hanson) Zopf, *Bact. allantoides* Trev. and *Bact. Balbianii* Billet are noted as particularly typical. The genus is divided into two primary groups, *Bacteria achroa* (22 species) and *Bacteria chromogena* (12 species). The first group is again sub-divided into *Species Zymogena* (8 species, mostly acetic acid bacteria), *Species saprophyticae* (13 species) and *Species anthrophobiae* (1 species). The second group is divided into *Species granula sulphuris nulla secernentes* (7 species) and *Species granula sulphuris secernentes* (5 species). Baumgarten (1890) abandoned the genus, including all rod forms in *Bacillus*. Billet (1890, p. 23) says "Formes rectilignes *Bacterium* element 1 à 5 fois plus long que large". Sternberg (1892, p. 18) states "This genus, established by Dujardin, is now generally abandoned, the species formerly included in it being transferred to the genus *Bacillus*". Ludwig (1892) included in *Bacterium* rod-shaped elliptical organisms, not in long filaments, not embedded in slime and differentiated from *Bacillus* by lack of endogenous spores. Freudenreich (1894) defines a *Bacterium* as a rod without endospores or with arthrospores.

Migula (1894, p. 236) radically emended the genus to include only the *non-motile* rods, i.e., those forms with shorter or longer cylindrical cells, sometimes forming filaments, without flagella, endospore formation observed in many species, in others apparently completely lacking. He states "*Bacterium* Cohn Zellen ohne Bewegungsorgane oft mit Endosporenbildung (Z. B. *Bacterium Anthracis*)". He notes that about 200 species are known.

Clements (1894, p 27) defined the genus *Bacterium* as follows "Cells very short, ellipsoid, rarely cylindrical, motile or non-motile, often imbedded in a gelatinous matrix and forming zoogloea "

Two species only are described, *Bacterium termo* and *Bact acet*

The genus name *Bacterium* is not used by Fischer (1895) at all Lehmann and Neumann (1896, p 181) emphasized lack of endospores as the most important character They say

Zellen mindestens $1\frac{1}{2}$ mal meist aber 2-6 mal als lang als breit, gerade oder in einer Ebene gekrümmt, zuweilen lange echte oder Scheinfaden bildend, mit oder ohne Geisseln Stets ohne Endospores, für einzelne Arten sind Arthrosporen beschrieben.

Macé (1897) retained the older conception of the genus. He states:

La longueur relative était le seul caractère qui le distinguerait du genre *Bacterium* Nous savons que ce caractère ne peut être considéré comme absolu, mais qui varie au contraire dans des limites très larges, pour des conditions de vie qui peuvent être considérées comme normales

Migula in 1897 again discusses the importance of organs of motility in a system of classification, and once more defined the genus as consisting of non-motile rods Chester in his Manual (1901, p 117) adopts Migula's definition of *Bacterium*, as does also Allan J. Smith (1902, p 270) and Kendall (1902, p 484)

Klocker (1903, p 332) evades the issue by using the generic name given by the author of the specific name of the species under discussion He states "For distinguishing purposes the spore formation and the flagella have been suggested, but these characteristics have proved to be insufficient "

Matzuschita (1902, p IV) does not use the genus name *Bacterium* Schmidt and Weis (1902, p 294) define the genus as follows

Die Zellen sind cylindrisch, stabförmig, frei oder zu Faden verbunden Geisseln fehlen stets, und diese Arten sind daher zu keiner Zeit ihres Lebens beweglich Endosporenbildung bei vielen bekannt Zur Gattung *Bacterium* gehört eine sehr grosse Anzahl (über 300) Arten, von denen viele pathogen sind

Erwin F Smith (1905, p 160) discusses at considerable length the history of the generic name *Bacterium*. He comes to the conclusion that *Bact termo* of Cohn can be identified with sufficient accuracy as a green fluorescent organism with polar flagella, developing in Cohn's nutrient solution when inoculated with beans He, therefore, formulates the following diagnosis

Bacterium Cohn Type The one-flagellate, green fluorescent schizomycete capable of growing in Cohn's nutrient solution To these should be added all the morphologically similar non-fluorescent and yellow species Synonym: *Pseudomonas* Migula

He creates the new genus *Aplanobacter* to replace the genus *Bacterium* of Migula.

Flügge (1908) includes in the genus all rods which develop no spores He divides the genus into nine groups which are worthy of careful consideration as generic groups.

Ellis (1909, p 6) follows Migula

Orla-Jensen (1909, p 344) places the genus *Bacterium* in his family *Acidobacteriaceae*, the type species being *Bact coli*

Conn (1909, p 12) defines *Bacterium* as a non-motile rod, contrasting *Bacillus* which is defined as a motile rod Heim (1911, p 25) uses *Bacterium* as a subhead under *Bacillus* to include non-spore-bearing forms

Frost (1911, p 57) follows Migula exactly Schneider (1912, p 23) includes "the old genus *Bacterium* in the genus *Bacillus*" Meyer (1912, p 2) uses *Bacterium* as a section of the genus *Bacillus* including those forms whose flagella are unknown

Lohnis (1913, p 45) differentiates *Bacillus* and *Bacterium* on the basis of endospore production

Vuillemin (1913, p 524) advocates the designation of *Bacterium* as a Genus conservandum with a prototype *Bacterium termo* Ehr and specific type *Bacterium pyocyaneum* This, it will be remembered, is practically the recommendation of E F Smith Vuillemin states that Ehrenberg differentiated *Bacterium* from other types of *Schizomycetes* by its oscillating movement Today, he states, we know this type of motion to be due to two morphological characters, short rods and polar flagella *Bact pyocyaneum* is a short rod with polar flagella, hence the designation of this species as the type

Winslow *et al* (Committee Society of American Bacteriologists (1917, p 561)) adopted the view of Orla-Jensen and gave the following generic definition

Motile or non-motile rods, staining evenly Easily cultivable Animal pathogens or saprophytes Often chromogenic Many forms decompose carbohydrates

The type species is *Bacterium coli* Escherich

Buchanan (1918, p 53) has likewise followed Orla-Jensen with the definition

Plump rods, without spores, Gram-negative, motile by means of peritrichous flagella, or non-motile, liquefying gelatin very slowly or not at all. Usually showing marked power to ferment carbohydrates, frequently with gas production.

The type species is *Bacterium coli* Escherich.

The genus is a large one, including many species. It is generally divided into subgroups based primarily upon the fermentative reactions. It may be convenient to recognize these as subgenera separated from each other by the characteristics noted in the following key.

Key to the subgenera of Bacterium

- A. Organisms which show a maximum of fermentative power, including fermentation of lactose, rarely pathogenic, some forms slowly liquefy gelatin.
Subgenus 1 *Aerobacter* (or *Eu-Bacterium*)
- B. Organisms not showing maximum fermentative power, never producing gas in lactose, frequently pathogenic, never liquefying gelatin.
 - 1. Producing acid and gas from glucose, sometimes other sugars, but not from lactose. Subgenus 2 *Salmonella*
 - 2. Producing gas from none of the carbohydrates, acid sometimes formed. Subgenus 3 *Eberthella*

Breed, Conn and Baker (1918, p. 445) suggest that the name be dropped, or at most regarded as a general designation for insufficiently described organisms.

Winslow *et al* (1920, p. 211) in the final report of the Committee use the following definition:

Gram-negative, evenly staining rods. Often motile, with peritrichic flagella. Easily cultivable, forming grape-vine leaf or convex whitish surface colonies. Liquefy gelatin rarely. All forms except *B. alcaligenes* and the *B. abortus* group attack the hexoses and most species ferment a large series of carbohydrates. Acid formed by all, gas (CO_2 and H_2) only by one series. Typically intestinal parasites of man and the higher animals although several species may occur on plants and one (*B. aerogenes*) is widely distributed in nature. Many species pathogenic.

Type species, *B. coli* Escherich 1885, p. 518.

Castellani and Chalmers (1919) have broken up the genus *Bacterium* into a large number of genera, apparently dropping this generic name entirely. In this they have been followed by Bergey *et al* (1923).

It is evident from the preceding analysis of the important literature bearing on the subject that the various diagnoses of the genus *Bacterium* with their principal advocates may be grouped as follows:

1. **Pre-Cohnian conceptions.** Relatively rigid cells or chains of cells, not flexible, motile, oscillating. First species described *Bacterium triloculare* Ehr. No type species designated. No species described at pres-

ent recognizable with certainty Ehrenberg (1828, 1830, 1836), Dujardin (1841), Perty (1852), Cohn (1854), Davaine (1864), Luders (1866), Davaine (1868)

2 Cohnian conception. Short cylindric or elliptic cells, never in chains or filaments. Alternating motility and non-motility. Shape and grouping the primary characteristics. No definite designation of a type species, although in the earlier writings *Bacterium termo* is frequently mentioned. Cohn (1872, 1875), Lister (1873), Lankester (1873), Dallinger and Drysdale (1874), Magnin (1878), Winter (1879), Luerssen (1879), Trevisan (1879), Grove (1884), Van Tieghem (1884), Escherich (1885), Flugge (1886)

3 Zopfian conception. Rod shaped organisms which produce no endospores Motility and grouping of cells regarded as secondary or non-essential characters No type species designated Zopf (1885), Hueppe (1895), Schroeter (1886), De Toni and Trevisan (1889), Ludwig (1892), Freudenreich (1894), Lehmann and Neumann (1896), Flugge (1908), Löhnis (1913)

4 Not recognized as a generic name. Forms usually included under *Bacillus* or other genera Baumgarten (1890), Sternberg (1892), Fischer (1897), Matzuschita (1902), Schneider (1912), Breed Conn and Baker (1918), Castellani and Chalmers (1919), Bergey *et al* (1923)

5 Migula's conception Non-motile, rod-shaped bacteria No type designated definitely, unless it be *Bacterium anthracis* (Davaine) Migula. Migula (1895, 1897, 1904), Chester (1901), Smith (1902), Kendall (1902), Ellis (1909), Heim (1911)

6 Smith-Vuillemin conception Rod-shaped organisms with polar flagella, type same one of the fluorescent bacteria, such as *Bacterium aeruginosum* (Schroeter) Cohn (*Bacterium pyocyaneum* (Gessard) Vuillemin) E F Smith (1905), Vuillemin (1913)

7 Conception of Orla-Jensen The genus to be defined to include *Bacterium coli* Escherich as the type Practically the colon-typhoid group Jensen (1909), Buchanan (1918), Committee (1918-1920)

Of the species of *Bacterium* named before Cohn's description in 1872, there does not seem to be a single one that can be identified with certainty Certain species of other genera can be recognized, but no organism designated *Bacterium* The diagnosis of the genus given by Cohn was soon found to include forms that were diverse, and proved impracticable However, certain of the species described and recognized by him can be identified

The generic diagnosis used by Zopf and by Hueppe, and more commonly used than any other includes in the genus the rod-shaped organisms which do not produce endospores. This was a natural emendation of Cohn's definition, as the *Bacilli* described by him were endosporous.

The definition of Migula excludes from the genus all motile rods. Certainly most of the organisms previously termed *Bacterium* by Cohn and his predecessors were motile forms. Furthermore, motility is one of the most variable of bacterial characteristics. This definition associates in the same genus forms so distantly related as the tubercle and the anthrax bacilli, and places in separate genera closely related organisms of the intestinal group such as the colon bacillus and the organisms of gassy milk.

It might seem best to select some organism early named *Bacterium* as the type and so word the generic diagnosis as to include this form and the related species. But it should be described by the inclusion of more than a single character, such as motility. E. F. Smith has attempted to identify the *Bacterium termo* of Cohn, evidently intended by this author to constitute the type of the genus. Smith states

Much discredit has been thrown on *Bacterium termo* in modern times, and it has been left out of many classifications. However, if one examines into the matter, there is no reasonable doubt as to what Cohn had in mind. His *Bacterium termo* was a small schizomycetous organism capable of growing freely in Cohn's nutrient solution, containing acid potassium phosphate and ammonium tartrate. It produced therein short rods (single, in pairs or fours, joined end to end) and roundish lobed white zoogloae, together with a greenish fluorescence. This is Cohn's statement and De Bary's. It did not appear in boiled fluids, i. e. was destitute of endospores (Cohn), and the motile rods were killed by short exposure to 58°C (Schroter). In other words, it was a non-sporiferous green-fluorescent organism possessed of a single polar flagellum, or, in some cases perhaps, provided with paired or triple polar flagella. If we start with Cohn's classification in the year 1872 we may keep the name *Bacterium* for schizomycetous organisms of this type, and at the same time we shall not be doing any violence to the older use of the word of Ehrenberg, who figures and describes this kind of an organism. Thus the writer proposes to do, substituting *Bacterium* (Cohn emend.) for *Pseudomonas* Migula and for more recent names proposed by others.

Smith has attempted to follow Cohn's procedure in getting cultures of *Bact. termo* and succeeded by inoculating Cohn's solution with water into which beans had been thrown. A green fluorescent organism with a polar flagellum and incapable of gelatin liquefaction was secured.

It is interesting to note that by an entirely distinct line of reasoning Vuillemin came to a similar conclusion. He notes that Ehrenberg

based the genus upon the type of movement of the cell. Today, he states, we know the characteristic type of motion to be due to the presence of polar flagella on short rods. He concludes that the *Bact termo* is closely related to *Bact pyocyaneum* and accepts this as the type of the genus. His result is a genus practically identical with that proposed by Smith.

A somewhat similar conclusion will be reached as the result of the examination of still another line of evidence, a study of the species of *Bacterium* other than *Bact termo* recognized by Cohn and his associates. Three species previously known were placed in the genus *Bacterium* by Schroter. One of these was named *Vibrio syncyaneus* by Ehrenberg (1840, p. 202), *Vibrio cyanogenus* by Fuchs (1841, p. 190) and *Bacterium syncyaneum* by Schroter. It is the organism which has been repeatedly described as the cause of a blue coloration in milk. It has the morphology characteristics of the genus *Pseudomonas* of Migula. The species *Bact aeruginosum* of Schroter is the earlier name of the blue pus organism variously called *Bacillus pyocyaneus*, *Pseudomonas pyocyaneas* and *Pseudomonas aeruginosa*, and chosen by Vuillemin under the name *Bact pyocyaneum* to constitute the type of the genus. The *Bact brunneum* resembles the two preceding in the formation of a water soluble pigment. The *Bact xanthinum*, or as it now more commonly known, *Bacillus synxanthus* is an organism which is frequently recorded as the cause of orange or yellow milk. It does not appear to belong to the same group with the other forms. It would seem that at least two and probably three of the organisms recognized by Cohn and Schroter in addition to the *Bact termo* forms which can be recognized today and studied readily, would conform to the generic diagnosis used by E. F. Smith and by Vuillemin.

A serious objection to this conception of *Bacterium* is the emphasis placed upon the flagella and motility. Smith would include, for instance, in his genus *Bacterium*, a green fluorescent monotrichiate rod but would exclude from the genus another organism which produced the same green fluorescence, had similar colonies and growth reactions in the various media, which in short, was culturally, physiologically, and morphologically identical except that it was non-motile. A genus ought not ordinarily to be constituted upon one character alone. It should group together closely related organisms. Disobedience to this principle would require the botanist to remove the Lombardy poplar from the genus *Populus* because of its constant failure to produce the seeds and fruits characteristic of the genus. Conformity with all other

characters of the genus, however, is of course regarded as outweighing the loss of a character which is generally emphasized as of great importance

Strict application of the code would seem to necessitate the designation of *Bacterium triloculare* Ehrenberg as the type. Certainly, then, the designation of a more recently described species as the type would seem to require international sanction. Probably the best solution will be to adopt the suggestion of Breed, Conn and Baker, and not attempt to include the genus in our classifications until the original species can be recognized. This seems to have been followed by Castellani and Chalmers (1919) and Bergey *et al* (1923) and may perhaps be recognized as a general tendency due to the advisability of splitting the unwieldy colon-typhoid series of bacteria into a number of genera.

Bacterius. A genus of bacteria proposed by Kendall (1902, p. 484) to include those rod-shaped bacteria which are motile, but on which the distribution of flagella is unknown. He states that this group would disappear as knowledge of the flagellation becomes more complete. The name has never come into use, and has not been applied to a species. It may be regarded as invalid.

Bacteroidae. A tribe (the sixth in the family *Bacillaceae* named by Castellani and Chalmers (1919, pp. 932, 959)) with the single (type) genus *Bacteroides*.

The tribal definition is

Bacillaceae with good growth on ordinary laboratory media, without endospores, fluorescence, or pigment formation, and obligatory anaerobes.

Type Genus *Bacteroides* Castellani and Chalmers, 1918

These authors again use this name in 1920 (p. 601).

Bergey *et al* (1923, p. 255) recognizes this as the ninth tribe of the family *Bacteriaceae* with the following description: "Motile or non-motile rods, without endospores. Show good growth on ordinary culture media, without pigment formation. Obligatory anaerobes."

One genus, *Bacteroides* is included.

Bacteroides. A generic name proposed by Castellani and Chalmers (1919, p. 959), for the sole genus of the tribe *Bacteroidae*, with the following description:

Definition *Bacteroidae* with the tribal characters

Type Species *Bacteroides fragilis* Veillon and Zuber

The type is found in abscesses from various parts of the body. Another species is the well-known *Bacteroides fusiformis* of Le Dantec and Vincent, found in

hospital gangrene, and in Vincent's angina and many other conditions, as well as in the mouths of healthy persons and in the tartar on teeth

It will be noted that the fusiform organism of Vincent's angina is included. Apparently the authors were unaware of the previous designation of *Fusiformis* (q v) as a generic name for this group. Eight other species are described as belonging to *Bacteroides*.

Whether *Bacteroides* is to be regarded as a valid genus depends on whether *Bacteroides fragilis* constitutes a generic type which should be placed in a genus other than *Fusiformis*.

Bactrella. A generic name proposed by Morren (1830) for an organism apparently to be placed among the bacteria. He states that the genus *Vibrio* of Bory-de-Saint-Vincent is divided into three subgenera "les lamellinaires, les gordioides et les oxyuroides." The first named of these is elevated to generic rank and named *Bactrella*. He (p 354) gives the following descriptions

Corpus simplex, elongatum, cylindricum, vel utroque extremo obtusum, vel antice tenuiter, postice e contra admodum attenuatum, undique clausum, vel omnino mobile

The genus contained *Bactrella undula* (*Vibrio undula* Mull), *Bactrella bacillus* (*Vibrio bacillus* Mull), and *Bactrella filum*. The description of the last species is

B corpore longissimo filiformi cylindrico aequali vel simplici vel bino dum procreat) extremis obtuso, rotundato vel purumper ancinato praesertim in motibus, libero, recto, extenso rigido immobile vel parum erret languide natanti, affixo autem flexuoso, plicatulo, gyris sese contorquenti apice mobiliore perarrequieto, forma paulisper mutabile Individua libera vel affixa, solitaria aut appropinquata immenso numero corporibus immersis affixa et intermixta

This generic name apparently has never been recognized by later writers. It may be regarded as invalid although possibly the *Bactrella bacillus* was a *Bacillus*. Trevisan (1879 p 137) states "Resterebbe a vedere, non avendo il libro allomano, se *Bactrella* di Morren sia identico con *Metallacter*, nel qual caso la denominazione proposta da Morren dovrebbe essere conservata." Because first described, *Bactrella undula* (Muller), Morren may be regarded as the type.

Bactridium. A generic name proposed by Kunze (1817) for a hyphomycetous fungus. It was later used by Wallroth (1833, p 315) for two species of fungi. The genus has been commonly accepted by subsequent writers on mycology. As a generic name for bacteria *Bactridium* was proposed by Fischer (1895, p 139) to include those rod-shaped bacteria

of the subfamily *Bactridei* which possess diffuse flagella and in which the cells containing endospores retain their cylindrical shape. Later (1897, p 32) the diagnosis read "Motile, peritrichous, in some spores as yet unknown. Very numerous representatives, e.g., *B. subtilis*, *B. megatherium*, *B. vulgaris* (old genus *Proteus*) *B. typhi* and *B. coli*." The second edition of the *Vorlesungen* (p 60) does not materially change the diagnosis.

Chudjakov (1896) named an anaerobe *Bactridium butyricum*.

Erwin F. Smith (1905, p 158) states that the generic name *Bactridium* has been used several times in botany as well as in zoology.

A *Bactridium hypolyticum* was named by Huss (1908, p 474). It is evident that upon grounds of priority Fischer's use of *Bactridium* is not tenable.

(For reason for not considering *Bactridium* as a homonym of *Bacteridium* see the latter.)

Bactrillei. A name used by Fischer (1895, p 139) for a subfamily of the *Bacillaceae*. It was defined to contain motile rod-shaped bacteria with lophotrichous flagella. The subfamily contains the genera *Bactrillum*, *Plectrillum*, *Clostrillum* and *Arthrobactrillum*.

The name is not in correct botanical form.

Bactrillus. A genus name proposed by Kendall (1902, p 484) to include rod-shaped bacteria motile by means of a single polar flagellum. The name has never been used, and has not been applied to a species. It is evidently invalid, (i.e., a hyponym).

Bactrillum. A genus name proposed by Fischer (1895, p 139) to include those rod-shaped forms belonging to the subfamily *Bactrillei* which are motile by means of a tuft of polar flagella and which produce spores without causing any enlargement of the mother cell. This genus was redefined in 1897 (p 32) as follows: "Motile rods with lophotrichous ciliation. Includes provisionally *Bact. cyanogenus*." In the second edition of his *Vorlesungen*, page 60, it is noted that the spores of many species are not known. Of forms belonging to this genus, the organisms commonly known as *Bact. syncyaneum* and *Bact. fluorescens* are noted. This genus is practically synonymous with *Pseudomonas* Migula. It has never come into general use, and is commonly regarded as invalid. It is rejected by Erwin F. Smith (1905).

Probably *Bactrillum pseudo-termo* Fischer is to be regarded as the type.

Bactriniei. A subfamily of the family *Bacillaceae* created by Fischer (1895, p 141) to include all rod-shaped organisms motile by means of a

single polar flagellum. Four genera are included, *Bactrinum*, *Plectrinum*, *Clostrinum* and *Arthrobactrinum*. This name is not in correct form.

Bactrinum. A genus of bacteria proposed by Fischer (1895, p. 41). This genus is one of four belonging to the subfamily *Bactrinea*. The organisms belonging to this genus are rod-shaped and motile by means of a single polar flagellum. Spores are produced without any alteration in the shape of the mother cell.

Bactrinum nov. gen. Beweglich, mit polarer Einzelgeissel und Endosporen in nicht aufgeschwollenen Stäbchen. Eine eingeisselige Form mit Endosporen ist noch nicht ausführlich beschrieben, dagegen kann mit grosser Wahrscheinlichkeit der *Bacillus pyocyaneus*, der nach Löffler eine Geissel trägt, hierher gestellt werden, ebenso der von Löffler abgebildete grosse Bacillus aus einem Pflanzenaufguss. Ich selbst habe bis jetzt keine solche Form gefunden. Auch die von Winogradsky beschriebenen Nitritbakterien von Zürich und Buitenzorg könnten hierher gehören.

Later the genus was defined (1897, p. 32) (1903, p. 60) as motile, monotrichous with a terminal cilium and including provisionally all monotrichous rods whose spores are as yet unknown, e.g., *Bact pyocyaneum*.

This genus name has never come into common use. It is rejected by Erwin F. Smith (1905). It is generally regarded as invalid, and a synonym of *Pseudomonas* Migula. Possibly *Bactrinum pyocyaneum* (Löffler) Fischer may be regarded as the type.

Bactrinus. A genus of bacteria proposed by Kendall (1902, p. 484) to include those rod-shaped organisms motile by means of a tuft of polar flagella. This name has never come into common use and has probably not been used for any species of bacteria. It may be regarded as invalid, and a synonym of *Pseudomonas*.

Baculogenae. A subfamily of the family *Schizomycetaceae* proposed by Trevisan (1885).

The diagnosis of this subfamily as given by De Toni and Trevisan (1889, p. 939) is as follows:

Adsunt tres status evolutionis h. e. baculi, filamenta, cocci. Baculi (status primarius, initialis seu archetypus) filamenta et cocci gignentes, filamenta (status secundarius transitorius) nunquam vaginata, nec basi distincta praedita, non affixa nec e puncto centrali radiantia, e prolongatione indefinita singuli baculi v. e. plurimorum baculorum conjunctione (ob divisionem imperfectam) efformati.

The subfamily is divided into the following tribes: *Bacilleae* (12 genera), *Klebsielleae* (4 genera). This designation of the subfamily

has apparently not been used by other authors. It is not in correct form. It is not derived from a generic name.

Baculogene. A variant of *Baculogenae* used by Trevisan (1889, p 11) for the second suborder of the bacteria. Two tribes, *Bacillee* and *Klebsiellee* are included.

Bakteriaceen. A name used by Hueppe (1886, p 142) to designate a "Gattung," of rod-shaped bacteria "mit Bildung endogener Sporen" with the two subgenera *Bacillus* and *Clostridium*. It may be regarded as a variant spelling of *Bacteriaceae* (q v).

Bakterium. An aberrant form of the spelling of the genus *Bacterium* used by some German authors as Hueppe (1891, p 30).

Balkanella. A generic name proposed by Castellani and Chalmers (1919, p 934) for the seventh genus of the tribe *Ebertheae*. It is defined (p 940)

Definition *Ebertheae* which ferment glucose completely with the production of acid and gas, lactose not fermented. Milk clotted.

Type Species *Balkanella coagulans* Castellani, 1916

It is also defined later (1920, p 609)

Batteriacee. A spelling of *Bacteriaceae* used by Italian writers, particularly Lanzi (1876, p 259) and Trevisan (1889, p 1).

Beggiatoa. A genus of bacteria created by Trevisan (1842, p 56) to include certain filamentous water bacteria. The generic definition given is

Thallus e filis muco obvolutis, liberis, oscillantibus, simplicibus, elasticis, rigides, arachnoideis, punctis asteriscefornibus, primum in fascias dispositis deinde inordinatis, notatis conflatus

Two species are described, the *B leptomitiformis* and *B punctata*. The genus is included under Sub IV, *Lyngbiondeae*, Tribus VII *Lyngbeae*. The genus is variously placed by different authors with the *Algae* and with the *Fungi*. Frequently it has been grouped with *Oscillatoria* among the blue-green algae. The type species may probably be designated as *Beggiatoa alba* (Vauch) Trev (named *Beggiatoa punctata* by Trevisan in 1842). Kuetzing (1849, p 237) included *Beggiatoa* as the first subgenus of *Oscillaria*, using the following subgeneric description "Trichomata tenuissima laba, obsolete articulata vel continua, substantia interna gonimica granulosa opaca." Seven species are described. Rabenhorst (1865, p 94) gives the following generic diagnosis

Trichomati simplicia, tenuissima, hyalina, mucro involuta (non-vaginata), libera, solitaria vel aggregata, rigida, oscillantia cytioplasma pallidissime albidum (subachromaticum), punctis asterisciformibus, primum in fascias dispositis, deinde inordinates, notatum

Seven species are listed

Cohn (1875, p 173) included this genus in the tribe *Nematogenae* with the description "Zellen in Faden geordnet, Zellfaden stets unverzweigt, frei oder verfilzt, faden cylindrisch, farblos, undeutlich gegliedert, starker, lang" Cohn also noted the presence of sulphur granules in the cells

Luerssen (1879, p 24) differentiates this genus from *Bacillus* by the relatively large, long, more or less definitely jointed filaments, more or less filled with dark granules, resembling in many ways *Oscillaria* among the *Cyanophyceae*, and showing a similar oscillating movement, forming slimy masses in sulphur springs, etc

Trevisan (1879, p 138) includes this in his subtribe *Eumbrioneae*, with the following description "Somatia cylindrica, plus minus distincte articulata valida, elongata, filiformia, recta, laxe aggregate" Six species are described

The genus is included among the bacteria by several writers before 1888, including Magnin (1879), Winter (1880, p 57), Van Tieghem (1884, p 1114), Grove (1884, p 36), Zopf (1885, p 95), and Flugge (1886, p 396) Hansgirg (1888, p 263) included this as the only genus of his subfamily *Beggiatoeae* He divided the genus into two subgenera *Eubeggiatoa* and *Chromobeggiatoa*

Winogradsky (1888, p 17) as a result of his study of the sulphur bacteria made the generic diagnosis somewhat clearer Previous descriptions were not all in harmony He described the genus as made up of exclusively colorless, sheathless, always freely motile filaments never attached to a substratum The filaments are quite variable in thickness, they show a uniform intercalary growth, and have no differentiation between base and tip Under normal growing conditions they always contain sulphur granules in quantities varying with the conditions These granules appear in the form of round oily drops, which consist of a semi-fluid or soft sulphur, crystals of sulphur are never observed in the living filament The definite jointing of the filaments can be seen in cells full of sulphur only after the sulphur has been dissolved out The differentiation of good species on the basis of size Winogradsky considers to be difficult

The diagnosis of Winogradsky has been quite generally accepted by subsequent writers on systematic bacteriology, among others by Schroeter (1886, p 171), Trevisan (1889, p 10), De Toni and Trevisan (1889, p 935), Hueppe (1891, p 30), Ludwig (1892), Clements (1894)

Migula (1894, p 238) stated

Beggiatoa Trevisan, Zellen mit Schwefelkörnchen Diese letzere Familie und Gattung schliess sich auch in ihrem anatomischen Bau so eng an die *Oscillarien* an, dass sie besser mit diessen vereinigt und die Schizophyceen zugewiesen wird

The definition of Winogradsky is also used by Chester (1889, p 64), Fischer (1897, p 32), Migula (1900, p 1040), Chester (1901, p 379), Schmidt und Weis (1902), Kendall (1902, p 484), Fischer (1903, p 62), E F Smith (1905, p 162), Orla-Jensen (1909, p 334), Frost (1911, p. 60) and Heim (1911, p 254)

The following diagnosis of E F Smith taken from Migula will give the modern conception of the genus

Threads destitute of a sheath, formed of flat discoidal cells, free, *i e*, not attached Multiplication by folding and separation of the threads Motile by means of an undulating membrane as in *Oscillaria* The organism creeps along, but at the same time rotates around the long axis, mostly with a swinging of one or both free ends Habitat, hot sulphur springs and other fluids in which hydrogen sulphide is developed No reliable method is yet known for the separation of the species The number and size of the included sulphur granules are not of specific value They depend on the amount of hydrogen sulphide in the water

Buchanan (1918, p 464) designated the type species as *Beggiatoa alba* (Vaucher) Trevisan

Bergey *et al* (1923, p 409) include this as the second genus of the family *Beggiatoaceae* with the following description

Threads sheathless, formed of flat, discoidal cells, not attached Multiplication by transverse splitting of the threads Show undulatory creeping Cells contain granules of sulphur

The type species is *Beggiatoa alba* (Vaucher) Trevisan

Beggiatoaceae. A family of bacteria created by Migula (1894, p 238) It included the single genus *Beggiatoa* The diagnosis given in (1895, p 41) follows "Fadenbakterien, faden ohne Scheide durch undulierende Membran wie *Oscillaria* beweglich Zellinhalt mit Schwefelkörnchen. Bildung von Conidien nicht sicher beobachtet"

In his "System der Bakterien" Migula (1900, p 1039) includes in the family those genera of sulphur bacteria that have the cells in filaments, destitute of bacteriopurpurin As thus defined it contains both the genera *Beggiatoa* and *Thiothrix*

This family name has been commonly used by recent bacteriologists who have discussed *Beggiatoa* (*qv*), among them Chester (1901, p. 379), Schmidt and Weis (1902, p. 92), Kendall (1902, p. 486), Migula (1904, p. 145), Corsini (1905), Erwin F. Smith (1905, p. 162), Clements (1909, p. 7), Frost (1911, p. 60), Schneider (1912, p. 23), and Bergey et al. (1923, p. 407)

Buchanan (1918, p. 463) included three genera, *Thiothrix*, *Beggiatoa* and *Thioploca*

Beggiatoeae. A subfamily of the family *Leptothrichaceae* proposed by Hansgirk (1888, p. 263) The group is defined as follows: "Faden mit lebhaften, oscillarien-artiger Bewegung, im protoplasmatischen Zellinhalte Schwefelkörnchen und nicht selten auch einen rosenrothen oder violetten Farbstoff enthaltend" One genus only, *Beggiatoa*, is included

Betabacterium. A generic name proposed by Orla-Jensen (1919, p. 175) for certain lactic rods The genus is described in a somewhat informal fashion, but the following quotations will probably serve to identify the group

The betabacteria (Table XXXI) are in most respects so closely allied to the betacocci that they may be regarded as the analogous rod forms, and they can, like the betacocci, be divided into those which ferment arabinose (Nos 1-20) and those which do not (Nos 21-33) The *Bacterium casei* [varieties] formerly described by me are typical representatives of these groups, and as they also differ in morphological respects—the arabinose-fermenting rods being generally shorter than the others—we consider ourselves justified in establishing the species *Betabacterium breve* and *Betabacterium longum* Besides occurring in vegetable matter, they are also found in cheese, faeces and kefir grains The rods forming the tissue of the kefir grains (Nos 1 and 2) are, however, so different from the remaining betabacteria that they must undoubtedly be reckoned as a distinct species which we will call *Betabacterium caucasicum*, as the kefir rods are now for the most part known under the name of *Bacterium caucasicum* This last name is also erroneously used for streptobacteria, which are likewise also found in kefir (though not as a rule in the grains themselves), and which are considerably easier to obtain in pure cultures

Most of the betabacteria are not altogether killed by heating until 75° They form inactive lactic acid, at times with a surplus of dextro-lactic acid The great majority of strains develop gas (carbonic acid and more or less hydrogen) When the development of gas is strong, succinic acid is also formed The power to form any considerable quantity of gas, and thus succinic acid, is, however, soon lost in artificial nutritive substrates, and in most of our strains, gas development could only be observed by sowing out closely in sugar agar tubes From laevulose, some few strains can form a small amount of mannite, and as the mannite-forming mash bacterium *Lactobacillus fermentatum*, which has been closely studied by Jan Smit, has the greatest resemblance to the betabacterium (judging from

its fermentation of sugar, with *Bbm longum*) it should doubtless be reckoned under this head. Possibly the mannite bacteria so dreaded in the making of wine also belong here. Some of these form, like the betacocci, slime from cane sugar, which the betabacteria investigated by us never do.

The betabacteria grow poorly in milk, and do not as a rule attack casein at all. When cultivated at a freshly isolated state, were able to curdle milk, and even with a slight development of gas, but this power was soon lost. The frequent occurrence of these bacteria in cheese is due to the fact that they are better able to utilize lactate of lime as a source of carbon than are most other lactic acid bacteria.

A characteristic feature in the betabacteria is their lack of ability to ferment salicin and alcohols, and, with the exception of the *Bbm longum* forms, their preference for pentoses. The betabacteria have also always a slight fermentation of mannose. Some of the strains (as for instance Nos. 6 and 7) which in a freshly isolated state fermented mannose comparatively strongly, lost this power later on.

Betacoccus. A genus proposed by Orla-Jensen (1919, p. 146) for certain lactic acid cocci. He says,

the betacocci are found in green vegetable matter and juicy roots. They are introduced with vegetable food into the intestinal canal of animals, and pass thence into the milk. In the retting process we always encounter arabinose-fermenting betacocci, which might be connected with the fact that pectin substances always contain an arabinose group. As the betacocci are far more variable in all respects than the streptococci, it is very difficult to divide them up into clearly defined species, and I therefore prefer to treat the genus *Betacoccus* under one head and merely note in conclusion what features might seem to justify our uniting certain strains into independent species.

The betacocci can as a rule stand heating to 60° but rarely to 65°. In a slimy state, however, they can stand higher temperatures, as the slime protects them, and it has been observed at sugar factories that thin syrup which had been heated to 80° to 85°, and could not possibly have become infected afterwards, has grown slimy (No. 11 had formed zooglea masses under such conditions). The optimal temperature lies at about 30° or under, a single strain (No. 14) was even found to grow best at indoor temperature, and this temperature is, as in the case of *Sc cremoris*, the most favorable one for slime formation. The maximal temperature is 35° to 37° (rarely 40°) and the minimal 5° to 7°. Some few strains (Nos. 1, 45, 46, and 47) grow, however, at 45°, but on the other hand thrive but poorly below 15°.

The betacocci always form laevo-lactic acid, more rarely also equivalent quantities of dextro-lactic acid, so that we find inactive lactic acid (Nos. 43-47).

The betacocci also as a rule develop gas, (carbonic acid with more or less hydrogen). The gas development is strongest in laevulose solutions, and next in cane sugar solutions.

In most strains, however, the gas development is so slight that it can only be observed by sowing out strongly in tall sugar agar tubes, and a few strains (Nos. 1, 2, 3, 38, 36, 37 and 43-47) do not seem to develop gas at all. Several of these last grow chiefly in the upper part of the agar tube, though they are not otherwise obligatorily aerobic. On

ordinary cane sugar agar, the slime develops but poorly, but appears in a very characteristic manner on cane sugar gelatin. Large colonies, clear as water, appear on the plates, resembling the colonies of certain aerogenes species (the slimy aerogenes forms produce, however, slime from all the sugars which they ferment) and in stabs, we get very characteristic pictures. Though these bacteria do not liquefy ordinary gelatin, and are not provided with other proteolytic qualities, several of them can, after some length of time, liquefy cane sugar gelatin, which figure is indicated in the tables by 1 (*liquare*).

As the genus *Betacoccus* contains all possible degrees of sliminess and liquefying power, we cannot attach too much importance to these characteristics and we may find cases where of two strains, otherwise entirely alike (as Nos 38 and 39) which were, moreover, found in the same sample of material, one will liquefy and the other not.

When isolated from vegetable matter, the betacocci thrive as a rule but poorly in milk, when isolated from milk, on the other hand, or from dairy products, and sometimes from dung, they can form comparatively large quantities of acid in milk, and even dissolve some casein (Nos 29 and 34). The power of souring milk, however, is comparatively soon lost, but can be regenerated by continued transference from milk to milk. The bacteria are often abundantly supplied with lactase, and it may happen that nearly all the lactose of the milk is hydrolyzed without any considerable quantity of it being fermented, which shows that the proteins of the milk are a poor source of nitrogen. When isolated from beets, they prefer beet juice to casein peptone (Nos 11 and 12).

The betacocci exhibit a certain preference for pentoses. Strains isolated from vegetable matter for the most part ferment both xylose and arabinose whereas those isolated from dung, milk, or dairy products, will as a rule ferment only one of the two, or sometimes no pentoses at all. Of the hexoses, they often prefer laevulose, and of the disaccharides, often saccharose. They frequently ferment raffinose, but of true polysaccharides, only a little dextrin at the outside. With regard to salicin, the different strains vary considerably. When they do not ferment raffinose, then in most cases they will not ferment salicin either. The betacocci do not as a rule attack alcohols to any perceptible degree, only a few strains ferment a little mannite (No 20 even a little sorbite).

No other bacteria have proved so variable with regard to the sugars as the betacocci.

Without cane sugar, the betacocci form small colonies, and are in all cultural respects indistinguishable from the streptococci. Morphologically also, they resemble the latter though the faculty of dividing in two directions is often more developed.

The above mentioned morphological differences between the AO and A + X-forms on the one hand, and the X- and O-forms of the other, render it likely that we have here to deal with two distinct species. As the former always (at any rate unless in a weakened state) ferment arabinose, we will term them *Betacoccus arabinosaceus*, and as the latter (especially the typical X-forms) can be isolated from most cowdung after enrichment in acid sugar broth, we will call them *Betacoccus bovis*. As the betacocci are for the most part known under the name of *Streptococcus mesenteroides*, it would have been reasonable to use the name *Betacoccus mesenteroides* for one of the species, had it not been that both comprise

strains which do not form slime, and consequently also no mesentery. It is possible that the betacocci should be divided into more than the two mentioned species. With all experiments in this direction, however, I have felt that I was working on treacherous ground, as these bacteria exhibit such great variability in almost all respects.

Winslow (1920, p. 102) regards this as synonymous with *Leuconostoc* Von Tieghem.

Billetia. A genus of bacteria founded by Trevisan (1889, p. 11) upon the species *Bacterium Laminariae* of Billet (1888, p. 293). The genus was reduced to synonymy with *Kurthia* by De Toni and Trevisan (1889, p. 931). The genus name has apparently never come into use. It should probably be considered a "*genus incertae sedis*" until the original species can again be identified and its right to generic rank decided. The generic description reads "*Kurthia* sporifera con vagine distante." It is included as the sixth genus of his tribe *Leptotrichiae*.

Bollingeria. A genus proposed by Trevisan (1889, p. 26) but without description. One species, *Bollingeria equi* is included. The diagnosis given by De Toni and Trevisan is as follows (1889, p. 1034).

Discomyces Rivolta (1884), *Botryomyces* Bollinger (1886) nomen tantum et ut *Discomyces* valde ineptum, *Botryococcus* Kitt (1886) nec Kuetz (1849). Cocci globosi v. divisionis tempore globoso ovoideis, cystidibus specialibus destituti, numerosissimus, in muco matriciali nidulantes, segregati, in familias globosas magnas, singulas cystidibus universalibus amplis crassis lamellosis summe firmis cartilagineis involutas, inordinate consociati. Glebae globosae ovoideae vel piriformes, omni aetate intus solidae, in acervos permagnos solidissimos. Ruborum fructus externam faciem quodammodo simulates, densissime cumulate. Coccorum divisio in tres directiones.

Two species are given, *Bollingeria equi* Trevisan (synonyms *Zoogloea pulmonis equi* Bollinger 1870, *Discomyces equi* Rivolta 1884, *Micrococcus ascoformans* Johne 1885, and *Micrococcus botryogenes* Rabe (1886)) and *B. Vacchetae* Trevisan (*Discomyces pleuriticus canis familiaris*, Rivolta 1884). The type is *Bollingeria equi* Trev.

The genus has never come into common use, and may probably be suppressed as a synonym. It is rejected by Erwin F. Smith (1905).

Borrelia. A generic name proposed by Swellengrebel (1907, p. 582) to include those forms of the family *Spirochaetaceae* in which the cells are provided with peritrichous flagella. The type species is given as *Borrelia (Spirochaeta) gallinarum*. If this organism should stand as a generic type, the name would appear to be valid but possibly it should be reduced to a synonym of *Spiroschaudinnia*,

Treponema or *Spirochaeta* It is probably based upon a false morphological assumption (peritrichous flagella). Gleitsmann (1913, p. 31) says

Da Sambon erst in August 1907 mit der Bezeichnung *Spiroschaudinna* herauskam, Swellengrebel dagegen bereits in Juni desselben Jahres auf Grund der Borselschen Ausführungen das Virus *Borreha* taufte, so muss der Streit über den neuen Namen des Virus zugunsten der *Borreha* entschieden werden

Botryococcus. A generic name first proposed by Kuetzing (1849, p. 892) for an algal genus in the *Chlorophyceae* It has been used by most subsequent writers on this group of algae, among them Rabenhorst (1864, p. 42), De Toni (1889, p. 674) and West (1904, p. 237) It must be regarded as a valid genus of algae containing several well known species

The name *Botryococcus* was introduced independently by Kitt (1888, p. 247) for the organism *Micrococcus ascoformans* The name as thus used is invalid It is not infrequently used as a casual designation of the organism causing botryomycosis It is given as a synonym of *Bollingeria* by De Toni and Trevisan (1889, p. 1039)

With the spelling botryocoques this organism is discussed by Nicolle and Cesari (1914, p. 229) These authors conclude that *Botryococcus* is a synonym of *Staphylococcus* It is evident that it is not a valid bacterial generic name

Botryomyces. A generic name proposed by Bollinger (1888, p. 177) for the organism variously termed *Micrococcus botryogenes* Rabe *M. ascoformans* Johne and *Discomyces equi* Rivolta The description is summarized by Enlows (1920, p. 24) as follows

Similar macroscopically to *Actinomyces* Under strong magnification the grapelike colonies consist of conglomerate, roundish, varying-sized broom-like groups or heaps of micrococci (4 to 4.5 μ) Each colony (150 to 250 μ) is surrounded by a homogeneous capsule-like membrane, and in this respect resembles the *Ascococcus* of Billroth Ray-fungus-like, fibrous masses produced at times in host

Type species (monotypy) *B. ascoformans* (Johne) (Bollinger's earlier name was *Zoogloea pulmonis equi*) Syn (Bollinger) *Micrococcus botryogenes* Rabe, 1886, *M. ascoformans* Johne, 1885, *Discomyces equi* Rivolta and Micellone, 1879

The name is termed inept by De Toni and Trevisan (1889, p. 1039) and made a synonym of *Bollingeria*

The name is rejected by E. F. Smith (1905)

Botulobacillus. A genus of bacteria proposed by Orla-Jensen (1909, p. 343) as the second in the family *Putribacteriaceae* It is defined

as including anaerobic bacteria with peritrichous flagella, producing an exotoxin which exerts specific action upon the central nervous system. The species do not attack lactose. The organism causing botulism and tetanus belong here. Probably *Botulobacillus botulinus* Orla-Jensen may be regarded as the type.

The genus name must yield priority to *Clostridium* and *Plectridium*, and is probably to be regarded as invalid, unless the type species is to be separated.

Brachybacterium A generic name used by Troili-Petersson (1903, p 138) to include certain lactic acid bacteria. The organisms are separated into two genera, *Bacterium* and *Brachybacterium*. The diagnosis of the latter genus is "Short, oval or ellipsoidal rods whose length is not more than double their width, round forms are present in addition to the rods." In *Bacterium*, on the other hand, the rods are in length more than double their width.

Nine species of *Brachybacterium* are described. Most of them are designated by number only. No 22 is the *Bacterium lactis acidii* of Leichmann or *Bact. lactis* of Lister. One species, No 26 is named *Brachybacterium apiculatum*.

It is possible that this genus should be reduced to a synonym of *Lactobacillus* (qv), or of *Streptococcus* (qv). Apparently no type species has been designated.

Brachybacterien. A term used by Troili-Petersson according to Lohms (Handbuch, p 198).

Brucella. A generic name proposed by Meyer and Shaw (1920, p 176) for the causal organisms of Malta fever and infectious abortion. They state,

On morphologic grounds the organisms of undulant fever and of infectious abortion of domesticated animals must therefore be considered as identical and must be placed together in the genus *Bacterium*. For reasons to be given in detail in the second paper it is proposed in accordance with the suggestions made by Buchanan (1918, p 8) of the Committee on Classifications of the Society of American Bacteriologists that a genus, for which we propose the name "*Brucella*" be created in the family *Bacteriaceae* to separate and to distinguish properly these important pathogenic microorganisms from the other members of the recognized genus *Bacterium* which is already overburdened with representatives, which have little in common with the *B. melitensis* and *B. abortus*.

Evans (1923, p 1944) discussed in detail the generic designation of the group. She says

That the Malta fever and contagious abortion organisms are closely related was confirmed by Fleischner, Meyer and Shaw in 1919, and later by a number of other investigators. Meyer and Shaw (1920) proposed the generic name "*Brucella*," in the family *Bacteriaceae* to include the Malta fever and contagious abortion organisms. That name has met with general approval, and has been used by foreign investigators (Khaled, Archibald).

Meyer and Shaw did not, however, give a generic diagnosis for the genus *Brucella*, and they did not consider other species besides the *melitensis*-*abortus* group which would logically belong to the new genus. They were apparently unaware that Castellani and Chalmers had already described a newly created genus, "*Alkaligenes*," which, according to its definition, would include the *melitensis*-*abortus* group. The definition is as follows: "*Bacillaceae* growing well on ordinary laboratory media, not forming endospores, aerobes, and often facultative anaerobes, without fluorescence, pigment formation, or gelatin liquefaction, without polar staining, Gram-negative, without a capsule. Milk not clotted, glucose and lactose not fermented." The type species of the genus "*Alkaligenes*" as established by "original designation," is *A. faecalis*, a common intestinal saprophyte. Castellani and Chalmers left the "*Micrococcus melitensis*" unclassified generically—"Incertae sedes"—because they were doubtful as to whether it should be considered a coccus or a rod form.

Bacteriological nomenclature is passing through an experimental, transitional stage, and criteria which should serve for valid specific or generic distinctions have not been established in detail. The characteristics which are available at present for bacterial classification are few, they are more or less variable according to varying conditions, and they are not well correlated. Hence comes the difficulty in classifying bacteria into a system comparable with that of the higher organisms. It appears that, in many cases, distinction must be made by means of the sum total of differences, rather than by differences in particular stable characteristics, such as obtains in the classification of higher organisms.

If *melitensis* is considered cogenetic with *faecalis*, then the Law of Priority demands the acceptance of the generic name *Alkaligenes*. If, however, the differences between the two species are sufficient for generic distinction, the valid name of the genus is *Brucella*. The writer is of the opinion that a generic distinction should be made between the organisms under discussion, which are characteristically invaders of the tissues of animals, and the type species of *Alkaligenes*, which is characteristically a saprophyte. Morphologically, also, *melitensis* is readily distinguished from *faecalis* by its smaller size and by its great numbers of coccoid cells.

There is given herewith a general description of *melitensis*, the type species of the genus *Brucella*.

Minute rods with many coccoid cells, (the cells of two-day cultures grown on the surface of plain agar and stained with carbol fuchsin appear about 0.5 of a micron wide and 0.5 to 2 microns long), not forming endospores, non-motile, aerobic, or preferring a slightly reduced, partial pressure of oxygen, without gelatin liquefaction, Gram-negative, parasitic, invading animal tissues, neither gas nor acid production from the carbohydrates.

Other species, which vary somewhat from the foregoing description, will logically be allocated to the genus *Brucella*. This genus should include a variety of

small rod-forms commonly present in freshly drawn cow's milk. These forms were described in an earlier publication (Evans 1918) as bacteria related to "*Bacterium abortus*". Some of them vary from the typical *Brucella* in the production of a slight amount of acid from the carbohydrates. The genus *Brucella* should also include the species *bronchisepticus*, which varies from the typical *Brucella* in being motile. A number of other species should also probably be classified in the genus *Brucella*.

Meyer and his associates have continued the use of the abbreviation "B" for the generic designation of *meliensis* and *abortus*. Since that is the accepted abbreviation for the genus *Bacillus*, there should be some other for the *Brucella*. The abbreviation *Br* is proposed for that genus.

But the point may be made here that whenever a generic name is referred to in any paper it should be printed in full the first time it is cited in the article.

Every investigator since 1918 who has compared the Malta fever and the contagious abortion organisms has found a close relationship between them. The literature on this subject is reviewed in the following paper. The accumulating evidence of the close relationship between the strains of bovine and human origin has culminated in the conclusion by Burnet, that *meliensis* and *abortus* are not distinct bacteriological species, but merely distinct serological varieties or subspecies of one and the same species. All the literature on the subject leads to this conclusion. Observing priority of publication as determining the nomenclature, we must adopt as the specific name for the *meliensis*-*abortus* group the name *meliensis*.

Butyl-Bacillus. A casual name used by Fitz (1878, p. 48) to designate bacteria capable of producing butylic fermentation.

Butyribacillus. A generic name proposed by Orla-Jensen (1909, p. 342) to include the butyric acid bacteria. It is the first genus of the family *Butyrobacteriaceae*. These forms are anaerobic peritrichous rods which as a rule split certain kinds of sugar with the development of gas and butyric acid, and do not ferment pectin or cellulose, and do not attack proteins actively.

Nach Pringsheim¹ ist *C. Pasteurianum* nur ein einzelner Repräsentant einer weitverbreiteten Gattung von Erdbakterien, und laut einer vorläufigen Mitteilung von Bredemann² können alle echten Buttersäurebakterien zur Stickstoffassimilation befähigt werden. Eine Gruppierung derselben in solche, die sich mit anorganischen Stickstoffquellen begnügen können, und in solche, die nur gedeihen, wenn sie organische Stickstoffquellen zur Verfügung haben, ist deshalb nicht statthaft, sondern wir müssen alle diese Organismen, wozu nach Grassberger und Schattenfroh auch *Bacillus Chauvoei* gehört, unter der Gattungsbezeichnung *Butyribacillus* sammeln. Wahrscheinlich sind die ältesten obligat anaeroben Bacillen auch die ältesten Sporenbildner, denn ohne die Fähigkeit zur Erzeugung von Endosporen wurden diese Bakterien im allerhöchsten Grade der

¹ Centralbl. f. Bakt., Abt. II, Bd. XX, 1908, p. 248

² Berichte der Deutschen botanischen Gesellschaft, Bd. XXVIa, 1908, p. 362

Vernichtung durch den Luftsauerstoff ausgesetzt sein, sobald der Abbau des Substrats, in welchem sie sich entwickelt haben, vollbracht ist

Apparently no type species has been designated, although *Butyribacillus Pasteurianus* Orla-Jensen is the first mentioned

Butyribacteriaceae. A family name proposed by Orla-Jensen (1909, p 343) to include the genera *Butyribacillus*, *Pectobacillus* and *Cellulobacillus*. These organisms are all anaerobic, peritrichous, rod-shaped bacteria which ferment carbohydrates with production of gas and butyric acid

Butyriclostridium. A generic name suggested by Orla-Jensen (1921, p 273). He says

The use of the term *Clostridium* as a generic name presents the inconvenience that under the same we must group together not only the clostridia but also the plectridia. The division into true butyric acid bacteria (*Butyriclostridium*), the requirements of which in regard to nitrogenous nutriment are very moderate (they are able to assimilate even the nitrogen of the air) and anaerobic, putrefying bacteria (*Putriclostridium*) seems natural to me

No species is mentioned

Byolysis. A generic name suggested by Salisbury (1868)

Enlows states

A minute "algoid vegetation developing in and on the human body in typhoid fever". Flourishes with great luxuriance in the "agminated and solitary glandules of Peyer". The spores multiply by duplicative segmentation, and develop rapidly on and in the cells of the epidermic and mucous surfaces. Spores frequently found in the colorless corpuscles, destroying their contents and dilating their walls, so that the cells are often from 2 to 4 times normal size

Several of the figures on the plate illustrating this genus resemble streptococci. Salisbury refers to a filamentous stage in this genus only in the figure descriptions where he states that the more mature stage is filamentous. Marchand (Bot Crypt, t 1, 1883, p 471) thinks this genus is synonymous with *Crypta* Salisbury

Calymmatobacterium. According to Enlows (1920, p 25) a generic name proposed by Beaurefare-Arago and Vianna (1913, p 222). She says

Type species (monotypy) *C. granulomatis*. Encapsulated coccus 0.2 to 0.3 μ in diameter, or rods with rounded ends of 0.5 to 2 μ in length, also encapsulated. Prior to division the rod presents a median constriction, appearing as a diplococcus. Found in granulomata

Carboxydomonas. A genus proposed by Orla-Jensen (1919, p 311) to include autotrophic bacteria which are able to oxidize carbon monoxide. This is the second genus of his family *Oxydobacteriaceae*

Winslow *et al* (Committee Soc Am Bact, 1917, p 551, 1920, p 201) use the following description "Autotrophic rod-shaped cells capable of securing growth energy by the oxidation of carbon monoxide (forming carbon dioxide)" The type species, *Carboxydomonas oligocarbophila* (Bejerinck and van Delden) Jensen, is described as non-motile

Bergey *et al* (1923, p 33) use this description, including this as the third genus of the tribe *Nitrobactereae*

Carphococcus. A generic name used by Hohl (1902, p 338) to include a bacterial species which he termed *Carphococcus pituitoparus*. The organism was isolated from straw It is a coccus varying from 0.7 to 1.5 μ in diameter It grows readily upon culture media Milk develops a very marked slimy consistency and within fourteen hours the upper layers become decidedly viscous Acid is not produced and the milk is not digested

It does not seem that Hohl had sufficient justification to warrant the creation of a new genus The name has not been accepted by other writers and may be regarded as a synonym of *Micrococcus*

Carteria. Merrill and Wade (1919, p 64) state

The adoption of this new name (as "*Carteria*") was tentatively suggested by Musgrave and Clegg as possibly advantageous for the purpose of avoiding further controversy, although they did not definitely advocate this highly informal procedure

It was proposed as a substitute for *Actinomyces* See *Carteria*

Carteria Musgrave, Clegg and Polk (1908, p 470) in a discussion of the validity of *Actinomyces*, *Streptothrix*, etc, conclude *Streptothrix* to be the correct name They say "In making this decision we are fully aware of the rights of those who favor *Actinomyces* or *Nocardia*, and under the circumstances are tempted to introduce a new name (*Carteria*) for the genus"

See also *Carteria*

Caryobacterium A generic name proposed by Mori (1913 p 1) for a species *C. equi* found associated with Brustseuche in the horse. The organism is described as a coccobacillus, straight or slightly curved, isolated or in chains, 2.5 x 0.5 μ , motile, staining readily Gram positive and without capsule Aerobic and facultative anaerobic, gelatin not liquefied, no indol, milk not coagulated, does not attack maltose, but ferments lactose with production of gas and acid In cells stained with methylene blue, or better with Giemsa a definite nucleus is visible,

whence the name *Caryobacterium*. The nucleus is not permanent and is visible only from cultures in peptone-carbohydrate (preferably maltose) and a polyvalent alcohol (mannitol)

Caseobacterium A genus proposed by Orla-Jensen (1909, p 311) to include those gram positive rod-shaped lactic acid bacteria which for the most part attack or degrade the casein, forming monoamino acids. These forms are facultative or anaerobic. All are non-motile. The only species definitely ascribed to this genus is *Caseobacterium casei* which may be regarded as the type. It is placed in the family *Acidobacteriaceae* of the order *Cephalotrichinae*.

This generic name may be a synonym of *Lactobacillus* which has priority over it. The name has not come into general use.

Cellulobacillus A genus proposed by Orla-Jensen (1909, p 343) to include those anaerobic peritrichous bacteria which ferment cellulose, and are capable of producing butyric acid and gas from sugars. No species is definitely ascribed to this genus by the author. It is included as the third genus of the family *Butyribacteriaceae*.

The author states

Von der Gattung *Butyribacillus* müssen wir ebenfalls die cellulosevergärenden Bakterien, die Gattung *Cellulobacillus*, ableiten. Während diese interessanten Mikroorganismen sich einerseits mit Ammoniak als einziger Stickstoffquelle begnügen können, zeigen sie andererseits eine eigentümliche Spezialentwicklung dadurch, dass sie imstande sind, ein so widerstandsfähiges Kohlenhydrat wie die Cellulose anzugreifen, und dass sie von den anaeroben Bakterien die Kohlenhydrat am besten ausnutzen (siehe die Methangärung). Da *Cellulobacillus* somit einen Kulminationspunkt bezeichnet, können wir nicht ohne zwingende Gründe hiervon andere Gattungen ableiten sondern müssen diese Gattung als einen insolierten Seitenzweig betrachten.

Enlows (1920, p 26) gives the type species (monotype) as *Cellulobacillus methanigenes*.

Cellulomonadeae. A name proposed for the third tribe of the family *Bacteriaceae* by Bergey *et al* (1923, p 154). The description follows:

Short rods, occurring in soil, having the property of digesting cellulose. Motile, or non-motile. Chromogenic or non-chromogenic. Growth on ordinary culture media often not vigorous. Gram-negative.

There is a single genus *Cellulomonas*.

Cellulomonas. A name proposed by Bergey *et al* (1923, p. 154) for the single genus of the tribe *Cellulomonadeae*.

The description is as follows

Small rods, with rounded ends, non-spore-forming, motile or nonmotile, occurring in soil and having the property of digesting cellulose

The type species is *Cellulomonas brazoteus* (Kellerman)

Cenomesia. A generic name proposed by De Toni and Trevisan (1889, p 1039) with the following diagnosis

Cocci globosi vel divisionis tempore globosovoides, modice numerosi, in mucos matricali nidulantes, segregati, in familias globosas parvas singulas cystidibus universalibus ampliusculis, crassiusculis homogeneis, non lamellosis, firmis, gelatinosis involutas, sociati Familiae e coccis ad peripheriam cumulatis compositae, demum intus medio inanes Cystides speciales nullae Coccorum divisio, initio generationum serierum, in duas directiones

Obs *Coelosphaerium* Naegeli, *Algarum* Phycochromophycearum genus, exacte analogum

The genus is included in the subfamily *Coccogenae* and the subtribe *Eu-Ascococceae* Two species are described, both named by Trevisan, *Cenomesia alba* and *C lilacina* Both contain sulphur granules and the latter contains bacteriopurpurin

This name does not seem to have come into common use The writer has not been able to find any other author who has discussed these species It is possible that the descriptions are not sufficient to make possible identification, and that the name *Cenomesia* is to be regarded as a *nomen nudum* More probably it is to be regarded as a synonym of the genus *Lamprocystis* It is rejected by Erwin F Smith (1905, p 174)

No type species has been designated, although *Cenomesia alba* Trevisan from its position may be regarded as such

Cephalotrichinae. The name given the first order of bacteria created by Orla-Jensen (1909, p 325) He includes in this order microorganisms with spherical, rod-shaped or spiral cells Endospores are found only in a few sulphur-free spirilla The cells may be motile or non-motile, if the former either *monotrichous* or *lophotrichous* They are typically water bacteria They secure their growth energy almost exclusively by oxidative processes Many of the members of this group develop poorly in laboratory media because the organic constituents are present in too concentrated a form

The order is divided into seven families as follows *Oxydobacteriaceae* (7 genera), *Actinomycetes* (4 genera), *Thiobacteriaceae* (4 genera), *Rhodobacteriaceae* (12 genera), *Trichobacteriaceae* (7 genera), *Luminobacteriaceae* (5 genera), *Reducibacteriaceae* (2 genera)

Chatinella. A generic name proposed by Roze (1898, p 69) for an organism observed microscopically in decaying plant tissue (straw in stable manure and horse dung) The cells measured 12 to 27μ in diameter, multiplying by fission A membrane 1 to 3μ in thickness enveloped the cells One species (type by monotypy) is *Chatinella scissipara* In spite of the unusual size and characteristics, Roze assigns this organism to the fission fungi Its position is doubtful

Chionyphe. A generic name given to a fungus closely related to *Mucor* by Thienmann (1839) Its only interest to bacteriology is through the association by Carter of a *Chionyphe Carteri* Berkeley with madua foot as a probable etiologic factor Castellani and Chalmers (1919, p 2114) say

He placed some black particles, taken from a food, on cotton soil moistened with animal juices and enclosed in a stoppered bottle, which he left unopened for two and three-quarter years, when he found a thin reddish film had appeared Other black particles sown on rice paste for the same length of time remained unchanged, but on opening the bottle a red mould speedily made its appearance

With reference to this mould, he says "It had not, however, a clear connection with the fungus particles, but seemed to spring up independently of them upon the rice whenever this was exposed to the air"

This statement is of importance, as he grew a fungus from the white variety which was pink in colour and produced sporangia resembling those of a species of the genus *Mucor* Micheli, 1729, but differing therefrom in the absence of a columella, which should have brought it under the genus *Mortierella* Coemans, 1863, but Berkeley, who examined the growths from a botanical point of view, classified it under the genus *Chionyphe* Thienmann, 1839, calling it *Chionyphe carteri* Berkeley, 1862, and defining it as "Hyphasmate ex albo flavorubroque, sporangium demum coccineis, sporis breviter fusiformibus"

The genus *Chionyphe* however, was never recognized by mycologists generally, as its species came under the genera *Mortierella* or *Mucor*, while *Chionyphe carteri* was most undoubtedly a contamination, as its connection with the black or white grains was never proved, as we have noted above with regard to the former

Chlamydatomus A genus of cocci in the subtribe *Sarcineae* created by Trevisan (1879, p 137) The generic diagnosis is as follows

Cellulae globosae, divisionis tempore ovoideae, inordinate in colonias conglobatas pluristratas densissime consociatae, 1-4 tegumentis propriis gelatinosis crassiusculis confluentibus obvolutae Coloniae tegumento communi desituta

One species is listed, *Chlamyda Bergelii* Trevisan He later (1889, p 27) included it as the fifth genus of the tribe *Ascococceae*

De Toni and Trevisan (1889, p 1042) give the following description

Cocci globosi, divisionis tempore ovoidei, in muco matricali nidulantes, singuli v 2-4 cystidibus specialibus gelatinosis, crassiusculis, persistentibus involuti. Cystides in acervos numerosos conglobatos, intus solidos, dense et inordinate aggregati. Cystides universales nullae. Coccorum divisio in unam directionem. Sporae ignotae.

The genus is included in the subtribe, *Gaffkyaee*, of the tribe *Ascococceae* of the subfamily *Coccogenae*. Two species are described *Chlamydatomus Beigelii* (Kuch. Rabenhorst) Trevisan (Synonyms, *Pleurococcus Beigelii* Kuchenmeister and Rabenhorst 1867, *Sclerotium Beigelianum*, Haller 1868, *Zoogloea Beigeliana* Eberth, 1873, *Hyalococcus Beigelii* Schroeter (1886)) and *Chlamydatomus cellaris* (Hansg.) Trevisan (Synonym, *Hyalococcus cellaris*, Hansg. (1880)).

Chlamydatomus Beigelii is the type (monotypy). It is now not regarded as a bacterium at all, but as a fungus, a member of the genus *Trichosporum*. Castellani and Chalmers (1919, p. 1103) recognize *Trichosporum beigelii* Vuillemin 1901 as the cause of piedra, in nodosities on the hairs of the moustache.

It would seem that Vuillemin was in error in substituting the new generic name *Trichosporum* for the older generic name *Chlamydatomus*.

Chlamydobacteriaceae. A family name proposed by Migula (1894, p. 237) to include those genera of bacteria which are characterized by the possession of firm membrane or sheath surrounding the filaments. The description is as follows:

Familie *Chlamydobacteriaceae* nov. Fam. Formen von sehr verschiedener Entwicklungsstufe, aber alle ausgezeichnet durch eine feste Hülle oder Scheide, welche die zu verzweigten oder unverzweigten Fäden vereinigten Zellen umgibt.

The filaments may be branched or unbranched. The family as first described contained the following genera: *Streptothrix* Cohn emend., *Phragmidothrix* Englel., *Crenothrix* Cohn, *Cladothrix* Cohn, and *Thiothrix* Winogr. This family name was used by Chester (1897, p. 64). In the second volume of his System, Migula (1900, p. 1030) gives the following diagnosis of the family:

Zellen cylindrisch zu Fäden angeordnet, die von einer Scheide umgeben sind. Vermehrung erfolgt durch bewegliche oder unbewegliche Conidien, welche direkt aus den vegetativen Zellen hervorgehen und, ohne eine Ruheperiode durchzumachen, zu neuen Fäden auswachsen.

The family is used without material alteration in definition by Chester (1901, p. 369), Schmidt and Weis (1892, p. 92), A. J. Smith (1902, p. 272), Kendall (1902, p. 484), Migula (1904, p. 145), E. F.

Smith (1905, p 161), Ellis (1909, p 7) Clements (1909, p 7) and Frost (1911, p 59)

Vuillemin (1913, p 526) places this family among the *Schizophyceae* and not among the bacteria

Buchanan (1918, p 301) and Bergey *et al* (1923, p 390) included this as the only family of the order *Chlamydobacteriales* q v. Five genera were listed, *Leptothrix*, *Didymohelix*, *Crenothrix*, *Sphaerotilus* and *Clonothrix*

Castellani and Chalmers (1919, p 924) list this as the fourth family of the order *Eubacterales*

It should be noted that this name does not conform to the nomenclatural rule that the name of a family should be fashioned from one of the constituent genera

Chlamydobacteriales A name proposed by Buchanan (1917, p 162) for the second order of the class *Schizomycetes* The description given follows (1918, p 301)

Filamentous bacteria, alga-like, typically water forms, frequently sheathed, without true branching although false branching may be present The sheath is frequently impregnated with iron Conidia may be developed, but never endospores Sulphur granules or bacteriopurpurin never present Mature cells or filaments not motile, not protozoan-like

The order contains a single family *Chlamydobacteriaceae*

The name was accepted by Winslow *et al* (Committee Society of American Bacteriologists (1917, p 551)), and by Bergey *et al* (1923, p 390)

Chlamydobacterinae A variant of *Chlamydobacteriaceae* used by Schmidt and Weis (1902, p 375) as a family name

Chlamydothrix A genus name proposed by Migula (1900, p 1030) to replace the name *Streptothrix* previously used by this author, and to include the two species commonly known as *Leptothrix ochracea* Kützinger and *Gallionella ferruginea* Ehr

Fischer (1903, p 61) used the following description "Nicht fest gewachsen, schwarmende Cylinderomiden, Beispiel *Chl ochracea*"

The genus is characterized by Smith (1905, p 162) as follows

Cells cylindric, non-motile, arranged in unbranched threads, with a sheath of varying thickness Frequently the septation of the threads is only demonstrable after the use of reagents Reproduction by means of non-motile roundish or ovoid conidia, which arise directly from the vegetative cells Synonyms *Streptothrix* (Cohn) Mig, *Leptothrix* Kütz *ex p* and *Gallionella* Ehrenberg *ex p*

Ellis (1909, p 156) regards the genus as synonymous in part with *Leptothrix* (*Chlamydothrix ochracea*) and *Gallionella* (*Ch ferruginea*) Mohs (1910, p 21) recognizes the genus *Gallionella* as distinct, and includes Migula's species *Chlam ochracea*, and a new species which he names *Chlam sideropous* His description of *Chlam ochracea* is as follows

Faden bestehend aus cylindrischen farblosen Zellen, mit anfangs farbloser dünner, später, dicker, gelb- und braunwerdener Scheide Junge Faden 0.9 μ Durch Inkrustation mit Eisenoxydhydrat wird die Scheide allmählich dicker, gelb bis rostbraun und der Faden erreicht heerdurch die doppelte oder manchmal sogar die mehrfache Dicke der Zellen Durch Behandlung mit 2-5 per cent Salzsäure löst sich das Eisenoxyd, die Gallerthülle wird undeutlich und die Zellen, von einer zarten Innenschicht der Hülle umgeben, treten scharf hervor Die nach einiger Zeit zur Ruhe kommen, sich gewöhnlich festsetzen und wieder zu Faden auskeimen mit deutlicher Spitze und Basis

Die Schwärme setzen sich häufig an schon vorhandene Faden fest, keimen hier zu Faden aus und tauschen hier Verzweigung kommt vor, aber selten und nie so regelmässig wie bei *Cladothrix*

Alte Scheiden entleeren ihre Zellen, bleiben als rostrote Röhrchen übrig und bilden dann oft kolossale Mengen ockeriger Massen in eisenhaltigen Wassern und mitunter auch in Rasenerzen

Frost (1911, p 59) gives the following generic description

The cells are cylindrical, non-motile, and arranged in unbranched threads and surrounded by a sheath of varying thickness in different species, being the same diameter at apex and base Reproduction takes place by means of gonidia, which are round and arise directly from the vegetative cell This genus is called *Leptothrix* by Kutzing and *Streptothrix* by Cohn

It is probable that *Chlamydothrix* should be reduced to a synonym of *Leptothrix*, (*g v*) for the latter name is apparently valid as a generic name for this form of iron organism Orla-Jensen (1909, p 331) takes this view

Enlows (1920, p 21) designates the type in the following statement

Cells cylindrical, non-motile, arranged in unbranched threads surrounded by a sheath of varying thickness Septation of threads often demonstrable only after the use of reagents Reproduction by means of round or ovoid, nonmotile conidia, which arise directly from the vegetative cells Syn (Migula) *Streptothrix* (Cohn) Migula, *Leptothrix* (Kutzing) exp, *Gallionella* Ehrenberg exp

Type species *C ochracea* (Kutzing) He also places here *C ferruginea* (Ehrenberg), *C hyalina* Migula, *C epiphytica* Migula, *C fluitans* Migula

Chlorobacterium. A genus named by Guillebeau (1890, p 32) for the species *Chlorobacterium lactis*, isolated from an inflamed udder.

The organism is actively motile, cells about 1μ by 3μ . Gelatin is rapidly liquefied. Green pigment is produced upon potato and nutrient agar.

The type species (monotypy) was designated by Enlows (1920, p. 27) as *Chlorobacterium lactis*.

Chlorobacteriaceae. A family proposed by Lauterborn (1913, p. 99) to include the chlorophyll-producing bacteria.

His reasons for its establishment he gives as follows:

Wenn man den sehr schwefelwasserstoffreichen Schlamm am Grunde von Characeenteichen durchmustert, fallen neben den blaugrünen Filzen der Cyanophyceen sowie den roten Wucherungen der Purpurbakterien vielfach auch kleinere oder grössere Flocken von gelbgrüner Farbe auf. Dieselben bestehen neben Fäden von *Oscillatoria chlorina* Kützinger, *O. putrida* Schmidle und *O. Lauterbornii* Schmidle besonders aus Ansammlungen kleiner Zellen von gelbgrüner Farbe in gallertigen, bisweilen ziemlich lockeren Verbänden, die bisher, soweit beschrieben, der Gattung *Aphanotheca* zugezählt wurden. Hierher gehören *Aphanotheca luteola* Schmidle sowie *A. clathrataformis* Szafer. Ein genaueres Studium dieser Formen hat in mir die Überzeugung befestigt, dass dieselben nicht zu den eigentlichen blaugrünen Algen gehören, sondern sich direkt den Schizomyceten anschliessen und hier eine den Purpurbakterien durchaus gleichwertige Gruppe bakterienartiger Organismen mit gelbgrünem Farbstoff (Bakteriochlorin) darstellen. Wir hatten es hier also mit einer neuen Familie der Schizomyceten zu tun, die man als *Chlorobacteriaceae* den *Rhodobacteriaceae* gegenüberstellen kann.

He included in this family the genera *Chlorochromatium*, *Pelodictyon*, *Schmidlea*, *Pelagloea* and *Pelochromatium*. The spelling should be *Chlorobacteriaceae*. It is probably invalid.

Chlorobium. A generic name proposed by Nadson (1906, p. 190) according to Enlows for

A green chlorophyll containing organism, which author thinks belongs with the bacteria or close to *Stichococcus bacillaris*.

Type species (monotypy) *C. limicola*. Cocci 0.4 to 0.5 μ in diameter, round or elliptical, or short rods. Non-motile. Multiplication by cross division. Long chains common in both the round and rod forms. Involution and apochlorotic forms occur.

Chlorochromatium. A genus of bacteria proposed by Lauterborn (1906, p. 197) in the following description:

Chlorochromatium aggregatum nov. gen. nov. spec. Gestalt ungefähr elliptisch bis spindel- oder tonnenförmig, an den Enden etwas verschmälert und abgestutzt. Farbe grüngelb wie bei *Oscillatoria chlorina*. Bei schwacher Vergrößerung der Korperrand scheinbar etwas krenuliert und dunkler grün gefärbt als die Mitte. Bei sehr starker Vergrößerung erscheint der Organismus zusammengesetzt aus

einer Anzahl kleiner gestreckt elliptischer, bakterienartiger Körper, welche eine starker grün gefarbte Rindenschicht und eine hellere centrale Partie erkennen lassen. Indem diese Körper, in etwas unregelmässigen lockeren Langsreihen angeordnet, mantelförmig einen axialen farblosen (wohl gallertigen?) Hohlraum umschliessen, kommt das oben geschilderte Aussehen bei schwacher Vergrösserung zustande. Freibeweglich mit Hilfe von Geisseln. Länge des Organismus 0,009–0,012 mm. Breite 0,005–0,007 mm.

Vermehrung durch Querteilung. Der Organismus streckt sich in die Länge und schnürt sich in der Mitte durch eine Ringfurche quer durch. Auch mehrfach Zerfall in die einzelnen bakterienartigen grünen Körper beobachtet, welche sich selbst wieder quer teilen können. Auf diese Weise kommen unregelmässige, bisweilen netzartig durchbrochene und dann im Habitus an *Lamprocystis rosea-persicina* erinnernde hellgrüne Kolonien zustande.

Later (1913, p. 99) he included this genus in his family *Chlorobacteriaceae*. Its position among the bacteria is questionable.

Chloronium. A generic name proposed by Buder (1914, p. 80) with the species *Chloronium mirabile*. The type species was designated by Enlows as follows:

Type species (monotypy) *C. mirabile*. Found in water in the Leipzig Botanic Garden. A cylindrical rod 0.7 to 1 μ by 1 to 2 μ with rounded ends, sometimes slightly curved, green, united into a zoogloeal mass, in the center of which is a colorless, spindle-shaped, 1-polar flagellate organism. The latter measures 2 to 2.5 by 5 μ . Multiplication by transverse division. The "peripheral component" is sometimes of coccus form also green of 0.75 μ diameter. Usually about 10 to 30 arranged in rather definite order about the colorless central organism. Perfiliev (J. Mic. Biol., v. 1, 1914, p. 223) says that the peripheral forms described here are identical with *Chlorochromatium aggregatum* Lauterborn. Buder is in doubt as to the systematic position of *Chloronium*. Perfiliev describes the "central organism" of Buder's *Chloronium* as *Cylindrogloea*, q. v.

Chondromyces. A genus credited to Berkeley and Curtiss by Berkeley (1857, p. 313) in an illustration, figure 70, which has the explanation "*Chondromyces crocatus* Berkeley and Curtiss, with its spores. Both magnified. From specimens on a decayed gourd from South Carolina." In the text the genus is not described, but is classified under the *Hyphomycetes*, in the family *Isariaceae* Corda-Stern. Under *Isaria* Fr. (p. 314) is the following statement:

Several genera which find their proper place in this highest group, and the last are, in fact, compound forms of organisms which appeared under other tribes—*Chondromyces* and *Stigmatella* are compound Mucedines.

The first description was published by Berkeley (1874, p. 64) as follows: "Stipes e floccis compactus ramosus induratus sporae apicales."

Saccardo (1886, p 576) gives the following diagnosis "Stipes ... hyphis compactus, ramosus, induratus, *Conidia apicalia*, capitulata, elongato-ovata, continua"

The true nature of this organism was first recognized by Thaxter (1892, p 401) He designated it as one of the genera of the *Myxobacteriaceae* (g v) His diagnosis is as follows "Rods forming free cysts in which they remain unmodified Cysts various, sessile, or borne on a more or less highly developed cystophore." This genus is distinguished from the related genus *Polyangrium* Lk. in that the latter forms its one or more cysts within a gelatinous matrix raised about the substratum, and from the genus *Myxococcus* in that no coccusslike spores are formed by the fruiting bodies

Thaxter gives the following synonymy "*Stigmatella* B and C in Berk, Introduct Crept Bot. p 313 fig 70, b (no description) 1857, do in Grevillea 3 97 1874 (first description)

(?) *Polycephalum* Kalch and Cke in Grevillea 9: 22 1880.

(?) *Cystobacter* Schroter in Kryptogamen Flora v Schlesiens Die Pilze, p 170 "

The species described by Schroter was named *Cystobacter erectus*. What is evidently a species practically or quite identical with *Chondromyces crocatus* was described by Zukal (1896, p 346) as *Myxobotrys variabilis* (q v) This author also regarded it as a fungus

Buchanan (1918, p 542) diagnosed the genus as follows

Rods forming free cysts in which they remain unmodified Cysts various, sessile or borne on a more or less highly developed cystophore

The type species is *Chondromyces crocatus* Berkeley and Curtiss

This description is also used by Bergey *et al* (1923, p 416)

Chorynebacterium. Apparently a misprint for *Corynebacterium* found in the index of Migula's System der Bakterien (1900, vol 2, p 1061)

Chromatiaceae. A subfamily of sulphur bacteria created by Migula (1900, p 1047) to include three genera *Chromatium*, *Rhabdochromatium* and *Thiospirillum* He states "Zellen frei, zeit lebens schwarmfahig" The subfamily is characterized as containing those sulphur bacteria in which the cells are free and capable of swarming at any time

The subfamily has been generally recognized by those who have discussed classification of the sulphur bacteria such as Migula (1904, p 146) E F. Smith (1905, p 164) and Frost, (1911, p 61)

It may be noted that the correct designation of the subfamily would be *Chromatioideae* (q v)

Chromatieae. The fifth tribe of the subfamily *Chromatioideae* named by Buchanan (1918, p 470) The diagnosis follows, "Sulphur bacteria in which the cells are not united into families, but free, and capable of swarming at any time" The tribe is the equivalent of Migula's *Chromatiaceae* q v Five genera were included, *Chromatium*, *Rhodobomonas*, *Thiospirillum*, *Rhodocapsa* and *Rhodotheca*

This usage is followed by Bergey *et al* (1923, p 400)

Chromatioideae. The first subfamily of the family *Rhodobacteriaceae* named by Buchanan (1918, p 464) with the description "Cells not filamentous, containing both sulphur granules and bacteriopurpurin

The subfamily is practically the equivalent of *Thiorhodaceae* q v

Five tribes are included, *Thiocapsee*, *Lamprocystee*, *Thiopedeae*, *Amoebobacterieae*, and *Chromatieae*

Bergey *et al* (1923, p 394) have followed in the use of this term

Chromatium A generic name created by Perty (1852, p 179) with the following diagnosis

Korper ausserst klein, cylindrisch, roth, braun, violett, grun gefarbt, im ausgebildeten Zustand mit innem Blaschen (Blastien) Ein Bewegungsfaden am Vorderende? Vermehren sich durch Quertheilung

Two species were described, *C Weissii* and *C violascens* He states the former to be related to the *Monas Okenii* of Weiss, but smaller The genus was recognized and more fully described by Schroeter (1886, p 156) as containing those organisms with cylindrical-elliptical or relatively thick cylindrical cells multiplying by fission, and with cell contents colored red, and containing black (sulphur) granules He states that this genus is placed with the bacteria only provisionally, for it has many resemblances to the monads Winogradsky (1888, p 86) includes the genus among his sulphur bacteria He accepts the Schroeter diagnosis of the genus and recognizes the best known species *Chromatium Okenii* In this species the cells are very large, averaging $15\text{ }\mu$ in length and $6\text{ }\mu$ in maximum width The cells are somewhat variable in shape, straight or more or less bent, short cells ovoid and longer forms more cylindrical The cells are motile by means of polar flagella Other species are *Chr weissii*, *Chr minus*, *Chr vinosum* and *Chr minutissimum*

The genus is included in classifications of sulphur bacteria in the subfamily *Chromatiaceae* by Migula (1900, 1904), E F Smith (1905,

p 164) and Frost (1911, p 61) Migula (1900, p. 1047) states, "Zellen cylindrisch-elliptisch oder elliptisch, verhältnissmassig dick" It should be noted that E F Smith (1905, p 205) also included *Chromatium* in his list of rejected bacterial names. Orla-Jensen (1909, p 330) says "Die sehr hellroter schwefelbakterien (wie Z. B. *Chromatium vinosa*) bilden den natürlichen Uebergang von *Thiomonas* zu *Rhodomonas*" He makes it a synonym (p 334) of *Rhodomonas*

Buchanan (1918, p. 470) and Bergey *et al* (1923, p 400) gave the following diagnosis

Cells cylindric-elliptical or relatively thick cylindrical Cell contents red, containing dark sulphur granules Cells somewhat variable in shape, straight, more or less bent, short cells ovoid and longer forms more cylindrical Motile by means of polar flagella

The type species is *Chromatium okenu* Perty

Chromatoideae. A variant spelling of *Chromatioideae* used by Bergey *et al* (1923, p 394)

Chromobacillus. The second section of the genus *Bacillus* recognized by Hansgirk (1888, p 265) with the following description "Einzelne Zellen scheinbar fast farblos, in grosserer Menge (Zoogleen) rosen bis blut roth, blau etc gefarbt" The *Bacillus sanguineus* Schroeter is given as the only species It was found "In Sumpfen bei Chlumec und Magdalena nachst Wittengau und bei Neu-Bestritz nachst Neuhaus in Sudbohmen"

It apparently should rank as a subgenus with the type *Bacillus* (*Chromobacillus*) *sanguineus* Schroeter

It is possible that the type organism belongs with the purple bacteria

Chromobacteriae. A tribe named by Winslow *et al* (Committee Soc of American Bacteriologists 1920, p 208) with the diagnosis "Water bacteria producing a red or violet pigment" The genera *Erythrobacillus* and *Chromobacterium* were included

The spelling probably should be changed to *Chromobacteriaceae*

Bergey *et al* (1923, p 85) recognize it as the first tribe of the family *Bacteriaceae* with the following description "Water and soil bacteria producing a red, yellow, violet or blue-green pigment"

Four genera are included.

Chromobacterium. A revision of the spelling of *Cromobacterium* Bergonzini (q v) used in a review (anonymous) of Bergonzini's work (Journ Roy Mic Soc, 1881, p 285) and by Grove (1884, p 26) in the synonymy of *Bacterium violaceum* This form is also quoted by De Toni and Trevisan (1889, p 978)

Buchanan (1918, p 52) emended the description of the genus as follows

Rod-shaped bacteria, without spores, aerobic, producing a violet chromoparous pigment soluble in alcohol but not in chloroform, motile or non-motile, Gram stain variable

The type species is *Chromobacterium violaceum* Bergonzini

The genus was recognized by Winslow *et al* (Committee Soc Am Bact, 1920, p 209) with the following description

Aerobic bacteria, producing a violet chromoparous pigment, soluble in alcohol but not in chloroform Motility and Gram reaction variable

Type species, *Chr violaceum* Bergonzini

Bergey *et al* (1923, p 117) include this as the third genus of the tribe *Chromobactereae*

If the violet bacteria should have generic recognition the name is apparently valid

Chromobeggiatoa. The second section (subgenus) of the genus *Beggiatoa* recognized by Hansgirk (1888, p 264) with the following descriptions "Faden von rosen-bis-pfirsch rother Farbe, rosen-oder blutrothe, violette bis violettbraunliche Ueberzuge an im Wasser faulenden Substanzen bildend, oder das Wasser roth farbend" The only species listed is *Beggiatoa roseo-persicina* Zopf

Chromococcus. A revision of the spelling of *Chromococcus* Bergonzini *q v* used in an anonymous review in Journ Roy Mic Soc, (1881, p 284)

This form is also used by De Toni and Trevisan (1889, p 1067) in the synonymy of *Streptococcus violaceus* The type species (monotypy) is *Chromococcus violaceus* Bergonzini

Cladochytrium. A genus of fungi named by Nowakowski (1876)

The name *Cladochytrium tuberculorum* was proposed by Vuillemin (1888, p 121) for the fungus which he believed was the primary cause of nodule formation on the roots of leguminous plants The bacteroids were concluded to be simple differentiations of the plasma The infection threads he believed to be fungal in nature See *Rhizobium*

Cladophytum. A generic name used by Leidy (1849, p 227) for two species of entophytes from the intestine of *Julus* (thousand leg)

The original description of the genus, and the first species follows

Filaments minute, attached by means of a roundish nucleus, simple or compounded near the base of attachment, with minute lateral ramuli, inarticulate, and with no evidence of interior structure

Cladophytum comatum Filaments delicate, regular, colorless, simple, more frequently branched near the base at very acute angles, growing in more or less dense bunches from a yellowish rounded or oval, attached, nuclear body varying in size from $\frac{1}{16}$ to $\frac{1}{8}$ of an inch. Lateral ramuli very minute, measuring in length from $\frac{1}{16}$ to $\frac{1}{8}$ of an inch, and passing off at acute angles. No indication of articulation or interior structure.

Length from $\frac{1}{16}$ to $\frac{1}{8}$ of an inch.

Habitat Growing more or less profusely from the mucous membrane of the small intestine of *Julus marginatus*, occasionally from the same surface at the commencement of the large intestine, from any part of the exterior surface of entozoa infesting those cavities, and also from any part of the surface of *Enterobrus elegans*.

A second species, *Cladophytum ramosissimum* was later (1849, p 250) described by Leidy, but still later (1851, p 37) was made a synonym of *Cl. comatum*. The generic description (1851, p 37) is "*Thallus* attached by means of one or more granules, filamentous, simple, with minute lateral ramuli, or branched, inarticulate, amorphous in structure.

This genus is described as follows by Robin (1853, p 358)

Nouveau genre voisin des Mycodermata. On sait que le genre *Mycoderma*, de Persoon, n'était pas formé par des espèces de Champignons comme il le croyait, mais bien par des Algues appartenant aux genres *Leptomitus* et *Hygrocrocis*.

Filaments petits fixes par un noyau arrondi, simples ou compliqués vers leur point d'insertion par de petits ramuscules latéraux inarticulés, sans structure intérieure définie.

Apparently the organism has not been studied by later writers. The exact position and relationships of this organism are uncertain. The type species is *Cladophytum comatum* Leidy.

Cladothriceae. A subtribe of the tribe *Vibrionieae* created by Trevisan (1879, p 137). The description given is "*Somatia libera Pseudo-ramosa*." Two genera are included, *Cladothrix* and *Streptothrix*.

Cladothrichaceae. A name proposed by Zopf (1884, p 45) with the aberrant spelling *Cladothricheen* as a family to include the single genus *Cladothrix*. He states "Sie zeigen Coccen-, Stäbchen-, Faden- und Schraubenformen. Die Fadenform ist mit Pseudoverzweigungen versehen." In 1885, p 51, he adds "Sporenbildung nicht nachgewiesen."

Schroeter (1886, p 173) also used this name with the spelling *Cladothrichacei* and the description "Faden durch falsche Astbildung verzweigt. Die weiteren Charaktere die der VI Ordnung."

Apparently Hansgurg (1888, p 229) was the first to use the correct spelling *Cladothrichaceae*. He included it as the first family of the order *Desmobacteria*, with the single genus *Cladothrix*.

Hueppe (1891, p 31) followed Zopf, as did Baumgarten (1900, p 129).

This family name has apparently not been recognized by later writers.

Cladothrichacei. An aberrant form of spelling the family *Cladothrichaceae* (*q v*) used by Schroeter (1889, p 173).

Cladothricheae. A tribal name used by Hueppe (1885) to include the genus *Cladothrix*. A variant of *Cladothricaceae* *q v*.

Cladothricheen. An aberrant method of spelling the family name *Cladothrichaceae* (*q v*) first used by Zopf (1884, p 45) to include the genus *Cladothrix* Cohn. This designation has also been used by Zopf (1885), Hueppe (1885), Baumgarten (1900), etc.

Cladothrix. This generic name was published in De Condollés *Prodromus* (1849) for a genus of the phanerogamic family *Amarantaceae*. It has since generally been recognized as valid in this sense by botanists.

Cohn (1875, p 185) used this as a bacterial generic name to include the single species *Cladothrix dichotoma*, an organism found in foul water. It is colorless, partly floating on the water and partly attached to decaying algae. The filaments are very slender, apparently unsegmented, straight or somewhat bent, similar to a colorless *Leptothrix*. The filaments of the latter, are however, unbranched while in *Cladothrix* they are repeatedly branched and with considerable regularity. By use of high magnifications he noted that it was not a true dichotomy, but false branching, as in the blue green alga *Scytonema*. The genus was included in the tribe *Nematogenae*.

Trevisan (1879, p 139) included this as the sixteenth genus of his family *Bacteriaceae* in the tribe *Vibriomeae* and subtribe *Cladothricae* with the following description:

Somatia cylindrica, leptotrichoidea, infeme vix articulata, pupeme distincte articulata, tenerrima, stricta vel subundulata pseudo-dichotoma, microgonidia e serie cellularum divisione longitudinali et transversa succedanea multipartitarum orta, globulosa, diaphragmatibus ruptis in pseudo-sporangium densissime congesta, demum e vagina lateraliter erumpentia et in somatia evoluta, foventia.

Magnin (1880, p 97) followed Cohn's description without modification.

Winter (1880, p. 59) described the genus as *Leptothrix*-like, very slender, colorless, not jointed, straight, slightly wavy or even partially irregularly spiral, with false branching. He states, however, that he is unable to find sufficient grounds for differentiation between *Cladothrix* and *Streptothrix* and concludes that both are doubtful genera. He cites in justification for this conclusion the work of Cienkowski. Grove (1884, p. 40) followed Winter in emphasizing the resemblance of *Cladothrix* and *Streptothrix*. Zopf (1885, p. 51) made this the only genus of the family *Cladothricheen*, characterized by having growth forms as cocci, rods, filaments and spirals, with the filamentous stage showing false branching, spore formation unknown. Schroeter (1886, p. 174) revised the diagnosis of the genus, including in it filaments surrounded by delicate sheaths, non-motile, and dichotomously branched, at first apparently not jointed, later fragmenting, and the segments developing into new filaments. Hansgirk (1888, p. 229) described a new variety, *Cladothrix dichotoma* var. *leptochaetoformis*.

De Toni and Tievisan (1889, p. 927) emphasize the development of arthriospores. Their generic diagnosis reads "Filamenter basi ab apice superiore distincta, vagina crassa obducta, cylindrica, aetate protracta, pseudoramosa. Arthrosporangia binae in singulis microbaculis ellipsoideis oetae."

Hueppe (1891) defines the genus as made up of filamentous organisms, sheathed, branched, and with the filaments differentiated into base and tip. Ludwig (1892) agrees with Hueppe in not stating that there is "false" branching. He says that the branches are not swollen at the tips, free, not embedded in slime. Sauvageau and Radais (1892, p. 246) state with emphasis that the genera *Streptothrix* and *Cladothrix* are very distinct. Migula (1894, p. 237) says

Cladothrix Cohn Zellen zu pseudodichotom verzweigten Faden verbunden. Teilung nur nach einer Richtung des Raumes. Vegetative Vermehrung durch polar begeißelte Schwarmzellen, die sich fest setzen und zu neuen Faden auswachsen. Endosporenbildung bei *Cladothrix dichotoma* einmal beobachtet. Hierzu gehören auch die Gattungen *Sphaerotilus* Kütz., und *Actinomyces*.

Migula (1895, p. 40) defined the genus in somewhat greater detail as filaments with delicate sheaths, attached when young, non-motile, cells cylindric, breaking occasionally through the sheath by intercalary growth and forming false branches, a pseudo-dichotomy, multiplication through swarming gonidia, which have a cluster of flagella attached laterally near one pole. Fischer (1897), Migula (1897, 1900)

Chester (1897, 1901), Hewlett (1898), A. J. Smith (1902), Fischer (1903), Jordan (1908), Orla-Jensen (1909), Ellis (1909) and Heim (1911) have all used the term in a somewhat similar sense. Migula later decided that the genus name *Cladothrix* Cohn is synonymous with *Sphaerotilus* Kutzing. In this he has been followed by many recent writers. However, it would seem that the original descriptions of *Sphaerotilus* (q. v.) and *Cladothrix* may be sufficiently distinctive to warrant the separation of the genera. *Sphaerotilus* has its filaments disposed in considerable numbers in gelatinous branched threads, the filaments lying parallel to each other. This massing of filaments into threads does not seem to be characteristic of *Cladothrix* though doubtless the two genera are closely related.

That *Cladothrix* is worthy of generic recognition is argued by Molisch who states

Migula stellt *Cladothrix* zu *Sphaerotilus*. Ich habe vorläufig davon Abstand genommen, weil Bau und Entwicklung der letzteren Gattung noch der *Sphaerotilus natans* und nach einem genaueren Studium dieser vielleicht doch herausstellen dürfte, dass die Aufstellung der Gattung *Cladothrix* neben *Sphaerotilus* erzeuge dichotom verzweigte Fäden ohne Gegensatz von Spitze und Basis. Aber die Fäden der *Cladothrix dichotoma*, die er zu *Sphaerotilus* stellt, lassen doch deutlich Spitze und Basis unterscheiden.

Molisch's description of *C. dichotoma* is as follows

Meist farblose Flockchen, festsitzend, aus Fäden bestehend, die niemals echt, sondern in weiten Abständen wiederholt unecht dichotom verzweigt sind. Die Fäden setzen sich aus stabförmigen, ovalen oder langlichen Zellen zusammen, die von einer dünnen festen Mangelhaft bekannt sind und weil sich nach Gewinnung von Reinkulturen Scheide umgeben sind. Fäden durchschnittlich 2 μ dick, mit Gegensatz von Spitze und Basis. Indem die Stäbchen sich innerhalb der Scheide teilen, werden die vorderen nach der Spitze des Fadens gedrängt oder sie wachsen unter entsprechender Auftreibung der Scheide aneinander vorbei, wobei die beiden oder auch nur die eine der einseitig freier werdenden Polzellen selbständig weiter wachsen und die nun entstehenden Äste gewöhnlich in losem Verband verbleiben. Auf diese Weise entsteht die falsche Dichotomie. Die Vermehrung geschieht durch unbewegliche und durch schwärmende Gonidien, die gestaltlich den vegetativen Zellen gleichen und sich aus diesem dadurch entwickeln, dass sich seitlich unterhalb des einen Endes ein Büschel von Geißeln bildet. Sie schwärmen nach dem Verlassen der Scheide einige Zeit umher, setzen sich dann fest und kennen zu neuen Fäden aus. Ein namentlich in mit organischen Substanzen verunreinigten Wassern sehr häufiger und weit verbreiteter Organismus.

More or less confusion has arisen due to loose definitions of the genus such as that of Kendall (1902, p. 484) who records the genus as

that group of the family *Chlamydobacteriaceae* in which the cells are more or less branched Macé (1897, p 1024) came to the conclusion that the *Cladothrix dichotoma* of Cohn did not exist as a well defined type, and that it had been interpreted in different ways by various investigators He therefore proposed that *Cladothrix* be substituted for *Streptothrix* Cohn, because the latter generic name is a homonym of *Streptothrix* Corda His definition reads

Ces sont des Bactéries filamenteuses, a éléments droits ou sinueux dépourvus de gaine, produisant latéralement des ramifications disposées d'une façon irrégulière, dans certaines conditions, les filaments se segmentent ou très courts batonnets ou articles sphériques ou ovoïdes que l'on doit probablement considérer comme arthrospores

The conclusions of Macé have not been generally accepted, and his statements have but served to confuse nomenclature

Kligler (1915, p 327) described a *Cladothrix placoides* from the teeth Some fifty-eight strains were studied The characters are given as follows

Morphology The morphology of this organism is very variable It often appears like a coccus, sometimes like a diphtheroid rod, or short club-end thread Usually one finds all three in the same preparation The coccus-like forms are probably spores, and the others are pleomorphisms of the thread Microscopic pictures are often obtained which look identical with *L. innominata* of Miller

Chromology It stains readily with ordinary anilin dyes and is usually Gram-positive Often the spore retain the stain while the threads are decolorized when treated by Gram's method *Biological properties* They are aerobic, non-motile, non-branching threads They grow best at 37°C, only very slowly at 20°C

Cultural characters On agar they form round, raised, lustrous, white colonies which can be picked up entire from the medium On glucose-agar the colonies are usually larger and more grayish Microscopically the surface colonies are convoluted and are often seen on the plates enclosing underneath them small colonies of other bacteria In broth the growth is very sparse, and the wall of the tube is usually covered with a granular layer of discrete colonies 0.5 mm or less in diameter which are not readily removed by shaking the tube In glucose-broth the growth is very abundant, giving both a heavy turbidity and a granular deposit on the wall Gelatin was liquefied by some strains but not by others They all fermented glucose, most of them fermented sucrose, while some also fermented lactose Gas is not produced They generally reduce nitrates and fail to produce indol or ammonia When first isolated they should be kept on glucose-agar and transferred every two weeks

Buchanan (1918, p 302) rejected the name as invalid as a bacterial genus because of its use for a phanerogam

Castellani and Chalmers (1919, p 1068) recognize and describe *Cladothrix dichotoma* Cohn

It would seem that the genus *Cladothrix* Cohn with the description as emended by Migula is not valid, because of prior use by De Candolle. The name should not be used for organisms which show true branching, as *Actinomyces*, *Nocardia*, etc. It may probably be regarded as a synonym of *Sphaerotilus*.

Cladotricaceae. A tribal name proposed by Trevisan (1879, p 15) with the following diagnosis, "Sporae (arthrospora) in filamentis normalibus obvenientes Filamenta pseudoramosa, basi ab apice superiori distincta, vaginata vel evaginata." Later (1889, p 926) De Toni and Trevisan included the following genera, *Sphaerotilus* Kutz, *Cladothrix* Cohn and *Nocardia* Trevisan.

This name (with the spelling *Cladotricheae*) was also used by Sternberg (1892, p 19) to include the genus *Cladothrix*.

Cladotricaceae. A variant of *Cladotricaceae* used by Trevisan (1889, p 9) for the first tribe of the suborder *Tricogone*.

Cladotricheae. A name used by Sternberg, a variant of *Cladotricaceae*, q v.

Cladotricheen. A variant of *Cladotricaceae* q v used by Zopf (1885, p 51) to include the genus *Cladothrix*.

Clathrococcus. A generic name proposed by Schmidt and Weis (1902, p 29) to replace *Clathrocystis*, q v as a name for a bacterial genus. The generic description is (p 292),

Die einzelnen Zellen sind kugelig oder durch den Druck der sie umgebenden etwas eckig. Die Teilungen geschehen erst nach drei Raumrichtungen, später nur nach zwei, und aus den anfangs soliden Zellverbänden geht daher eine hohlkugelige Wuchsform hervor mit zuletzt netzformig durchbrochener Wand. Die Zellen sind anfangs unbeweglich, allmählich aber lösen sie sich gruppenweise ab und schwärmen mittelst Geisseln davon.

Apparently these authors have overlooked the genus *Lamprocystis* created by Schroeter for this same purpose.

The type species (monotypy) is *Clathrococcus roseo-persicinus* (Cohn) Schmidt and Weis.

Clathrocystis. A genus of blue green algae proposed by Henfrey (1856, p 52) for the species *Cl. aeruginosa* (Kützting) Henfrey. This genus was used by Cohn (1875, p 156) to include an organism which is now generally placed among the bacteria. This was named *Clathrocystis roseo-persicina*. It was found upon twigs and other plant

remains which were decaying on the bottom of standing water. It occurred in the form of cherry red flecks, which under the microscope showed a loose aggregate of small spherical or oval cells. According to Cohn, Kutzing (p. 341) first described this organism as *Protococcus*. Rabenhorst (1856, p. 28) used the name *Pleurococcus roseo-persicinus*. Cohn described the genus *Clathrocystis* as having spherical cells, united into families with reticulate structure without regular arrangement, and places it in the tribe *Gloeogenae*. The genus was noted by Flügge (1886).

Winter (1880, p. 37) separated this species from the algae, creating the generic name *Cohnia* q v. Schroeter (1886, p. 151) called attention to the fact that this name was preoccupied, and proposed *Lamprocystis* q v.

This name was accepted by Winogradsky (1888, p. 67) for a certain type of blue green algae, but it should not be used for a bacterial genus.

Clitridium. Apparently a casual name used by Billet (1890, p. 54). Enlows states,

Bacterium of average length in the form of a biscuit, "c'est-à-dire en train de se segmenter" (Possibly refers to a division stage—merely states the "Clitridium des auteurs," without any further reference.)

Cloaca. A generic name proposed by Castellani and Chalmers (1919, p. 943) for the second genus of their tribe *Proteae*. The generic description is "Slow gelatine liquefiers, ferment lactose, Gram negative."

The type species is designated (p. 958) as *Cloaca cloacae* (Jordan) Castellani and Chalmers.

Clonothrix. A genus of iron bacteria described by Schorler (1904, p. 691) to include the species *Clonothrix fusca*. The generic diagnosis is: Filaments dichotomous or irregularly branched, fixed, with base and tip contrasted, tapering somewhat to the free end. Sheath constantly present, thin in the young filaments, later thicker, and becoming impregnated with iron or manganese oxide. Cells cylindrical. Multiplication by means of small non-motile spherical gonidia which arise from the vegetative cells.

The species *Cl. fusca* produces gray to dark brown and black masses similar to *Crenothrix* and is often in association with it. He concludes that all *Cladothrix* descriptions of iron bacteria belong here. The genus is recognized by Ellis (1909, p. 158) and by Mohs (1910).

G. S. West (1904, p 335) in his discussion of *Phormidium* says "*Clonothrix gracilluma* W & G S West is one of the Schizomycetes of the genus *Cladothrix* "

Buchanan (1918, p 305) gives the following description

Filaments with false dichotomous or irregular branching, attached, with contrast of base and tip, tapering to the tip Sheath always present, thin on young filaments, later becoming thicker and encrusted with iron or manganese Multiplication by small non-motile gonidia of spherical form, formed from the disk shaped cells near tip by longitudinal division on rounding up

The type species is *Clonothrix fusca* Schorler

This is included as the fifth genus of the family *Chlamydobacteriaceae* by Bergey *et al* (1923, p 393)

Clostridiaceae. A family name proposed by Heller (1921, p 550) as a sixth family of the *Eubacteriales* with the following description

Eubacteriales that are rod-like, not spiral, that will not grow within 7 mm of the surface of a shaft of clear tissue-free agar medium contained in a tube 12 mm or more in diameter, incubated in air, in which they are able to grow in the depths They may or may not possess peritrichial flagella, they may or may not form endospores Most members of the group are characterized by their energetic catalytic action on proteins or on carbohydrates or on both of these types of substances

Two subfamilies were included, *Clostridoideae* and *Putrificoideae*

Clostridae. A subfamily of bacteria proposed by Fischer (1897, p 32) to include those forms of the family *Bacillaceae* in which the sporebearing rods are spindle-shaped Two genera are included, *Paracloster* A Fischer and *Clostridium* Prazmowski The same designation is used in the second edition of the *Vorlesungen* (1903)

This subfamily designation does not seem to have been used by other authors It may be noted that as a subfamily the designation should be *Clostridoideae*

Clostridoideae. A subfamily named by Heller (1921, p 449) as the first of the family *Clostridiaceae* The first description follows "*Clostridiaceae* which on meat medium produce after twenty days incubation under oil at 37° a reaction of pH 7.0 or a more acid reaction, the reaction being read after the culture has been boiled "

Later (p 550) the type genus is designated as *Rivoltillus* Heller

A list of fifteen genera were described and included, with designation of type species. The genera included are *Clostridium*, *Omelianskillus*, *Macintoshillus*, *Douglasillus*, *Henrillus*, *Flemmingillus*, *Vallorillus*,

Multi fermentans, *Hiblerillus*, *Welchillus*, *Stoddardillus*, *Rivoltillus*, *Arloingillus*, *Meyerillus* and *Novillus*

Clostridium. A generic name proposed by Prazmowski (1880, p 23) The origin of the name is given in the following words:

Um die Synonymik der Bakterien nicht mit einem neuen Worte zu bereichern, habe ich den von Trecul für eine Habitusform des Buttersäureferments zuerst angewendeten Namen *Clostridium* zur Bezeichnung meiner Gattung gewählt

Trecul's name (as a form genus or casual name) was given in 1865 (p 435)

Two species, *Clostridium butyricum* and *Cl. polymyxa* were described The genus was differentiated from *Bacillus* by the production of spores in spindle shaped cells In this article Prazmowski disposed of the prevalent idea that *Bacillus subtilis* causes butyric acid fermentation Zopf (1885, p 61) described the genus as containing organisms which resemble *Bacillus*, but producing the spores in characteristically broadened cells The genus was also recognized by Schroeter (1886, p 166) with the single species *C. butyricum*

Trevisan (1889, p 22) included *Clostridium* as the eighth genus of his tribe *Bacillae*, with eight species *Bacillus amylobacter* Van Tieghem 1877 is given as a synonym of *Clostridium butyricum* Prazmowski

De Toni and Trevisan (1889, p 1002) described nine species The generic description given is "Baculi dimorphi, plasmate uniformiter diffuse Sporae (endospores) microsomae in baculis specialibus intumescensibus oetae, ovoideae vel ellipsoideo-cylindraceae Among the species were *C. polymyxa* Piazm, *C. butyricum* Piazm, and *C. feveri* Trev (the cause of blackleg in cattle)

Hueppe (1891) and Sternberg (1892) used Zopf's diagnosis Macé (1897, p 468) concludes that the enlargement of the cell at the time of sporulation (the spindle shape) is not sufficiently distinctive for generic differentiation He cites the organism of symptomatic anthrax as one which shows great variation in the cell shape during sporulation

By most writers of the last two decades the genus *Clostridium* has been submerged in *Bacillus* Fischer (1895, p 143) divided the spindle shaped bacteria into four genera, defining *Clostridium* as the type in which the cells are motile by means of diffuse flagella Later (1897, 1903) he recognized two genera only of spindle shaped bacteria, *Paracloster* non-motile, and *Clostridium* with peritrichous flagella.

Erwin F. Smith (1905, p 174) rejects this generic name Orla-Jensen (1909) abandons this generic name, using *Butyribacillus* Vuillemin (1913, p 525) suggests that this genus be made a "*Genus conservandum*" with *Clostridium butyricum* Prazmowski as the type

Winslow *et al* (Committee Soc Am Bact, 1917, p 562) used this as a designation for all spore bearing anaerobes with the following description

Anaerobes Often parasitic Rods frequently enlarged at sporulation, producing clostridium or plectridium forms

The type species is *Clostridium butyricum* Prazmowski

Buchanan (1918, p 38) recognized both *Clostridium* and *Plectridium* for the spore bearing anaerobes The following generic description of *Clostridium* was given

Cells rod-shaped, straight or at least never spiral Frequently showing granules Endospores usually produced in cells showing some enlargement, usually the cells become spindle-shaped Anaerobic or microaerophilic Usually Gram-positive

The type species is *Clostridium butyricum* Prazmowski

Orla-Jensen (1921, p 273) suggests that the genus *Clostridium* as described by the Committee (Winslow *et al*) might well be split into two genera *Butyriclostridium* and *Putriclostridium*

Heller (1922, p 9) radically emended the generic definition, making it the first of the fifteen genera included in her family *Clostridioideae* *q v* The definition is

Clostridium Prazmowski 1880, emend Heller, *Clostridioideae* that do not liquefy gelatin Most species cause stormy fermentation of milk Highly saccharolytic, many even splitting pectins Produce considerable amounts of butyric acid as a split-product of carbohydrate fermentation May fix nitrogen Readily derive their nitrogen from inorganic nitrogen salts Large rods which are frequently polymorphic and form large ovoids and oval spores Frequently store up starch Form in deep agar large lenticular or modified lenticular colonies Common destroyers of plant tissue (not cellulose) in soil Used in the retting of flax to split pectins

Type species, *C amylobacter* van Tieghem as described by Winogradsky (1896) Probable synonyms *Amylobacter* Trécul, *Clostridium butyricum* Prazmowski, *Amylobakter* Gruber, *Granulobakter saccharo-butyricum* Beijerinck, *B saccharo-butyricus* von Kelcki, *Granulobacillus saccharobutyricus mobilis nonliquefaciens* Grassberger and Schattenfroh, *B amylobacter* von Hibler, *B amylobacter* Arthur Meyer and Bredemann Most of these are today incapable of accurate definition and several of them represent groups and not entities For this reason the type description chosen is that of Winogradsky instead of that of Prazmowski

Trécul (1865) gave the bacteria that contain starch the name of *Amylobacter*. He (1867) declared the *Amylobacter* to be heterogenetic, formed of minute particles that organise themselves into bacilli in decaying plant tissue. Van Tieghem (1877) named *Bacillus amylobacter* bacilli which contained amorphous starch during their growth. He believed such organisms to be the agents of cellulose destruction. Prazmowski (1880) described and figured *Clostridium butyricum*, which though it was in impure culture, was evidently of the type described above. Winogradsky (with Friebes) (1896) first defined a type that can be considered a species, he assigned no name to the organism. He declares that it does not split cellulose, but pectin. It ferments glucose, sucrose, lactose and starch in peptone media.

Apparently the type species of the genus is *Clostridium amylobacter* (Van Tieghem) Winogradsky unless it can be shown that the specific name *butyricum* antedates *amylobacter*.

Bergey *et al* (1923, p. 216) follows Winslow in including this genus as the second in the family *Bacillaceae*.

Clostrillum. A generic name proposed by Fischer (1895, p. 144) to include those rod-shaped bacteria which become spindle shaped when sporulating and which have a tuft of polar flagella. No species has been named, the generic designation is therefore a *nomen nudum* and invalid. It was abandoned later by Fischer himself (1897 and 1903).

Clostrinum. A generic name proposed by Fischer (1895, p. 144) to include those rod-shaped organisms which become spindle shaped when sporulating and which are motile by means of a single polar flagellum. No species were described, and the name is a *nomen nudum*. The name apparently was later abandoned by Fischer himself (1897 and 1903).

Clostrydium. Apparently a typographic error, a misprint for *Clostridium* found in the index to the second volume of Migula's *System der Bakterien* (1900, p. 1061). Enlows states (1920, p. 30) that this variant spelling has also been used by others in literature.

Coccaceae. A family of bacteria proposed under the name *Coccaeen* by Zopf (1884, p. 45) with the description "Sie besitzen nur die Coccen- und die durch Aneinanderreihung von Coccen entstehende Fadenform". The only genus recognized in this edition was *Leucostoc*. The next year (1885, p. 50) however, he included four genera, *Streptococcus*, *Micrococcus*, *Merismopedia* and *Sarcina*. The family was defined to comprise only cocci and filamentous forms consisting of rows of cocci, in which spore formation had not been recognized with certainty, and in which cell division occurred in one or more directions.

of space Schroeter (1886, p 143) recognized the family (with the spelling *Coccaceri*) making it the only family of his fourth order of bacteria (*Coccobacteria*) He states

Vegetative Zellen kuglig oder kurz elliptisch zu keiner Zeit ihrer Entwicklung zu Stäbchen oder Fadenformen auswachsend, immer unbewegt Die einzelnen Individuen im Substrat resp der von ihnen gebildeten Schleimmasse zerstreut oder in typischer Art zu bestimmt geformten Familien vereinigt Die einzelnen Individuen oder auch die Familien oft mit scharf begrenzten Gallert-hüllen umzogen—Sporen, soweit wie bekannt, aus einem vollständigen Zell—Individuum hervorgehend (Arthrosporen)

The genera recognized were increased to nine

Hansgirk (1888, p 266) listed this as a synonym of his *Mycococcaceae* Migula (1894, p 236) says,

Familie *Coccaceae* Zopf emend Mig Zellen im freien Zustande völlig kugel rund, Teilung nach ein, zwei oder drei Richtungen des Raumes, in dem sich jede Kugelzelle in Kugelhaften, Kugelquadraten oder Kugeloctanten teilt, die wieder zu Vollkugeln heranwachsen Endosporenbildung selten

The genera included were *Streptococcus*, *Micrococcus*, *Sarcina*, *Planococcus* and *Planosarcina*

Most recent writers on the classification of bacteria have used the name Among these may be cited Hueppe (1895, p 37), Lehmann and Neumann (1896, p 101), Fischer (1897, p 32), Macé (1897), Migula (1897, p 116, 1900, p 1, 1904, p 144), Chester (1897, p 62, 1901, p 55), A J Smith (1902, p 268), Schmidt and Weis (1902), Kendall (1902), Klocker (1903, p 330), Fischer (1903, p 59), E F Smith (1905, p 159), Flugge (1908), Ellis (1909, p 3), Frost (1911, p 56), Schneider (1912, p 22), Benecke (1912) The Winslows (1908, p 249) have emended somewhat the family diagnosis as follows

Cells, in their free condition, spherical, during division somewhat elliptical In the latter condition, division has already set in, although it may not be apparent Division in one, two or three planes without previous elongation of the cells If the cells remain in contact after division they are frequently flattened in the plane of division Motility is present only in a few forms Formation of endospores appears to be absent or very rare

These authors subdivide the family into the subfamilies *Paracoccaceae* and *Metacoccaceae* (qv)

Winslow *et al* (Committee Soc Am Bact, 1977, p 557) use the following description

Cells in their free conditions, spherical, during division somewhat elliptical. Division in one, two or three planes. If the cells remain in contact after division they are frequently flattened in the plane of division. Motility rare. Endospores absent. Metabolism complex, usually involving the utilization of amino-acids or carbohydrates.

Buchanan (1917, p. 603) emended the description as follows

Cells usually spherical, sometimes somewhat elongated just before division, or when occurring in pairs or groups cells may be somewhat flattened on sides in contact. Never containing granules of sulphur, nor with bacteriopurpurin. Many species develop a pigment. Some species are motile by means of flagella. Cells frequently remaining in groups after division, forming chains, regular packets or irregular masses. Spore production doubtful, having been reported for but one species. Pseudoplasmodium never produced. Growth energy not secured by the oxidation of ammonia or nitrites.

Three tribes were included, *Streptococceae*, *Micrococceae* and *Siderocapseeae*.

Castellani and Chalmers (1919, p. 924) accept the Committee's (Winslow *et al.*) emendation. They recognize the tribes *Streptococceae* and *Micrococceae*. They designate *Micrococcus* Hallier 1866 emendavit Cohn 1872 and Winslow and Rogers 1905, as the type genus.

Winslow *et al.* (1920, p. 205) slightly modified their original (1917) definition by including "forming chains, packets or irregular masses".

Bergey *et al.* (1923, p. 41) recognize *Coccaceae* as the second family of the order *Eubacteriales*.

It may be noted that objection might be raised to the fact that the family *Coccaceae* does not include any genus *Coccus*. Its use, however, is well established, it leads to no confusion, and there would seem to be no good reason for abandoning it.

Coccaceen. An aberrant form of spelling of the family *Coccaceae*. This was first used by Zopf (1884, p. 45), and later by other German writers.

Coccaceés. An aberrant form of spelling of *Coccaceae* used by certain French writers, as Macé (1897, p. 334).

Coccacei. An aberrant form of spelling the family name *Coccaceae*. This was used by Schroeter (1886, p. 143).

Coccobacillus. De Toni and Trevisan (1889, p. 966) state that Leube (1885) used the designation *Coccobacillus zymogenes*. They refer also to Cornil and Babes (1885). A careful study of Leube (1885, pp. 540-570) fails to show any such term. He describes a

Bacterium ureae and a *Micrococcus ureae* together with other unnamed organisms Cornil and Babes (1890, p 153) in the third edition of "Les Bactéries" figure a "*Coccobacillus zymogene* de Leube" The figure is of two distinct organisms, No 1 and No 3 of Leube Apparently De Toni and Trevisan are in error in ascribing a *Coccobacillus zymogenes* to Leube or to Cornil and Babes Bergey *et al* (1923, p. 265) state that *Coccobacillus ancidus* was used by Perroncito (1879, p 22). A careful examination of the original shows no such use

This name was also introduced independently by Gamaleia (1888, p 167) in the name *Coccobacillus ancidus* used for the organisms which caused bird septicemia (including fowl cholera) The name in this sense, however, is a synonym of the earlier name, *Pasteurella* Trevisan, which has priority.

Tissier (1900, p 70) described a *Cocco-Bacillus anaerobius perfoetans* from infant diarrhea

Tissier (1905, p 114) notes a *Coccobacillus perfoetans* from the intestines This organism is of two varieties, one which ferments glucose and saccharose, the other glucose, saccharose and lactose Inactive lactic acid is produced, and small quantities of butyric and valeric acids Does not form indol This generic name is rejected by Erwin F Smith (1905, p 174)

The name has also been used as a casual designation of the conjunctival organisms commonly known as diplobacilli This is the use made by Ruata (1909, p 630) who designates the organism as *Coccobacillus conjunctivae* A species, *Coccobacillus fusiformis* has also been named by Kawacki (1909, p 679)

Coccobacillus acridorum in a name applied by D'Herelle (1914, p 408) to an organism causing a disease in locusts This is Gram-negative motile bacillus, fermenting glucose, levulose, maltose and galactose It apparently belongs with the colon typhoid series of bacteria.

Cocco-Bacillus. A variant of *Coccobacillus qv* used by Tissier (1899, p 70) for the species *Cocco-Bacillus anaerobius perfoetans* As it was part of a trinomial, it may be regarded as invalid.

Coccobacteria. This generic name was introduced by Billroth (1874, p 1) for the species *Coccobacteria septica* This author conceived all bacteria to belong to one very pleomorphic species He describes it as a kind of plant which consists partly of spherical and partly of rod-shaped cells which may show considerable variation in size Each form may change into the other, although there is a certain

amount of constancy in that a coccus will for a long time form cocci and bacteria similarly produce bacteria. Both types of growth forms may produce a gelatinous capsule, forming cell plates (*Petalococcus* and *Petalobacteria*) or cell masses (*Gliacoccus*) or cocci may enlarge greatly and the contents divide forming a cell mass inside a capsule (*Ascococcus*). The rod-shaped forms may multiply in the same manner. Chains of cells (*Streptococcus* and *Streptobacteria*) may be formed. At certain times any of these cell types may show motility when not surrounded by capsular material and when not too large.

Billroth's conceptions of bacterial classification and growth have been disproved by time. The name *Coccobacteria* as a generic designation has never come into use.

The name was also proposed by Schroeter (1886, p. 143) for his fourth order of bacteria and as a synonym for *Sphaerobacteria* Cohn. It contained one family only *Coccaceae*. It included all bacteria in which the cells are spherical or short elliptical, always non-motile, frequently united in slimy masses or definite families, spores if any, arthrospores. The diagnosis given by Schroeter is as follows:

Vegetative Zellen kuglig oder kurz elliptisch, zu keiner Zeit ihrer Entwicklung zu Stäbchen- oder Fadenform auswachsend, immer unbewegt. Die einzelnen Individuen im Substrat resp. der von ihnen gebildeten Schleimmasse zerstreut oder in typischer Art zu bestimmt geformten Familien vereinigt. Die einzelnen Individuen oder auch die Familien oft mit scharf begrenzten Gallert-hüllen umzogen. Sporen, soweit sie bekannt, aus einem vollständigen Zell-Individuum hervorgehend (Arthrosporen).

The name has not been used by other authors as a bacterial order *Coccobacteriaceae*. A name proposed by Orla-Jensen (1921, p. 264) in the statement:

As to the family names of the bacteria, it will be convenient to let all of them end in *-Bacteriaceae*, by which it will be seen directly what is in question. If there are to be formed families of the cocci and spirilla, they must consequently be termed *Coccobacteriaceae* and *Spirillobacteriaceae* (or by the older name of Zopf, *Spirobacteriaceae*).

Coccobacterium. According to Trevisan (1887, p. 105) this term as a generic designation was first used by Rivolta. The former author states:

Rivolta sembra che miri a risuscitare l'appellativo *Coccobacterium* per un genere di cui non fa conoscere ne caratteri, ne limiti, di cui cita solo una seconda specie, il *Cocco-batterio* della tubercolosi, che per ogni altro e un *Bacillus*, e naturalmente non potrà essere seguito da veruno.

Rivolta (1888, p 1) says

"Nel N 1 Maggio 1887 nel giornale l'Allevatore descrissi un microbo del gatto, *coccobacterium felis* il quale, moltiplicandosi prodigiosamente nel sangue delgatto, fu causa di una ma lattia letale," also "Coltivazione Si semino il contagio nell'infuso di carne peptone gelatina, si moltiplico bene ed il liquido divenne torbido Si semino pure il virus nel brodo di carne di pecora per conservarlo "

Descrizione del virus Il virus venne studiato nel sangue di conigli e nelle culture ed e rappresentato da cocco-bacterini mobili in alto, in basso e qualcuno sull' asse trasversale, sono formati da due cellule minime o da tre talor come fuse insieme, quelli che si presentano all' osservatore per una delle loro estremita appaiono come micrococchi, sono un po' splendenti, della lunghezza di $0^{mm}00142$, altri sono piu corti ed alcuni pare che arrivino a $0^{mm}00285$

This organism was later named *Bacillus felis* (Riv) Trev by Trevisan (1889, p 14)

Klinger (1912, p 188) described a *Coccobacterium mucosum anaerobium* from pus. The organism is an anaerobic gas producing rod or coccus, growing only at blood heat and in serum medium The generic name is probably invalid

Schmidt and Weis (1902, p 10) again introduced the term for a generic name with the species *Coccobacterium prodigiosum* (Ehrenberg) S. & W This is a synonym of the older *Serratia* q v

Cocco-Bacterium. A generic name proposed by Rivolta (1888, p 1) for the species *Cocco-Bacterium felis* See *Coccobacterium*

Coccogenae A subfamily of the bacteria proposed by Trevisan (1885, p 92) to include all bacteria which have spherical cells, at least some stage in their life history The diagnosis is "Status unicus evolutionis coccus sistens" It is the equivalent of the family *Coccaceae* (q v) as commonly used

Later De Toni and Trevisan (1889, p 1035) include the following tribes *Ascococceae*, *Sarcineae*, *Streptococceae*, and *Micrococceae*

Coccogene. A variant of *Coccogenae* used by Trevisan (1889, p 26) for the third suborder of *Bacteriaceae* Four tribes, *Ascococceae*, *Sarcineae*, *Streptococceae*, and *Micrococceae* were included

Coccoglia. A synonym of *Zoogloea*, introduced by Billroth (1874, p 7) to include those cocci which occur embedded in gelatinous slime It apparently is synonymous with *Gliacoccus* It was not proposed as a generic name, but as a growth form of his pleomorphic *Cocco-bacteria septica* It has not been used by other authors

Coccomonas. A generic name suggested by Orla-Jensen (1921, p 268) in the following statement

For the Gram-negative, non-acid-forming or, at most, very slightly acid-forming cocci (among which must probably be reckoned the gonococci and the meningococci too, as well as Gram-negative streptococci, if such exist) we might simply use the generic name *Coccus*, or, if they should turn out to have *terminal* flagella, *Cocomonas*

Coccus. A name proposed by Billroth (1874, p 4) for a growth form of his pleomorphic *Coccobacteria septica* which is characterized by the production of spherical cells. In the form "coccus" it has been commonly accepted as a morphological term. It is not a generic designation in any strict sense. It is rejected by Erwin F. Smith (1905, p 174)

Coccothrichaceen. An "Abteilung" created by Lutz (1886, p 22) to include his genus *Coccothrix*. It has apparently not been used by other authors.

Coccothrix. A generic name proposed by Lutz (1886, p 22) for the organisms causing leprosy and tuberculosis. The diagnosis given by Lutz is as follows:

Kleine, runde, kokken-ähnliche Zellen, die sich ohne Mitbeteiligung der Zellmembran nur in einer Richtung teilen und daher einzeln oder in Reihen getroffen werden. Dieselben sind von den gedehnten, allmählich gallertig verquellenden, Zellhäuten überzogen, zwischen den Durchmesser an Grösse übertreffen. Durch Färbung der innersten Schichten der membranösen gallertigen Hülle entstehen Bilder, welche wie auf Fäden angereihte Perlen oder wie Stäbchen aussehen. Es finden sich ausserdem grössere zum Teil ovale und doppelt konturierte Zellen, teils frei, teils am Ende der Zellreihen. In dieses Genus ist vorderhand mit Sicherheit der Lepra- und Tuberkelpilz einzurichten.

It is true that Lutz misinterpreted some of his observations. To be accurate, his genus requires emendation in order to include the organisms of tuberculosis and leprosy. However, if the acid fast bacteria of which these are types are to be given generic recognition, *Coccothrix* would appear to be valid, with the type species either *Coccothrix tuberculosis* Lutz or *Coccothrix leprae* Lutz.

The name is rejected by Erwin F. Smith (1905, p 174).

Vuillemin (1913, p 527) notes this name, but believes that it resembles *Coccotrichum* Wallroth (1833, p 314) too closely. It would seem that these names are sufficiently unlike so that they would not be confused, and that both would be valid under a strict interpretation of the botanical code.

Synonyms of *Coccothrix* are *Sclerothrix* Metschnikoff and *Mycobacterium* Lehmann and Neumann. By many authors (as De Toni and Trevisan 1889, p 943) it is regarded as a synonym of *Bacillus*.

Coccus. This name with the spelling *Coccos* (*qv*) was first used by Billroth (1874, p 6) to designate the growth form of his *Coccobacteria septica* in which the cells assumed a spherical form. It was not used in a generic sense, however.

As a genus name *Coccus* has been accepted by several writers, but in most cases examined, they have been used in a trinomial combination and without any adequate generic diagnosis. Among such trinomial names may be mentioned *Coccus cumulus minor* Black (1886) (probably a *Sarcina* according to Miller 1892, p 78), *Coccus salivarius septicus* Biondi (1887, p 217) and *Coccus albus non-liquefaciens* Fick (1887).

Apparently the first use of *Coccus* in a binomial was by Fick (1887) for a *Coccus candicans* from the conjunctival sack. De Toni and Trevisan (1889, p 1078) rename this organism *Micrococcus Fickii* Trevisan.

It has also been used by Nissen (1889, p 487) for a *Coccus aquatilis*.

Billet (1890, p 24) accepts *Coccus* as a generic name with the statement "Formes arrondies, Coccus, élément isolé." With the spelling *Kokkus* this term has also been used by Eisenberg (1886) and other German writers. It is given as a synonym of *Micrococcus* by De Toni and Trevisan (1889, p 1076). Baumgarten (1890, p 127) says "1 Gatt Kokken, Bacterienarten umfassend, welche nur kugelege Vegetations formen bilden." The name is rejected by Erwin F. Smith (1905).

Orla-Jensen (1921, p 269) states

For the Gram-negative, non-acid-forming or, at most, very slightly acid-forming cocci (among which must probably be reckoned the gonococci and the meningococci too, as well as Gram-negative streptococci, if such exist) we might simply use the generic name *Coccus*, or, if they should turn out to have terminal flagella, *Cocomonas*.

The status of *Coccus* as a generic name is uncertain. Possibly the type *Coccus candicans* Fick cannot be identified.

Cohnia. A valid name given by Kunth (1850) to a genus of the family *Labiaceae*. The same name was given by Reichenbach, (1852) to a genus in the family *Orchidaceae*, now a synonym of *Cohniella* Pfitz.

This name was later proposed by Winter (1880, p 48) to include the bacteria which had been assigned to the algal genus *Clathrocystis* Henfrey. In this genus the cells are spherical, embedded in gelatin in a simple peripheral layer, forming hollow, round or irregular bladders or vesicles, which finally become reticulate. The diagnosis given follows:

Zellen rundlich, in einfacher peripherischer Schicht von gemeinsamer Gallerte umschlossen, hohle kuglige oder später unregelmässiger geformte Blasen oder Sacke bildend, die schliesslich netzformig durchbrochen sind Vermehrung der Zellen durch wiederholte Zweitheilung der Familien durch Furchung und Abgliederung von Tochterfamilien Die Cohn'sche Gattung *Clathrocystis* umfasst nach meiner Auffassung Algen und Pilze Da nun der Gattungsname zuerst für eine Alge (*Cl aeruginosa* Henfry) angewendet wurde, so ist es rathsam, ihn dieser zu lassen, und die zu den Pilzen gehörige Art zur Repräsentantin eines neuen Genus zu machen, dem ich, zu Ehren des um die Erforschung der Schizomyceten so hoch verdienten Professor Dr F Cohn in Breslau den Namen *Cohnna* gegeben habe

This genus is also recognized by Grove (1884, p 17). The name is rejected by Erwin F Smith (1905, p 174) and as used for bacteria is generally regarded as a synonym of *Lamprocystis*

The type species is *Cohnna roseo-persicina* (Cohn) Winter

Cohnistreptothrix. A genus of *thread bacteria* proposed by Pinoy (1913, p 929) to include those forms which differ from *Nocardia* by anaerobic or microaerophilic growth and by absence of conidia (gonidia)

The genus is accepted by Castellani and Chalmers (1919, p 251 and p 1063) They give the following statement (p 1063)

Definition *Nocardaceae* growing best anaerobically, but can grow aerobically, usually difficult to cultivate and do not produce arthrospores

Type Species *Cohnistreptothrix israeli* (Kruse 1896)

Historical In 1891, Wolff and Israel published a beautifully illustrated account of a streptothrix, which they had isolated from two cases of actinomycosis in man—viz, from the lungs and from a retromaxillary growth This organism was considered to differ from *N bovis*, in that it grew best anaerobically, that branching was absent, and that its injections into animals were regularly positive in their result These three characteristics induced Kruse, in 1896, to make a new species for it under the name *Streptothrix israeli* In 1911, for reasons already set forth, Pinoy founded a new genus *Cohnistreptothrix*, with Israel's organism as the type species, and therefore its name becomes *Cohnistreptothrix israeli* (Kruse, 1896)

These authors list a considerable number of species

Coleothrix. Veley and Velez (1898, p 154) according to an abstract (Kochs Jahresber, 9, 154, 1898) named a branched filamentous organism, sometimes appearing as cocci, *Coleothrix methystes* It is one of the important causes of faulty rum in the West Indies

Colibacterium. A generic name proposed by Orla-Jensen (1921, p 272) in the following statement

The genus *Bacterium* will undoubtedly dissolve into several genera, of which I may especially mention the *Colibacterium*, and *Aerogenesbacterium* The

reason why I am now inclined, in contrast to my earlier opinion, to consider the coli- and aerogenesbacteria as two different genera, is because they differ not only in morphological, but, as later researches have shown, also in biological respects. I have myself proved (1914) that the Aerogenesbacteria completely oxidize the carbohydrates when the nutrient matter offers a sufficient buffer effect, and they thus correspond with their name in forming more gas than do other bacteria, and Rogers, Clark and Davis (1914), have shown that in the gas developed by the Colibacteria there is proportionately more hydrogen than in that developed by the Aerogenesbacteria.

Conidiothrix. A name suggested by Benecke (1912, p. 489) as a generic designation. In a discussion of the work of Ellis on the iron bacteria, he says:

Es leuchtet ein, dass diese Form von der *Leptothrix ochracea*, wie sie oben beschrieben worden ist, ganz ausserordentlich abweicht. Ich mochte für den Fall, dass die Richtigkeit der Beschreibungen sich auch später hin bestätigen sollte, — was vorläufig recht zweifelhaft, erscheinen muss, den Namen *Conidiothrix* für sie vorschlagen.

The suggestion has apparently not been adopted by other authors. It may be regarded as a synonym of *Leptothrix*.

Coniothecium. According to Trevisan (1879, p. 144) the name *Coniothecium syphiliticum* was given by Hallier to an organism which he regarded as causing syphilis. Loeffler (1886, p. 143) quoted several paragraphs from an article by Hallier (1868, p. 327) in which he mentions this organism.

The organism evidently was so inadequately described that the name is probably invalid.

Cornilia. A genus described by Trevisan (1889, p. 21) as the seventh of the tribe *Bacillae*. The description reads: "Baculi con protoplasma uniformemente diffuso. Spore macrosome provenienti nelle parti mediane intumescanti, di regola non mai nelle estremità, di baculi normali immutati." Two subgenera are included, *Eucornilia* and *Pleurospora*. Fourteen species are listed, the first being *Cornilia alvei* (*Bacillus alvei* Flugge). De Toni and Trevisan (1889, p. 998) give the following diagnosis: "Baculi plasmati uniformiter diffuse foeti Sporae (endosporae macrosomae) in partibus medianis tumefactis baculorum normalium immutatis exorientes numquam (in baculis) apicales."

Two subgenera are included, *Eucornilia* with sixteen species and *Pleurospora* with two.

What species should be regarded as the type is uncertain. It is possible that it should be *Cornilia Pasteuri*, the name given by Trevisan to

the vibron septique of Pasteur, or to the first species *Cornelia alvei* (Cheshire and Cheyne) Trevisan

Altogether eighteen species are listed. The generic name is rejected by E. F. Smith (1905, p. 174) and by Vuillemin (1913, p. 521).

Corynebacterium A genus of bacteria founded upon the diphtheria bacillus by Lehmann and Neumann (1896, p. 350) with the following description:

Kulturen, durchaus den Charakter echter Bakterienkulturen tragend, weich, den Nährboden flach und locker aufliegend. Der Organismus färbt sich mit den gewöhnlichen Bakterienfärbemitteln gut. Mikroskopisch, Stäbchen, die an den Enden häufig keulig angeschwollen sind, aus verschiedenen farbigen Scheiben aufgebaut erscheinen und in manchen Kulturen durchweg eine unzweifelhafte echte dichotome Verzweigung zeigen.

The genus is accepted by Chester (1897, p. 63). It was rejected by Erwin F. Smith (1905, p. 174). This is suggested as a *Genus conservandum* by Vuillemin (1913, p. 527) and placed among the *Microsiphonés*.

Winslow *et al.* (Committee Soc. Am. Bact., 1917, p. 555 and 1920, p. 199) include this as the second genus of the family *Mycobacteriaceae* with the description:

Slender, often slightly curved, rods with tendency to club formation, branching cells occasionally seen in old cultures. Barred irregular staining. Not acid-fast. Gram-positive. Non-motile. Aerobic. No endospores. Some pathogenic species produce a powerful exotoxin. Characteristic snapping motion is exhibited when cells divide.

The type species is *Corynebacterium diphtheriae* (Loeffler) Lehmann and Neumann.

Buchanan (1918, p. 55) includes this as the tenth genus of the family *Bacteriaceae*. He gives the following description:

Rods which stain interruptedly (striped) with weak staining solutions. Not acid-fast. Clubbed, wedge-shaped and pointed rods frequent. Gram-positive. Non-motile. No spores. Aerobic.

The type of the genus is *Corynebacterium diphtheriae* Lehmann and Neumann.

Bergey *et al.* (1923, p. 380) recognize this as the second genus of the family *Mycobacteriaceae*, and use the diagnosis proposed by Winslow.

It probably may be regarded as a valid generic name if the diphtheria bacillus and related forms are to be accorded generic recognition.

Corynemonas. A genus of bacteria proposed by Orla-Jensen (1909, p. 344) to replace *Corynebacterium* Lehmann and Neumann (q v) It has as its type species *Corynemonas diphtheriae* and is placed in the family *Actinomycetes* as its second genus It is to be regarded as an invalid synonym of *Corynebacterium*

Corynethrix. A generic name used by Bongert (1901, p. 449) in the title of a paper, but not used by him in the article It is possible that the name has been used previously, but no earlier reference has been found

The organism *Corynethrix pseudotuberculosis murum* is described as a non-motile rod in young cultures on agar 0.5, 1 to 2 μ Older cultures show great numbers of involution forms On potatoes the cells may become 15 μ in length, and cultures three to four weeks old are almost *Actinomyces*-like The older rods stain irregular, and showing pronounced metachromatic granules In young cultures the cells are said to be gram negative, but when the granules appear they become gram positive Neisser's stain shows the metachromatic granules distinctly The organism is grouped quite definitely with the diphtheroid forms

The generic name occurs in a trinomial, and should probably be regarded as invalid, or at most as a synonym of *Corynebacterium*

Corynocladus. A genus of filamentous bacteria proposed by Leidy (1849, p. 250) The generic description and the description of the single species *Corynocladus radiatus* are as follows

Filaments hyaline, inarticulate, very compound, branches thicker than the trunk, clavate, without ramuli, growing from rounded or oval granular masses

Corynocladus radiatus Comatose, growing in very dense bunches, occasionally straggling, branches spreading, terminal ones very long, simple, clavate, $\frac{1}{16}$ inch long

Length $\frac{1}{16}$ inch diameter of trunk $\frac{1}{100}$ inch branches $\frac{1}{16}$ inch

Habitat Growing profusely in the ventriculus of *Passalus cornutus*, from the mucous membrane

This organism was again described and figured by Leidy (1851, p. 38)

Apparently the genus has not been recognized since the original description It may be included among the *genera incertae sedis*

Crenothrichaceae. The second family of the order *Desmobacteria* recognized by Hansgirk (1888, p. 229). He says

Diese bisher mit den Leptothricheen vereinigte Spaltpilzgruppe muss aus denselben Gründen wie die ihr unter den blaugrünen Algen (Schizophyceen)

entsprechende Familie der *Chamaesiphonaceae*, welche früher (noch von Rabenhorst) zu den Oscillariaceen, den Parallelförmigen der Leptothricheen gezählt wurde, von diesen getrennt werden

One genus only is included, *Crenothrix* Cohn

Crenothrix. A genus of filamentous bacteria proposed by Cohn (1872, p 130) for the species *Crenothrix polyspora* Cohn Cohn gives the following description

Trichomata plus minus stricta arcuata vel contorta in caespitulos libere nantes intricata libera vel alia alius affixa, in modum Oscillarium cylindrica elongate filiformia basi tenuissima sursum paulatim incrassata subulata vel subclavata divisione transversa succedanea articulata vaginata hyalina, cellularum plasmate homogeneo intus saepe cavo non granuloso, vagina tenerrima hyalina demum indurata nec non ferro intersuscepto flava

Sporangia terminalia apice trichomatum vagina intumescente elongato-claviformia gonidiis subglobosis numerosissimis densissime repleta, gonidia duplicis generis, saepissime in filis diversis formata

1 Microgonidia e serie cellularum divisione longitudinali et transversa succedanea multi-partitarum orta, rotundata et diaphragmatibus ruptis in sporangium terminale densissime congesta, demum ex apice vaginae erumpentia, in aqua motu lento circumvoluta secedentia vel in cumulos gelatinosos zoogloeis consumiles coacervata, cili, destituta globosa ovalia elliptica transverse plusminus constricta vel divisa, demum in trichomata evoluta

2 Macrogonidia, singula e cellulae contento toto indiviso, vel bivel quadri partito orta rotundata, ex apice vaginae vix inflatae erumpentia secedentia, motu forma microgonidius similis sed majora et minus numerosa, demum germinantia Spores? ex articulo trichomatis terminali elongata aucto formata plasmate denso repleta, quod e vagina erumpere et in trichoma Oscillariae forme evolvi videtur

Genus unter Lyngbyam et Chamaesiphonem intermedium nec von ad Bangias accedens Oscillariaceas cum Florideis connectit

The genus was included by Trevisan (1879, p 138) in his subtribe *Euribroniae* with the following description

Somatia cylindrica distincte articulata, valida, elongata, arcuata vel contorta, in caespitulos intricata, microgonidia e serie cellularum divisione longitudinali et transversa succedanea multipartitarum orta, globulosa, diaphragmatibus ruptis in pseudosporangium terminale densissime congesta, demum ex apice vaginae erumpentia et in somatia evoluta, foventia

The genus has been generally recognized by subsequent writers Winter (1880) and Grove (1884, p 54) used the following definition

Threads cylindrical, somewhat clavately thickened upwards, articulated, provided with a sheath Multiplication by means of the joints, which escape

from the sheath and grow into threads. Reproduction by spores, which are formed in the sheath by further subdivision of the joint-cells. The spores either grow directly into threads, or form by continued subdivision gelatinous colonies of roundish cells, which afterwards produce threads.

Other writers who recognized the genus were Van Tieghem (1884), Zopf (1885), Flugge (1886), Schroeter (1886, p 172), Hansgirk (1888, p 230), Trevisan (1889, p 10), De Toni and Trevisan (1889) and Hueppe (1885, 1891). Hansgirk (1891, p 313) discussed the synonymy of the genus, and included within it the genus *Phragmidiothrix* Engler. He divided the genus into two sections, as follows:

Section 1 *Phragmidiothrix* (Engler) nob. Plantae marinae 1 *Crenothrix-foetida* (Fior-Mazz.) nob. (*C. marina* Hansg.) cum aliis synonymis. Section 2 *Eucenothrix* nob. Plantae aquae dulcis 2 *Crenothrix Kuhniana* (Rbh.) Ciard. (*C. polyspora* Cohn) cum synonymis.

Ludwig (1892), Hueppe (1895, p 38), Migula (1895, p 40, 1897, p 47, 1900, p 1033, 1904, p 145), Fischer (1897, p 32, 1903, p 61), Chester (1897, p 64, 1901, p 372), Schmidt and Weis (1902, p 380), A. J. Smith (1902, p 272), Kendall (1902), Erwin F. Smith (1905, p 162), Ellis (1909, p 9), Orla-Jensen (1909, p 334), Molisch (1910), Frost (1911, p 59) and Heim (1911, p 254) have all described this genus.

Migula (1894, p 238) says:

Zellen zu unverzweigten Faden vereinigt, mit anfang nur nach einer Richtung vor sich gehender Teilung. Später teilen sich die Zellen nach allen drei Richtungen des Raumes, die Teilungsprodukte runden sich ab und werden zu Fortpflanzungszellen.

The later diagnosis of the genus by Migula gives the modern conception of the genus.

Cells united into unbranched filaments, attached, and gradually enlarging toward the free end, i. e., with a distinction between base and apex. Sheath rather thick. In iron waters the old and empty sheaths are permeated with iron oxide hydrate. Cells cylindric or flat discoidal. Multiplication by non-motile (mostly roundish) conidia, which arise from the vegetative cells by division and rounding off. For this purpose the cell of the thicker threads divide in three directions of space, those of the thinner threads only perpendicularly to the long axis of the thread. The conidia are discharged and germinate immediately, often on the sheath of the mother thread.

Molisch (1910, p 16) used practically the same description. He discusses *Crenothrix polyspora* Cohn at length. He states that there

has been confusion as to the correct specific name Migula (1900, p. 1033) notes that he examined the exsiccata of Rabenhorst's *Leptothrix Kuhniana* (Algens Sachsens No 284), but that it contained no organisms of the type of Cohn's *Crenothrix polyspora*, but a mixture of *Leptothrix ochracea* and *Gallionella ferruginea*. Zopf's use of the name *Crenothrix Kuhniana* he considers invalid

Buchanan (1918, p 305) included this genus as the third in the family *Chlamydo bacteriaceae* with the following description

Filaments unbranched, showing differentiation of base and tip, attached, usually thicker toward the tip Sheaths plainly visible, usually colorless, brownish from iron oxid in old filaments Cells cylindric to spherical Multiplication by non-motile, spherical, conidia, cells dividing in 3 planes to form conidia

The type (and only) species is *Crenothrix polyspora* Cohn

This description is also followed by Bergey *et al* (1923, p. 392)

Crenotrichaceae. A family named by Hansgurg (1890, p 20) for a group of *Mycophyceae*, including the genus *Crenothrix*

Crenothriceae. A tribe of the family *Trichogenae* proposed by De Toni and Trevisan (1889, p 925) with the following diagnosis "Sporae (arthrospora) in specialibus filamentorum partibus (pseudosporangis) obvenientes Filamenta simplicia, basi ab apice superior distincta, vaginata"

Crenothrix was the only genus included

Cristispira. A genus of spirochetes proposed by Gross (1910, p 44) to include those forms found in the crystalline style of mussels The body of the organism is circular in cross section, more or less spirally wound and possessing a longitudinal "comb" or "crest" which does not extend quite to the end This crest has been compared to the undulating membrane of the trypanosomes, though Gross does not consider them to be homologous organella Gross states that the organism is divided by cross walls into separate chambers or cells His generic diagnosis (p 89) reads "Spironemacea mit spiralig gebogenem Koerper und Crista Theilung durch einfache Durchschnurung oder Ausbildung einer Scheidewand, meist mit vorhergehender Incurvation" Two species are described *C veneris* and *C. interragationis*

This name is used as a subgeneric designation by Doflein (1911, p 356). It is recognized as a valid genus by Dobell (1911, p 536) and by Gonder (1914, p 678) The latter author summarizes the conclusions of various writers relative to the morphology of the organisms belonging to this genus.

Winslow *et al* (Committee Soc Am Bact, 1917, p 563) give this as the second genus under the heading of forms intermediate between bacteria and protozoa. The description is:

Cristispira Gross. Giant forms with undulating body and peculiar flattened ridge erroneously called an "undulating membrane" which runs the length of the body. Parasitic in molluscs.

The type species is *Cristispira balbranni* Certes, from the crystalline style of the oyster.

Buchanan (1918, p 544) makes this the fourth of the family *Spirochaetaceae*.

Spiral organisms known only from the crystalline style of mussels. The body of the organism is circular in cross section, more or less spirally wound and possessing a longitudinal comb or crest which does not extend quite to the tips.

The type species is *Cristispira veneris* Gross.

Bergey (1923, p 423) recognized this as the third genus of the family *Spirochaetaceae* with the following description: "Giant forms with undulating body and peculiar flattened ridge erroneously called an 'undulating membrane' which runs the length of the body. Parasitic in molluscs."

It would appear that *Cristispira* Gross constitutes a valid genus name. It is questionable, however, whether these forms are to be considered among the bacteria or the protozoa. Most authors include them among the latter. *C. veneris* Gross is the type.

Cromobacterium. A generic name proposed by Beigonzini (1881, p 153) for the species *Cr. violaceum*. The organism was first observed in a solution of egg white, producing a dark violet scum. Microscopic examination showed it to be made up of motile rods two to three times as long as broad. The solubility of the pigment in alcohol, but not in water, was noted. The original diagnosis reads:

Elementi cellulari cilindrici per lo piu isolati, 2 a 3 volte così lunghi come larghi, dotati di movimento oscillante, colorati in violetto da una sostanza speciale insolubile nell'acqua. Groschezza da 0.6 a 1 μ , lunghezza da 2 a 3 μ . Vegetano specialmente nelle soluz. d'albume d'ovo.

This organism has been variously named *Bacillus violaceus* and *Bacterium violaceum*. This genus with the spelling *Cromobacterium* does not appear to have been generally recognized by other writers, but with corrected spelling *Chromobacterium* *qv* it has been suggested as a generic name for the violet bacteria.

Cromococcus. A generic name used by Bergonzini (1881, p. 149) for a violet organism *Cromococcus violaceus* which he found in egg albumen. The original description is as follows

Osservai adunque che il liquido del bicchierino, che per evaporazione era ridotto a meno della metà, da bianco opalino era diventato giallo leggermente verdastro ed era coperto da una pellicola molta densa di un colore violetto scuro molto manifesto

Dapprima dubitai che fosse caduto qualche cosa accidentalmente entro il liquido, probabilmente dell'annilina, ma poi pensando che da due mesi non era stata toccata annilina nel mio studio, e che la colorazione violetta era limitata alla superficie del liquido, anzi ad una pellicola superficiale affatto identica a quelle di color bianco sporco che si presentano sopra tutti i liquidi di putrefazione, pensai ad una vegetazione batterica colorata, e probabilmente al *Cromococcus violaceus* l'unico batterio che presenta questa colorazione quantunque esso non sia mai stato osservato sull'albumina d'uovo

Ma dovetti ben presto ricredermi per due ragioni. Primo perché staccato un frammento di questa pellicola violetta e triturato nell'acqua essa non vi si scioglieva punto, mentre le zooglee del *Cromococcus violaceus* sono solubili nell'acqua. In secondo luogo poi perché l'esame microscopico invece di lasciarmi vedere la pellicola formata d'una zooglia di micrococchi me la mostrò composta di un numero straordinario di microbatteri violetti immobili riuniti a sciami (micoderma) i quali pigliavano un movimento distintissimo quando venivano disaggregati e potevano nuotare liberamente nel liquido

This generic name has apparently not been adopted by other writers. The description is scarcely sufficient to enable one to recognize the original species

The spelling was corrected to *Chromococcus* q v

Crypta. A generic name proposed by Salisbury (1868, p. 19) with the following description "Minute, transparent, highly refractive algoid filaments, which develop in living organic matter from spores"

Enlows (1920, p. 34) discusses the species as follows

C. syphilitica n. sp. A homogeneous filament, with extremities obtusely rounded. No transverse markings, except in early stage of development. Filaments either straight, coiled, or arranged in curves. They develop from spores which may be active or inactive in the connective tissue, and may be transplanted from one individual to another. Believed to be the cause of syphilis. *C. gonorrhoeae*. Spores very minute and well defined. Often in twos and sometimes in fours, undergoing the process of duplicative segmentation. They occur and develop rapidly in and among the parent cells of the mucous surfaces affected. In some instances the pus cells become filled with the spores of this "vegetation." The filamentous stage of this plant is frequently met with in and among the epithelial cells. In their embryonic stages a moniliform arrangement may be seen at times. In later and more advanced stages they are usually homogeneous

throughout their entire length. Occur singly or in little knots. Limited in its invasion to the epithelial tissue. Bergonzini (Lo Spallanzani, An 12, f 10 Modena, 1884) places these species among the Schizomycetes.

Cryptococcus. A genus of "algae" described by Kützting (1833, p 374), later used as a genus of fungi, to which occasionally bacteria have been referred, as *Cryptococcus xanthogenicus* Freiere (See Koch's Jahresber, 1, 346, 1885).

Cryptodesma. A genus of entophyte named by Leidy (1849, p 250) with the following description "Filaments ribbon-like, growing from attached granular masses. Consisting of a single cell, with a very delicate cell wall, and minute granular contents." One species *C. tenuis* was described from the ventriculus of *Passalus cornutus*. The specific description is as follows:

Filaments hyaline, compressed, attenuated at both extremities, growing in dense bunches from rounded granular masses. Cell wall very thin and delicate, granular contents of cell very fine and indistinct, measuring from $10\frac{1}{100}$ to $20\frac{1}{100}$ inch with a few coarser granules, and occasionally a few globules, measuring $1\frac{1}{100}$ inch. Length of filaments $2\frac{1}{100}$ to $7\frac{1}{100}$ inch greatest breadth $1\frac{1}{100}$ inch.

Leidy (1851, p 39) later questioned the ascription of name to this organism, doubting the entophytic nature of the form. The name is probably to be regarded as of doubtful validity.

Cylindrogloea. A genus of bacteria described by Perfiliev (1914, p 223). Enlows (1920, p 35) states:

Type species (monotypy) *C. bactifera*. A bacterium living symbiotically with *Chlorochromatium aggregatum* Lauterborn. A cylindrical, colorless zoogloeal mass, consisting of an axial filament formed by colorless cells, rectangular and numbering 20 to 35, measuring 0.7 to 0.8 μ by 2 to 4 μ , separated by interstices and surrounded by a mucilaginous envelope. In this outer mucilaginous zone the green bacteria lie, which Perfiliev says react in the same manner as the green form described by Lauterborn. The axial filament is not easily visible, because of the overlying *Chlorochromatium*.

Cystobacter. A genus of bacteria proposed by Schroeter (1886, p. 170) to include two species, *C. fuscus* and *C. erectus* Schroeter. The generic description reads:

Zellen in Form kurzen, dünner Stäbchen, in eine verbreitete Schleimmasse eingebettet, später in längeren Fäden zusammenhängend. Die Schleimmasse theilt sich in unregelmässige rundliche Klumpen, welche sich später mit einer festenhornartigen structurlosen Hülle umgeben.

The genus was accepted by De Toni and Trevisan (1889, p 1035) and Ludwig (1892) Thaxter (1892, p 394) concluded that this genus is a synonym of *Chondromyces* B and C (qv) and subsequent authors have accepted his findings. The generic name is therefore to be regarded as invalid.

Cystococcaceae. The first subfamily of the family *Mycococcaceae* proposed by Hansgirk (1888, p 266) He gave the following diagnosis "Zellen oder Zellfamilien von bestimmt umgrenzten Gallerthüllen (Cysten) umgeben" The following genera were included, *Leuconostoc*, *Ascococcus*, *Mycotheca*, *Leucocystis*, *Hyalococcus*

Cytobacter. A casual name suggested by Duclaux (1895, p 814) to designate certain bacteria which destroy the cellulose wall of the potato cell without attacking the starch, as distinguished from *Amylobacter* which destroys starch but not cellulose

Deazotonitrazobacterium. A name proposed by Ambroz (1913, p 10) for organisms which are capable of producing free nitrogen from nitrates One species is noted, *Deazotonitrazobacterium thermophilum* Ambroz

It would seem that a name of this length (12 syllables) would rather transcend good usage

Deazotonitranitrazobacterium. A name suggested by Ambroz (1913, p 10) to designate bacteria which set free atmospheric nitrogen from nitrates and nitrites See preceding

Deazotonitrazobacterium. A name suggested by Ambroz (1913, p 10) to include bacteria which can free elementary nitrogen from nitrites See preceding

Denitrobacillus One of several names applied by Ambroz (1913, p 6) to *Denitrobacterium thermophilum*, Ambroz Probably to be regarded as a casual designation

Denitrobacterium. A genus of bacteria proposed by Orla-Jensen (1909, p 314) to include those organisms which show active denitrifying powers, and have peritrichic flagella He notes the type to be the *Bacillus denitrificans* II of Burni and Stutzer, which has the power of reducing nitrates to nitrites and of freeing elementary nitrogen The genus is regarded as an intermediate type between the two orders *Cephalotrichinae* and *Peritrichinae*, and is placed tentatively as the first genus in the family *Acidobacteriaceae* of this latter order

Ambroz (1913, p 3) named a *Denitrobacterium thermophilum*

Enlows (1920, p 35) states "Type species (monotypy) *D. agile* (*Bacillus denitrificans agilis* Ampola and Garino 1896)."

If organisms of this type are to be accorded generic recognition, this genus name would appear to be valid

Denitromonas. A genus of bacteria proposed by Orla-Jensen (1909, p 314) for those organisms belonging to his family *Luminobacteriaceae* which show active denitrifying powers The organisms of this genus are usually fluorescent and differ from *Liquidomonas* in their lack of ability to liquefy gelatin The type species is *Denitromonas fluorescens* (*Bacterium fluorescens nonliquefaciens*) The name *Denitromonas* Orla-Jensen may be regarded as a synonym of *Pseudomonas* Migula and probably invalid

Enlows (1920, p 36) states *Bacillus denitrificans* Burri and Stutzer, to be the type

Desmobacteria. A tribe of bacteria proposed by Cohn (1872, p 173) to include the two genera *Bacillus* and *Vibrio* It comprises all filamentous bacteria, which consist of elongated cylindrical joints, which multiply by transverse fission and which occur in longer or shorter chains or filaments The cells are not constricted at the joints The original description reads

Alle Fadenbacterien bestehen aus verlängerten cylindrischen Gliedern, welche, wenn sie isolirt vorkommen, dem *Bact. Lineola* ähnlich sind, durch Quertheilung aber vermehrt, sich zu längeren oder kürzeren Ketten oder Faden aneinanderreihen, diese Faden sind jedoch nicht an den Gelenken eingeschnürt wie die Rosenkranzketten (Torulaform) der Oscillarien, an die sie sich zunächst anschliessen, sie werden in diesem Zustand als Leptothrixfaden bezeichnet Die Fadenbakterien bilden oft Schwärmern, nie aber, wie ich schon in einer Abhandlung von 1853 (Nov Act 1 c p 124) hervorgehoben, ZoogloeaGallert, doch wechseln auch bei ihnen bewegliche und unbewegliche Zustände, je nach der Anwesenheit oder dem Mangel an Sauerstoff, der Reaction des Mediums und andern noch unerforschten Bedingungen, gewisse Arten scheinen nie bewegt zu sein (*Bacteridium* Dav)

This grouping was also used by Schroeter (1886, p 170) as an order with the families *Leptotrichaceae* and *Cladothrichaceae*

Hansgirk (1888, p 229) also adopted this name for his first order of bacteria No description is given The three families *Cladothrichaceae*, *Crenothrichaceae* and *Leptothrichaceae* are included

With the spelling *Desmobacteriaceae*, this was used as a family name by Benecke (1912, p 188) This family he included in the order *Desmobacterinae*

Desmobacteriaceae. A name proposed by Benecke (1912, p 188) for the single family of the order *Desmobacterinae* This includes all

of the filamentous bacteria, usually grouped under *Trichobacteriaceae* and *Chlamydobacteriaceae*

Desmobacteries. The fourth order of bacteria, according to Maggi (1886, p. 173), including the genera *Bacillus*, *Actinobacter* and *Dispora*. No description is given

Desmobacterinae. An ordinal name used by Benecke (1912, p. 188) to include the single family *Desmobacteriaceae*

Desmobacterium. Included as a synonym of *Bacillus* by Klein (1885, p. 37) in the statement "*Bacillus* (*Desmobacterium* Cohn)"

Detoniella. A bacterial genus proposed by Detoni and Trevisan (1889, p. 929) with the following diagnosis

Filamenta cylindrica articulata, simplicia, basi ab apice superiori distincta, propter pulvinulum mucosum primitus affixa immobilia, serius libere natantia, lente oscillantia et in strata varie implexa Cocci constanter nulli Multiplicatio baculogonidis e vaginae apice egredientibus, primitus vivacissime mobilibus, cito immotis Arthrospora 4-5 in singulis articulis baculiformibus obvenientes OBS Genus a ceteris Kurthiensis arthrosporis in bucalis ortis coccorumque constanti defectu primo intuitu probe distinguendum

It is included in the tribe *Kurthiae* Trevisan (q v)

The genus includes three species of which the first described *Detoniella ochracea* (Roth) Trevisan may be regarded as the type The organisms included are those which are generally listed with *Leptothrix* and the genus may be regarded as a synonym of this term

Dialister. A name proposed by Bergey *et al* (1923, p. 271) for the second genus of the tribe *Hemophileae* with the description

Minute rod-shaped cells, occurring singly, in pairs and short chains Non-motile Strict parasites Growth occurs only under anaerobic conditions in media containing fresh, sterile tissue or ascitic fluid

The only species known is *Dialister pneumosintes* (Olitsky and Gates) comb nov

Diaschizae A subdivision of the group *Schizomycetes* coordinate with *Paraschizae* proposed by Jennings (1899, p. 315) to include all bacteria which multiply vegetatively by transverse division

Dicoccia. A genus of bacteria proposed by Trevisan (1889, p. 26) De Toni and Trevisan (1889, p. 1034) give the following diagnosis "Baculi Klebsiellae, capsulis, inclusi, plasmate polaris-diblastico foeti, spora, ignota Ob. Ut Pasteurella a Bacilio, hoc genus a Klebsiella distat" This genus is included in the tribe *Klebsielleae* Trevisan (q v) and the subtribe *Euklebsielleae* The cells resemble *Klebsiella* Trevisan (q v) except that the cells show bipolar characters similar

to *Pasteurella* Trevisan (q v) One species only is given *Dicoccia glossophila* Trevisan This is the "Bakterien menschlicher Mundsecrete, zweite Art" of Flugge (1886, p 258). It is probable that this organism should be included with the hemorrhagic septicemia forms although the specific determination is not practicable from Flugge's description This would make *Dicoccia* a synonym of *Pasteurella* The genus is rejected by Erwin F Smith (1905, p 174)

Didymohelix. A generic name proposed by Griffith (1853, p 438) for the organism termed *Gallionella ferruginea* by Ehrenberg, *Gloeotila* by Kützing and *Meloseira ochracea* by Ralfs He points out that each of these authors included this species with others of the genus among the diatoms This organism, as it is not a diatom, cannot be included either in *Gallionella* or *Gloeotila* He notes that the organism usually has the appearance of two intertwined filaments, and proposes the name *Didymohelix* For discussion of the synonymy, see *Gallionella*

Buchanan (1918, p 304) recognized this as the second genus of the family *Chlamydobacteriaceae* with the following diagnosis

Filament twisted, simple, or two filaments, twisted together Young cell colorless, later yellow brown to rust red through deposition of iron Simple filaments show no division into cells, even when iron is removed with acid and stain applied Sheath not demonstrable

The type species is *Didymohelix ferruginea* (Ehr) Griffith

Diffusionsbacillus. A casual name used by Beijerinck (1893, p 830) Enlows states "A species closely related to *B perlibratus* Probably did not intend to use it as a generic name"

Diplectridium. A genus of bacteria created by Fischer (1895, p 2) to include those rod-shaped organisms having diffuse flagella which become clavate when spores are produced, one spore being formed in each end of the cell No species apparently have been described within this genus and the name itself has been abandoned by Fischer in later publications (1903) This generic name is to be regarded as a *nomen nudum* and invalid It is rejected by Erwin F Smith (1905, p 174)

Enlows (1920, p 147) states the type to be "*B solmsi* Klein"

Diplobacillus. Maggi (1886, p 174) used this term in designating a growth form of a *Bacillus* Apparently it was first used as a generic name by Weichselbaum (1887, p 212) in the combination *Diplobacillus brevis endocarditidis* to designate a short rod which occurred in pairs and was apparently associated with a case of endocarditis The cells

characteristically show polar staining, only the youngest cells stain uniformly. The organism grew well at room temperature on ordinary media. The name as used, in the form of a trinomial, is non-valid. De Toni and Trevisan (1889, p. 997) included this organism in the genus *Pasteurella*, naming it *Pasteurella Weichselbaumii*.

Morax (1896, p. 337) described a diplobacillus associated with conjunctivitis, the same organism being discussed in greater detail later by Axenfeld (1897, p. 1). This organism has been commonly known since as the Morax-Axenfeld bacillus, or in some cases as *Diplobacillus Morax*. The latter designation is binomial in form, and perhaps might validate the generic name *Diplobacillus* for this group of organisms. The species is termed *Bacterium duplex* by Lehmann and Neumann and *Bacillus lacunatus* by Muir and Ritchie.

Petit (1898) described a related organism from a case of hypopyonkeratitis, as a "Diplobacille liquefiante." McNab (1904, p. 64) discussed this organism under the name "*Diplobacillus liquefaciens*" Petit, and the earlier organism as *Diplobacillus Morax-Axenfeld*.

Diplobacteria. This name was applied by Billroth (1874, p. 16) to a growth form of his *Coccobacteria septica* in which the rod-shaped organisms occurred in pairs. The name has never been used as a generic designation. It is rejected by Erwin F. Smith (1905, p. 174).

Diplobacterium. A casual name used by Haase (1887, p. 347) in the phrase, "Diplobacterium der Brustseuche." The name was also used as a morphological term by Billet (1890). It has not been used as a generic designation.

Diplococcus. A growth form of *Coccobacteria septica* named by Billroth (1874, p. 5) to indicate a grouping of cocci in which the organisms occurred normally in pairs.

The spelling has been changed to *Diplococcus* by later writers and used as a generic designation.

Diplococcus. The name *Diplococcus* was first used by Billroth (1874, p. 5) to designate a growth form of his *Coccobacteria septica* in which the spherical cells occurred in pairs. It was not used in a strict generic sense.

As a generic name *Diplococcus* came into use without any definite characterization. Bumm (1885, p. 16) used the term diplococcus, though not in a generic sense, to designate the characteristic grouping of the gonococcus and related forms. In no case was it used in a strict binomial designation. Fluegge (1886, p. 183) follows Bumm in the

use of the name *Diplococcus albicans tardissimus*, but as this is a trinomial it is invalid. In the same year Weichselbaum (1886, p. 506) gave to the pneumococcus the name *Diplococcus pneumoniae*, a valid binomial. This seems to be the first correct designation of a species of the genus, and the pneumococcus should therefore be regarded as the type. Several other species were assigned to this genus in the next two years, but of these only four appear to have binomial form and therefore valid. These are *Diplococcus roseus* and *D. subflavus* described by Bumm (1887), *D. luteus* Adametz (1887, p. 9) and *D. coryzae* Hajek (1888).

As the designation of a genus, *Diplococcus* has rarely found its way into formal classifications, although diplococcus is in common use as a casual designation. For example, Sternberg (1892, p. 17) under the heading of *Diplococcus* makes the following statement:

Association in pairs is common to all of the micrococci, inasmuch as they multiply by binary division. When such association has rather a permanent character it is customary to speak of the microorganism as a diplococcus, but we doubt the propriety of recognizing this mode of association as a generic character.

Winslow and Rogers (1905, p. 669) have included *Diplococcus* as one of the valid genera in their studies of the classification of the *Coccaceae*. In further contributions on the same subject the genus was accurately defined as follows (1908, p. 250):

Strict parasites, not growing, or growing very poorly, on artificial media. Cells normally in pairs, surrounded by a capsule. Fermentative powers high, most strains forming acid in dextrose, lactose, saccharose and inulin. Hemolytic power generally lacking. Characteristic group serum reactions.

They assign to this genus the pneumococcus, the gonococcus, the meningococcus and the coccus of catarrh.

If the Gram-positive cocci of which the pneumococcus is a type are to be grouped together into a genus, it would appear that the generic name *Diplococcus* is valid, if, however, the gram negative cocci of which the gonococcus is the type are to be included in the same genus, *Diplococcus* becomes a synonym of an earlier genus, *Neisseria* Trevisan. This latter genus was founded upon the gonococcus by Trevisan in 1885 (p. 16). Article 46 of the botanical code states "When two or more groups of the same nature are united, the name of the oldest is retained." Inasmuch as the genus *Diplococcus* as used by the Winslows contains also the type of the genus *Neisseria* Trevisan, and since the latter name has priority, *Neisseria* should replace *Diplococcus*.

Buchanan (1817, p. 612) included this as the third genus of the *Streptococceae* with the diagnosis

Parasites growing poorly or not at all on artificial media Cells usually in pairs, capsulated Fermentative powers high, most strains forming acid in dextrose, lactose, saccharose and inulin The cells are Gram-positive

The type species is *Diplococcus pneumoniae* Weichselbaum

Winslow *et al* (Committee Soc Am Bact, 1920, p. 206) describe the genus as follows

Parasites, growing poorly, or not at all, on artificial media Cells usually in pairs of somewhat elongated cells, often capsulated, sometimes in chains Gram positive Fermentative powers high, most strains forming acid in glucose, lactose, sucrose and inulin

Type species, *D pneumoniae* Weichselbaum

Castellani and Chalmers (1919, p 927) diagnose the genus as follows

Streptococceae usually in encapsulated pairs, forming acid in glucose, lactose, saccharose, and inulin Haemolysis usually absent Characteristic group serum reactions Soluble in bile salts Undergo autolysis in cultures and in emulsions of physiological saline

Bergey *et al* (1923, p 44) include this as the second genus of the *Streptococceae* and follow the Committee's description

It would appear that *Diplococcus* Weichselbaum is a valid generic name for the pneumococcus and the related forms, if they are worthy of generic recognition If the gonococcus and its relatives are to be included in the same genus, the name *Neisseria* Trevisan should be used

Diplokokken. Baumgarten (1890 p 127) states "1 Untergattung *Diplokokken*, die Zellen bleiben nach der Theilung zu je zweien im Zusammenhang "

Diplokokkus. An aberrant form of spelling of *Diplococcus* used by certain German authors, as Baumgarten (1890, p 127), Liebs (1887 p 321)

Diplopneumococcus. A casual name used by Krokiewicz (1904)

Diplostreptococcus. A generic name used by Lingelsheim (1912, p 494) in a discussion of "Brustseuche der Pferde" He states, "Als erreger hat man langere Zeit, den von Schutz (1887, p 27) zuerst gefundenen *Diplostreptococcus pleuropneumoniae* der sich vielfach in den Krankheitherden findet angesehen " This reference to Schutz as an authority for this generic name is erroneous as a careful search of the

text fails to show, by this author, any use of the name, *Diplostreptococcus pleuropneumoniae*

It is possible that it has been used previously to the article by Lingelshelm. It does not appear that this organism is sufficiently distinct from other streptococci to warrant separate generic recognition

This name has also been used as a casual designation for the lactic organism of the fermented milk "Yoghurt," by Luerssen and Kuhn (1908, p 241) It was not used, however, as a generic name

This designation has also been used by Bettencourt and others for streptococci showing diploid elements when taken from cerebrospinal fluid (1904, p 55) Perrone (1905, p 371) mentions a *Diplostreptococcus intestinalis* of Tavel

Diplovibrio. A name given by Maggi (1886, p. 176) to a growth form of *Vibrio*

It has also been used by Billet (1890, p 23)

A casual name

Discomices. Apparently a typographical error for *Discomyces* *q v* Migula (1900, p 116)

Discomyces. A generic name proposed by Rivolta (1878, p 208) to replace *Actinomyces* Harz, which he regarded as inept Rivolta and Micellone (1879, p 145) used the name *Discomyces equi* to designate the organism causing so-called "botryomycosis" of the horse Later Rivolta (1884, p 198) also described a species *Discomyces pleuriticus* from the dog This genus was at first regarded by Rivolta as identical with *Actinomyces* Harz, but later both genera were recognized as is evident from the following statement by Rivolta (1884, p 188)

Discomyces Il micelio di questo fungo é fatto da grandicellule otricellari granulose

Differisce dall'actinomyces bovis Harz per la grandezza e disposizione degli elementi che lo costituiscono e per la struttura più semplice

Esaminando questa specie, il Prof Harz si convincera come non possa venir confusa con l'actinomyces bovis perché ha nulla di raggianto E fatto da grandi cellule e non da filamenti Le si conviene adunque il nome di discomyces equi

Si può pertanto fondare un gruppo di discomiceti patogeni comprendente le seguenti specie

I *L'actinomyces bovis* Harz Sin Corpuscoli o cespugli discordi Rivolta, colle due varietà Actinomice granuloso e Actinomice degli ascessi

II *Il Discomyces pleuriticus canis familiaris*

III *Il Discomyces equi* Rivolta e Micellone

Discomyces Rivolta is listed as a synonym of the genus *Nocardia* Trevisan, by De Toni and Trevisan (1889, p 927) although the two

species of Rivolta are not included here, but under *Bollingera* Trevisan (p 1039) as *Bollingera equi* Trev and *B Vacchetae* Trevisan

This generic name has generally been disregarded because of confusion with the *Discomycetes* among the sac bearing fungi, because of the priority of other generic names and because of the lack of a careful characterization of the organisms The species described as *Discomyces equi* is generally regarded as a capsulated coccus and not a filamentous form related to *Actinomyces* at all

Blanchard (1895) came to the conclusion that the older generic name *Actinomyces* was invalid because of its prior use to designate an unrelated fungus, and proposed the use of *Discomyces* to replace it. In this proposal he has been followed by Brumpt (1910, p 842)

Discomyces is definitely rejected by Smith (1905, p. 174) and by Vuillemin (1913, p 527)

Merrill and Wade (1919, p 64) review the status of this name very completely, and conclude that it is the valid designation They say

Vuillemin, and more recently Chalmers and Christopherson, in advocating *Nocardia* as the valid generic name, hold that Rivolta's use of *Discomyces* was trivial and without botanical significance We do not agree with this argument, which is clearly refuted by Rivolta's original paper Here he distinctly proposes *Discomyces bovis* as the name for the organism called *Actinomyces bovis* by Harz in a manner that must be acknowledged as valid from the viewpoint of botany, even though it is not in conventional form and was advocated on irrelevant, inadequate grounds Therefore, it is in no sense a "medical genus," as Vuillemin asserts The fact that subsequently Rivolta erroneously referred other organisms to this genus has no bearing on the case His original application of it was to the organism of Bollinger and Harz alone, which is, therefore, the type of the genus Nor does the fact that to propitiate Harz, Rivolta later agreed to accept *Actinomyces* affect the question As Blanchard pointed out, a name once introduced is no longer the property of its originator to withdraw or modify at will

Finally, to argue, as do Chalmers and Christopherson, that *Discomycetaceae* a group name, invalidates *Discomyces*, as a generic name in the connection in which Rivolta used it, on the ground that the type genus of Fries's *Discomycetaceae*, published in 1836, should bear the designation *Discomyces*, indicates an erroneous conception of the principles of nomenclature and priority in technical names, a family name such as *Discomycetaceae* cannot invalidate a similar specific name This generic name was new with Rivolta, and there is no valid objection to its adoption in taxonomy

For reasons for the invalidity of *Discomyces* see *Actinomyces*

In the present work *Discomyces* is regarded as a synonym of *Actinomyces*

Dispora. A generic name proposed by Kern (1882, p 135) in a preliminary note for a rod-shaped organism found by him in kefir grains and regarded by him as the organism causing the development of the lactic acid In a later contribution (1882, p 141) he described the organism in greater detail

The organism in the kefir grains occurs in the form of zoogloeal masses associated with a yeast The vegetative cells when in active growth are described as 3.2 to 8μ in length by 0.8μ in diameter Under the influence of acids, of high temperatures, etc., the cells become *Leptothrix*-like and filamentous These filaments ordinarily contain spores The vegetative cells may be motile, when they are supplied at one end with a slender flagellum The spores are spherical, and are always found in pairs, each cell producing two spores, one in each end The spores are not large enough to cause any distortion of the cell Within the cells the spores are apparently 0.8μ in diameter, when free 1μ In the *Leptothrix* forms the spores appear to be in rows Careful examination, however, shows them to occur always in pairs within separate cells In germination an exo- and endosporium are recognizable.

Kern states that in the vegetative condition this organism resembles *Bacillus subtilis* Cohn, but that the presence of two spores in a cell is an adequate generic differential character

The generic name *Dispora* is accepted and used by Maggi (1886, p. 176)

De Toni and Trevisan (1889, p 1015) included this form in the genus *Pacinia* under the name *P. caucasica* (Kern) Trevisan Cornil and Babes (1890, p 183) recognized the genus by describing the species *D. caucasica* The genus has in general not been recognized by recent writers Either Kern worked with an organism which has not been since recognized, or he grossly misinterpreted what he saw Certainly the organism now regarded as the most important in Kefir manufacture does not conform to the description given by Kern

If the organism commonly called *Bacillus bulgaricus*, *B. caucasicus* or *B. lebanus* is to be made the type of a genus, it is important to determine the status of the generic name *Dispora* If Kern misinterpreted what he saw, and in reality named the high-acid bacillus of Kefir *Dispora*, the name, in spite of its lack of appropriateness would appear to be valid If, however, as seems probable, Kern was describing not the causal organism of Kefir but some other, the name can be rejected for this purpose Inasmuch as the organism described by Kern has

not been later recognized, the name *Dispora* may be regarded as a *nomen nudum*, or possibly as a synonym of *Bacillus* Cohn. It is rejected by Erwin F. Smith (1905, p. 174) and by Vuillemin (1913, p. 520).

Douglasillus. The fourth genus of the subfamily *Clostridioideae* proposed by Hellei (1922, p. 14) with the following description

Clostridioideae that do not liquefy gelatin. They produce little gas in meat medium. They may not clot milk. Gram-negative bacilli which are frequently fusiform, and may show peculiar involution forms. Spherical spores are formed in the bacilli, the rods are usually widened by the spores so that their sides are not parallel. Young spores may not be truly spherical. Occasionally found in moulds. Probably incapable of invading tissue.

Type species *sphenoides* (*Bacillus sphenoides* Douglas, Fleming and Colebrook), as described by the Committee (p. 43) the type which ferments glucose, maltose, galactose, lactose, salicin, mannitol, sucrose, dextrin and starch.

Drepanospira. A generic name used by Boris de Petschenko (1911, p. 249) to designate a species of bacterium, *Drepanospira* Muller, found to be parasitic within the bodies of paramoecia, and having a relatively complex life cycle. Enlows (1902, p. 39) is apparently in error in writing *Drepanospora*. She states

Order *Eubacteria*, fam. *Spirillaceae* occupying an intermediate place between *Spirosoma* Migula and *Microspira* Schroter. Cells with 2 spirals, one end pointed, the other somewhat rounded. No cilia or flagella. Helicoidal motion. "Pas de division cellulaire". Endospores. Regular spherical colonies formed by the individuals in certain stages of development. Measure 7μ long by 0.75μ wide. Cell membrane visible on the living cell. In the vegetative state protoplasm composed of an anterior, smaller, strongly refractive part, and a larger posterior or dull part. The anterior portion he considers to be nuclear.

Dysenteroides. The fourth genus of the tribe *Ebertheae* proposed by Castellani and Chalmers (1919, p. 934). The description (p. 935) is as follows:

Definition. *Ebertheae* fermenting glucose and lactose partially with the production of acid, but no gas. Milk not clotted.

Type Species. *Dysenteroides metadysentericus* (Castellani, 1917).

Ebertheae. The ninth tribe of the family *Bacillaceae* proposed by Castellani and Chalmers (1919, p. 932). The description given follows:

Bacillaceae growing well on ordinary laboratory media, not forming endospores, aerobes, and often facultative anaerobes, without fluorescence, pigment

formation, or gelatine liquefaction, without polar staining, Gram-negative, without a capsule

Type Genus *Eberthus* Castellani and Chalmers, 1918

Ten genera are included, *Alcaligenes*, *Eberthus*, *Shigella*, *Dysenteroides*, *Lankoides*, *Salmonella*, *Balkanella*, *Wesenbergus*, *Enteroides* and *Escherichia*

Eberthella. A subgenus of *Bacterium* proposed by Buchanan (1918, p 53) to include those forms "Not producing gas from any of the carbohydrates, acid may or may not be formed" The type species is designated as *Bacterium (Eberthella) typhi* Flugge.

Bergey *et al* (1923, p 221) erected *Eberthella* into a genus, erroneously ascribing the name to Castellani and Chalmers Possibly this is a misprint for the generic name *Eberthus* used by Castellani and Chalmers for the same group It is included as the fifth genus of the tribe *Bactereae* with the following description

Motile or non-motile rods, occurring in the intestinal canal of man, usually in different forms of enteric inflammation Attack a number of carbohydrates with the formation of acid but no gas Do not form acetyl-methyl-carbinol

The type species is *Eberthella typhi* (Eberth-Gaffky) Castellani and Chalmers

Eberthus. The second genus of the tribe *Ebertheae* named by Castellani and Chalmers (1919, p 934) The description (p 936) follows

Bacillaceae motile, partially fermenting glucose with the production of acid and no gas Lactose not fermented Milk not clotted

Type Species *Eberthus typhosus* (Zopf, 1885)

Later (1920, p 604) four species are described

Eccrina A generic name proposed by Leidy (1851, p 35) for an entophyte with "Characters same as *Enterobryus*, except that it divided into numerous cells at the free extremity" Two species were described, *Eccrina longa* from the mucous membrane of the intestine of *Polydesmus virginensis*, and *Eccrina moniliforma* from *Polydesmus granulatus*

Leidy (1851, p 29) later described the genus as follows

Thallus attached, consisting of a single very long tubular cell, filled with granules and globules, producing at its free extremity a succession of numerous globular or oblong cells, and growing at the other end from a relatively short, cylindrical, amorphous coriaceous pedicle, commencing with a discoidal surface of attachment

The description given for *E. longa* is as follows:

Thallus long, filiform, colorless, or brownish, transparent, cylindrical, usually not holding a constant relation of breadth to the length, forming a simple curve or single spiral turn, and then proceeding in a straight line or gently flexuose curve to the free extremity. *Pedicle* very short, columnar, expanded at the base. Principal cell very long, uniformly cylindrical. Secondary cells in various stages of development from ten to thirty in number, oblong, or short cylindrical, with obtusely rounded extremities when completed.

Whole length from 2 to 7 lines. Length of pedicle $\frac{1}{1000}$ to $\frac{1}{1500}$ of an inch. Breadth of principal cell, from $\frac{2}{1000}$ to the $\frac{1}{1000}$ of an inch. Length of secondary cells, from the $\frac{1}{125}$ to the $\frac{1}{3125}$ of an inch.

Habitation. Parasitic, growing in profusion from the mucous membrane of the posterior part of the intestinal canal of *Polydesmus virginicus*.

The generic name is probably valid, but it is questionable whether the organism belongs with the bacteria.

Enterbacterium. A casual designation used by Kuttner (1895, p. 263) for his *Pyobacterium fischeri*.

Enterococcus. A casual name used by German writers as a designation for pus coccus (Rosenbach 1884, p. 23).

Encapsulateae. The name proposed by Castellani and Chalmers (1919, p. 932) for the ninth tribe of the family *Bacillaceae*. The description (1919, p. 934) is as follows:

Bacillaceae growing well on ordinary laboratory media, without endospores, neither fluorescent nor chromogenic aerobes, not liquefying gelatin, possessing capsules in animal tissues.

Type Genus *Encapsulatus* Castellani and Chalmers, 1918.

One genus only, *Encapsulatus*, is included.

Bergey *et al.* (1923, p. 237) recognize *Encapsulateae* as the seventh tribe of the family *Bacteriaceae* with the following description:

Short rods, somewhat plump with rounded ends, mostly occurring singly. Encapsulated. Non-motile. Gram-negative. Ferment a number of carbohydrates with the formation of acid and gas. Encountered principally in the respiratory tract of man. Aerobic, growing well on ordinary culture media.

This probably should yield priority to *Klebsiellae qv*.

Encapsulatus. The only genus of the tribe *Encapsulateae* proposed by Castellani and Chalmers (1919, p. 934) with the description:

Encapsulateae with the tribal characters.

Type Species *Encapsulatus pneumoniae* (Friedlaender, 1883).

Later (1920, p 602) the following statement is made

Ce genre est actuellement le seul de la tribu, et il embrasse l'ancien groupe des bacilles encapsulés qui ont été passés en revue par Fricke en 1896, Clairmone en 1902, Perkins en 1904, Abel et Kallwacks en 1912 et Fitzgerald en 1914

Les espèces sont ordinairement des organismes courts, non mobiles, gram-négatifs, encapsulés, pléomorphiques, qui font fermenter le glucose et le lactose, mais en ce qui concerne ce dernier sucre, ils peuvent produire seulement de l'acidité quoiqu'ils habituellement ils forment de l'acide et du gaz

Bergey *et al* recognize this as the single genus of the tribe *Encapsulateae*

Apparently *Encapulatus* must yield priority to *Klebsiella* *q v* for both are based upon the same type species

Endobacterium. Lehmann and Neumann (1896 p 103) in a discussion of *Bacillus* state "Den uns besonders geeignet erscheinenden Namen 'Endobacterium' unterdrucken wir, um nicht weitere neue Namen zu benutzen"

Endostreptococcus. A name proposed by Hueppe (1885) for organisms having spherical cells in chains, producing endospores, zoogloea not formed. Apparently this has never been coupled with a species name, and is a "*nomen nudum*"

Endostreptokokkus. A German variant of *Endostreptococcus* *q v* used by Hueppe (1891, p 33)

Enterobrus. A genus of organisms described by Leidy (1849, p 225) to include certain parasitic organisms found in the intestines of milliped *Julus* and the beetle *Passalus*. Leidy regarded it as a genus of *Conservaceae*. The original description reads

Simple, attached, isolated filaments consisting of a long cylindrical cell (containing protoplasm, granules, and large translucent globules enveloped in a primordial utricle) with a distinct coriaceous peduncle or stripe of attachment, and at length producing at the free extremity one or two, rarely three, shorter cylindrical cells, (filled with the same matter as the parent cell)

The first species described was *Enterobrus elegans* from the intestine of *Julus marginatus*. Later (1848, p 249) he described *Enterobrus spiralis* from *Julus pusillus* and *Enterobrus attenuatus* from *Passalus cornutus*

Later Leidy published revised descriptions with drawings of these various species (1851, p 19-20). The revised description reads as follows

Thallus attached, consisting of a single, very long, tubular cell, filled with granules and globules, producing at its free extremity one, usually two, rarely

three, shorter tubular cells, and growing at the other end from a relatively short, cylindroid, amorphous, coriaceous pedicle, commencing with a discoidal surface of attachment

There is doubt as to whether the organism described and figured so carefully by Leidy should be placed with the bacteria. In many of its characteristics it seems to approach certain forms of the fungi

It should be noted that in the last mentioned publication of Leidy (1851, p 19-20) the generic name is spelled *Enterobryus*

Enterobryus. A spelling of the generic name *Enterobryus* used by Leidy (1851, p. 19)

Enterococcus A casual name given by various writers to cocci found in the intestinal tract. Apparently it was first used by Thiercelin (1898) in the form *Enterocoque*. Tissier (1900, p 65) pointed out that his organism is apparently identical with the *Micrococcus ovalis* of Escherich (1886). Tissier at this time gave a fairly complete description of the organism. It was also studied by Hollig (1904, p. 11). According to Tissier (1905, p 114) *Enterococcus* produces an acid fermentation in glucose, saccharose and lactose. Rougentzoff (1914, p 648) discusses an *Enterococcus saccharomyces* from the intestines of the rabbit

Enterocoque A casual name used by Thiercelin (1899) and Tissier (1899, p 65) for *Enterococcus* (q v) the *Micrococcus ovalis* of Escherich

Enteroides A name given to the ninth genus of the tribe *Ebertheae* by Castellani and Chalmers (1919, p 935) with the description

Ebertheae which ferment glucose and lactose completely with the production of acid and gas. Milk not clotted

Type Species *Enteroides entericus* (Castellani, 1907)

These authors name five species (1920, p 618)

Erebonema. Cohn (1883, p 193) gives the following account of this genus

Romer (Deutschlands Algen, S 70) hat in den tiefen Gruben bei Clausthal im Harz Schleimüberzüge gefunden, die von locker verwebten, gegliederten, farblosen Faden durchzogen waren. Er beschrieb sie unter dem Namen *Erebonema* und unterschied nach der Dicke und Gruppierung der Faden zwei Arten, *E. hercynicum* und *E. divaricatum*. F. T. Kützing, welcher Romer'sche Exemplare untersuchte, hat (Species Algarum 1849, S 157) die Gattung dahin charakterisiert: deutlich gegliederte, sehr locker verwebte, farblose, verzweigte Faden, zwischen einer schleimigallertartigen, aus sehr kleinen schleimigen Kugeln zusammengesetzter Grundmasse nistend, Glieder hohl, schlaff, letzte Verzweigungen erweitert.

ert Kützing hat also die Faden in der Gallert für die Hauptsache, die Coccen, die er gleichwohl schon auffand und in seinen *Tabulae phycologicae* abbildete, für nebensächlich gehalten

Professor Ferdinand Cohn untersuchte im Jahre 1857 eine rothlichgelbe Gallert, welche in einer 60 Lachter tiefen Grube bei Volpersdorf gefunden worden war Ihr Aussehen war ganz so, wie ich es bei dem Kellerschleim gefunden, sie hing auch hier von dem Gestein als Zapfen herab Professor Cohn erklärte den Schleim für identisch mit *Erebonema hercynicum* Kg und fand in ihm Faden, ähnlich wie sie Kützing beschreibt, nur waren sie nicht, wie dieser angiebt, an den Enden angeschwollen, sie stimmten also mit den Faden des Kellerschleims überein Zwischen den Faden, welche Cohn für eine *Leptomitrus*artige Alge erklärte, lagerten in structurloser Gallert zahllose ovalstabchenförmige, farblose Körperchen dicht nebeneinander, so dass sie den Bau einer *Palmella* darboten Er stellte es bereits als zweifelhaft hin, ob die Faden wirklich zum Begriff der Species gehörten oder nur zufällig in der *Palmella* als Parasiten existiren, und neigte sich der letzteren Ansicht zu

Schroeter (1883) termed the organism *Leucocystis cellaris*

This latter name is also accepted by De Toni and Tievisan (1889, p 1041) The name *Erebonema* is rejected by Edwin F Smith (1905, p 174)

Ermengillus. A genus of anaerobic bacteria proposed by Heller (1922, p 28) It is the twenty-fourth genus of the family *Clostridiaceae*, and is included in the subfamily *Putrificoraceae* The description is

Putrificoraceae that produce a yellowish coloration in meat medium, and later blacken and digest it They are more highly proteolytic than the organisms of the foregoing genera They ferment various sugars Gram-positive rods which form sub-terminal oval spores Their colonies in deep agar, are, when discrete, lenticular or kidney shaped, and may show tufted smooth or woolly polar projections and infrequently fine loose woolly radiations Fairly common in soil, grow readily in vegetable material and on meat Form a characteristic and powerful neuro-toxin

Type species *E. botulinus* (*Bacillus botulinus* von Ermengen) as described by K F Meyer and co-workers in a future paper

Erwineae A tribe named by Winslow *et al* (Committee, Soc Am. Bact, 1920, p 209) as the second in the family *Bacteriaceae*, with the following description "Plant pathogens Growth usually whitish, often slimy Indol generally not produced Acid usually formed in certain carbohydrate media, but as a rule no gas"

The single genus is *Erwinia*

Bergey *et al* (1923, p 168) have used the variant spelling *Erwiniae*. Probably the spelling should be *Erwineae*

Erwinia. A generic name proposed by Winslow *et al.* (Committee Soc Am Bact, 1917, p 560) with the following description: "Plant pathogens Growth usually whitish, often slimy Indol generally not produced Acid usually formed in certain carbohydrate media, but as a rule no gas."

Later (1920, p 209) it was included as the single genus of the tribe *Erwiniae*, and the type species designated as *Erwinia amylovora* (Burrill) Committee

Bergey *et al* (1923, p 168) follow the committee in placing the genus in the tribe "*Erwiniae*" with the following description "Motile rods, possessing peritrichous flagella The rods are white and few species form pigment"

Erwiniae. A variant spelling of *Erwiniae* or *Erwiniae* q v used by Bergey *et al* (1923, p 168). It is included as the fourth tribe of the family *Bacteriaceae* with the following description

Plant pathogens Growth usually whitish, often slimy Indol generally not produced Acids usually formed in carbohydrate media, forming acid or acid and gas Motile, or non-motile Gram-negative

Erysipelococcus. A casual designation apparently first used by Bender (1888, p 10) for Erysipelkokken of Fehleisen Enlows states

This name has been ascribed by numerous authors to Fehleisen In Sitz d Phys-Med Gesellsch z Wurzburg 1881-1885, p 126 (1881) and p 9 (1883) he described a micrococcus "der Mikrokoccus des Erysipels" In his second paper he uses in the title of the paper the word "Erysipelkokken," but in the text he uses "mikrokocken"

Erysipelococcus. Enlows (1920, p 41) states this is found in Lipp. Med Dict Phil, 1920, p 322, syn *Streptococcus erysipelatis*

Erysipelothrix A generic name proposed by Rosenbach (1909, p 367) to include the causal organisms of swine erysipelas, human erysipeloid and mouse sepsis As characters differentiating this genus from *Bacillus* are noted the formation of long filaments and the regular development of branches, the thickening of the filaments with the characteristic granules The species named are *Erysipelothrix porci*, *E erysipeloides* and *E murisepticus*

Buchanan (1918, p 55) included this as the ninth genus of the family *Bacteriaceae* with the following description.

Rod-shaped organisms with a tendency to the formation of long filaments which may show branching The filaments may also thicken and show charac-

teristic granules No spores Non-motile Gram-positive Does not produce acid Microaerophilic Usually parasitic

The type species is *Erysipelothrix rhusiopathiae* (*E. porci*), the causal organism of swine erysipelas

Winslow *et al* (Committee Soc Am Bact, 1920, p 198) emended the diagnosis as follows

Rod-shaped organisms with a tendency to the formation of long filaments which may show branching The filaments may also thicken and show characteristic granules No spores Non-motile Gram-positive Do not produce acid Microaerophilic Usually parasitic

The type species is *Erysipelothrix rhusiopathiae* (*Bacillus rhusiopathiae* *suis* Kitt 1893, *Mycobacterium rhusiopathiae* Chester 1901, *Erysipelothrix porci* Rosenbach 1909), the causal organism of swine erysipelas

Bergey *et al* (1923, p 371) follow the Committee and include the genus as the fourth in the family *Actinomycetaceae*

Erythrobacillus. A generic name used by Fortineau (1905, p 104) for an organism, *Erythrobacillus pyosepticus* This species is described as a motile coccobacillus, but *without flagella*, without spores, staining readily and gram negative In culture it grew well at 37° but showed pigment development best at 19° to 22° In bouillon a distinct clouding developed, and a delicate rose colored film On agar a vermillion mucus-like layer formed Serum and gelatin were both liquefied In Uschinsky solution there was growth but no pigment production Ammonia, hydrogen sulphid and indol were formed, nitrates reduced and sugar fermented feebly Pigment was produced only in presence of air It was soluble in water, alcohol, slightly soluble in chloroform and insoluble in ether, carbon disulphid and benzine, and decolorized by alkalis

Winslow *et al* (Committee Soc Am Bact, 1920, p 209) recognizes this genus with the following diagnosis

Small aerobic bacteria, producing a red or pink pigment, usually a lipochrome Gram stain variable It is possible that related yellow and orange chromogens should be included here as well

Type species, *Erythrobacillus prodigiosus* (Ehrenberg) Committee

It is probable that the committee's designation of *E. prodigiosus* is not valid, as this species was not mentioned by Fortineau The type is *E. pyosepticus* Enlows (1920, p 41) states

Type species (monotypy) *E. pyosepticus* A motile "coco-bacille," with no spores, flagella, Gram-negative, grows at 37°C, red pigment forms best at 19°

to 22°C Pigments soluble in water, the alcohols, slightly so in chloroform insoluble in ether, carbon bisulphide, benzine Pathogenic for the guinea pig etc Isolated from the chemise of a patient at the Hotel-Dieu, Nantes

It is not probable that this name is to be regarded as a valid generic name, although it is possible that it might be used for the red bacilli if they are to be accorded generic recognition providing *Serratia* and *Zooglossina* are invalid It is rejected by Erwin F Smith (1905, p 174)

Erythroconis. A generic name proposed by Oersted (1840, p 555) for an organism described by him from the Danish coast The description given is "Character genericus. Massa pulveracea parum mucosa ex corpusculis quadratis rigidis fragilibus per quaterna aggregates constans Genus e familia Diatomerarum analogon Palmellae et Tetrasporae"

His species, *Erythroconis littoralis* is described by Rabenhorst (1865, p 57) as follows "Thallo mucoso-grumoso indefinito olivascente familiae minimis e cellulis cubico-polygonis 32 compositis, cytoplasmate aerugineo-viridi Diam cell circ 1/300" = 0.00029" This later author regards *Erythroconis* as a subgenus of *Merismopedia* Meyen among the blue green algae, with the characterization "Species in aqua natantes vel in limo nascentes," and including with it all those species not belonging to "*Sarcina medicorum*"

Winter (1884, p 50) uses the name *Sarcina littoralis* with the following description

Zellen kuglig oder vor der Theilung oval, 1 2 μ , selten 2 oder mehr μ in Durchmesser, zu 4, 6, 8, etc., zu Familien verbunden, die ihrerseits grossere Colonien (bis zu 64 Tetraden in einer Colonie) bilden Plasma farblos, aber in jeder Zelle 1-4 rothe Schwefelkörner In faulenden Meerwasser

De Toni and Trevisan (1889, p 1049) regarded this generic name as a synonym of *Lamprospedia* Schroeter, though this latter author did not include Oersted's species

Migula (1900, p 1045) regards *Erythroconis littoralis* Oerstedt as a doubtful synonym of *Thiopodia rosea* Winogradsky.

The genus is rejected as invalid by Erwin F Smith (1905, p 174)

The type species is *Erythroconis littoralis* Oersted.

Escherichia. A name given to the tenth genus of the tribe *Ebertheae* by Castellani and Chalmers (1919, p 935) The description (p 941) follows

Ebertheae which ferment glucose and lactose completely, milk clotted
Type Species *Escherichia coli* (Escherich, 1886)

Some 16 species (1920, p 619) are listed.

Bergey *et al* (1923, p 194) includes *Escherichia* as the first genus of the tribe *Bactereae* with the following description

Motile or non-motile rods, commonly occurring in the intestinal canal of normal animals Attack numerous carbohydrates forming acid and frequently acid and gas Do not produce acetyl-methyl-carbinol

The type species is *Escherichia coli* (Escherich) Castellani and Chalmers

Eu-Ascococceae. A name given by De Toni and Trevisan (1889, p 1035) to a subtribe of the tribe *Ascococceae*, with the following diagnosis "Subtribus I *Eu-Ascococceae* Cocci in muco matricali segregati, in familias cystidibus universalibus obductas cumulati Cystides speciales nullae" It contains seven genera *Lamprocystis* Schroter, *Ascococcus* Cohn, *Bollingeria* Trevisan, *Leucocystis* Schroter, *Cenomesia* Trevisan, *Thiothece* Winogradsky, *Thiocystis* Winogradsky

This name does not appear to have been used by other investigators

Eubacilleae. The third subtribe of the tribe *Bacilleae* proposed by De Toni and Trevisan (1889, p 939) with the following diagnosis "Baculi (divisione semper in unam directionem transversalem) liberi, numquam spiraliter torti, Endosporae" Ten genera are noted *Mantegazzia* Trevisan, *Bacillus* Cohn, *Pasteurella* Trevisan, *Cornelia* Trevisan, *Clostridium* Prazmowsky, *Vibrio* Zopf, *Spirillum* Ehrenberg, *Spiromonas* Perty, *Pacinia* Trevisan, *Bacterium* Ehrenberg

Eubacillus. This name was first used by Hansgirk (1888, p 264) for the first section of the genus *Bacillus* with the following diagnosis "Zellen mit farblosen, feinkörnigen Plasma, ohne grossere Schwefelkörnchen, zu farblosen oder grau-bis gelblich-braunlichen Schleimmassen (Zoogloeen) vereinigt" *B subtilis* was included in this section

A genus of green bacteria proposed by Dangeard (1891, p 151) was also given this name This genus was created to include those organisms, rod-shaped in form, or variable length, with hyaline protoplasm, containing no granules and with a small amount of chlorophyll in the protoplasm, spore bearing filament may be simple or branched, the green color more pronounced in the swellings, spores

formed by the contraction of the protoplasm of the swellings, spores refringent, grouped or isolated, separated from each other by septa. The genus is divided into two sections, the first containing the type species, *Eubacillus multisporus* Dangeard. This organism is made up of filaments which are simple, long, and slender and of a faint green color. The spore bearing filaments are likewise quite long, producing the numerous spores isolated or grouped in pairs, threes, fours, separated by septa. Spores are elliptical in shape and of a decided green color with one or two refringent granules. Length of spores 5 to 8 μ , width 3 μ . Habitat: Developing in cultures of algae from fresh water.

In the second section, the author includes five species of bacteria which have been described by Kline, all of which are green in color.

Lehmann and Neumann (1896, p. 11) question the ascription of this genus to the bacteria. Migula (1897, p. 94) concludes that the organisms described under this name by Dangeard do not belong to the bacteria, both because the spore bearing filaments of his *Eubacillus multisporus* were branched at times and because of the formation of spores in chains which is not characteristic of bacteria.

Meyer (1912, p. 2) uses the name *Eubacillus* for the third section of the genus *Bacillus* with the statement "Enthalt alle ubrigen Spezies der Gattung, auch die, fur welche Sporangienbildung nicht gesehen wurde."

Doubt as to the generic validity of *Eubacillus* Dangeard is raised by the previous use of the term by Hansgirk, in practically a subgeneric sense. There is also some reasonable doubt as to whether Dangeard's organism belongs with the bacteria.

Eu-Bacillus. A subgenus of *Bacillus* proposed by Buchanan (1918, p. 38). A variant of *Eubacillus* Hansgirk. The description follows:

Motile, usually by means of peritrichous flagella. Spores not barrel-shaped without longitudinal ridges. Other characters those of the genus.

The type species of the subgenus is *Bacillus subtilis* Cohn.

Eubacteria. The fifth order of bacteria recognized by Schroter (1886, p. 154) with the following diagnosis:

Vegetative Zellen kurzer oder langer stabchenformig, ruhend oder bewegt frei oder in Schleim eingebettet. Häufig wachsen sie in lange Faden aus, welche aus einzelnen Gliedern bestehen, die bei der Sporenbildung wieder durch schärfere Theilung deutlich werden, sie zeigen keine deutliche gemeinsame Scheiden. Sporen meist einzeln innerhalb der Zellen gebildet und deren Raum nicht ganz

ausfüllend, seltener aus dem Inhalte eines Gliedes derart gebildet, dass die Spore diesem an Grösse gleichkommt oder dasselbe übertrifft

There are included in this order the single family *Bacteriaceae* Zopf, and nine genera, namely *Bacterium* Ehrenberg, *Chromatium* Perty, *Bacillus* Cohn, *Clostridium* Prazmowsky, *Spirillum* Ehrenberg, *Spirochaete* Ehrenberg, *Microspira* Schroter, *Myconostoc* Cohn, *Cystobacter* Schroter

Hansgurg (1888, p 264) used this as the designation of his second order of *Bacteria*, containing the two families *Bacteriaceae* and *Myconostocaceae*. The name was adopted by Migula (1900, p 1) for his first order of *Bacteria* with the following diagnosis "Zellen ohne Centralkörper, Schwefel und Bacteriopurpurin, farblos oder schwach gefärbt, chlorophyllgrün" He divides the order into four families *Coccaceae*, *Bacteriaceae*, *Spirillaceae*, *Chlamydobacteriaceae*

This name has been used in a similar sense by Erwin F Smith (1905, p 159) Frost (1911, p 56), and Meyer (1912, p 1)

Eubacteriaceae. A spelling of the order *Eubacteria* used by A J Smith (1902, p 270) This name is not in the correct form for the designation of an order

Eubacteriales. An ordinal name apparently first proposed by Buchanan (1917, p 347) for the first order of the *Schizomycetes* Following is the description used

This order includes the true bacteria, those forms which are least differentiated and least specialized The cells are usually minute, spherical, rod-shaped or spiral, not typically producing true filaments although the cells may occur in chains or other groups The cells do not have a well-organized or differentiated nucleus They may be motile by means of flagella, or non-motile, never notably flexuous Multiplication by transverse fission, never by longitudinal Some forms, particularly rod-shaped types, produce endospores, but never conidia Branching of cells occurs in a few forms which intergrade with the *Actinomyce-tales* Neither sulphur granules nor bacteriopurpurin are present, though the cells may be pigmented Chlorophyll is absent (with the possible exception of one genus) The cells may be united into gelatinous masses, but never form motile pseudoplasmodia, nor develop a highly specialized cyst-producing fruiting stage

Four families were included *Coccaceae*, *Bacteriaceae*, *Spirillaceae* and *Nitrobacteriaceae*

The description was emended by Winslow *et al* (1917, p 550) as follows

The order *Eubacteriales* includes the forms usually termed the true bacteria, that is, those forms which are considered least differentiated and least specialized

The cell metabolism is not primarily bound up with hydrogen sulphide or other sulfur compounds, the cells in consequence containing neither sulfur granules nor bacterio-purpurin. The cells apparently do not possess a well-organized or well-differentiated nucleus. The cells are usually minute and spherical, rod-shaped or spiral in shape, in most genera not producing true filaments, the filaments when formed not sheathed, and frequently branching, thus being differentiated from the iron bacteria. The cells may occur singly, in chains or other groupings. The cells may be motile by means of flagella, or non-motile, they are never *notably* flexuous. Cell multiplication occurs always by transverse, never by longitudinal fission. Some genera produce endospores, particularly the rod-shaped types.

Chlorophyll is absent, though the cells may be pigmented. The cells may be united into gelatinous masses, but never form motile pseudo-plasmidia nor develop a highly specialized cyst-producing fruiting stage, such as is characteristic of the *Myxobacterales*.

Castellani and Chalmers (1919, p. 924) use the simple definition, "*Schizomycetes* which contain neither sulphur nor bacterio-purpurin." They include five families, *Coccaceae*, *Bacillaceae*, *Spirillaceae*, *Chlamydo bacteriaceae* and *Mycobacteriaceae*.

In the final revision Winslow *et al.* (1920, p. 200) emended the description as follows:

The order *Eubacterales* includes the forms usually termed the true bacteria that is those forms which are considered least differentiated and least specialized. The cell metabolism is not primarily bound up with hydrogen sulphide or other sulfur compounds, the cells in consequence containing neither sulphur granules nor bacterio-purpurin. The cells apparently do not possess a well-organized or well-differentiated nucleus. These organisms are usually minute and spherical rod-shaped or spiral in most genera not producing true filaments and rarely branching. The cells may occur singly in chains or other groupings. They may be motile by means of flagella or non-motile, but are never *notably* flexuous. Cell multiplication occurs always by transverse never by longitudinal fission. Some genera produce endospores particularly the rod-shaped types. Conidia not observed. Chlorophyll is absent, though the cells may be pigmented. The cells may be united into gelatinous masses, but never form motile pseudoplasmodia nor develop a highly specialized cyst-producing fruiting stage such as is characteristic of the *Myxobacterales*.

Heller (1921, p. 449) makes this the first order of the class "*Eubacterieae*," with the definition "*Eubacterieae* whose cells are never in sheathed filaments. Conidia not observed. Free iron sulphur, or bacteriopurpurin never present. Multiplication always occurs by transverse fission. (Committee)"

Eubacterieae. A subtribe of bacteria proposed by Trevisan (1879, p. 136) with the following diagnosis: "Cellulae liberae vel in statu quiescente in colonias mucosas indeterminatas consociatae." It is

the first subtribe of the tribe *Bacterieae*, and contains the genera *Micrococcus* and *Bacterium*

Heller (1921, p 449) has proposed *Eubacterieae* as a class name coordinate with *Myrobacterieae* in the phylum *Bacteria*. This spelling is not in correct form for a class name

Eubeggiatoa. The first section of the genus *Beggiatoa* recognized by Hansgurg (1888, p 263). He gives the following description "Faden farblos, zu kreideweissen bis graugelblichen schleimigen Massen vereinigt". Two species are listed, *Beggiatoa alba* (Vauch) Trev and *B. arachnoideae* (Ag.) Rbh var *uncinata* Hansg.

Eucoccaceae. The second subfamily of the family *Mycococcaceae* described by Hansgurg (1888, p 267) with the following description "Zellen oder Zellfamilien ohne bestimmt begrenzte, Gallerthüllen, nackt oder von einer gemeinsamen Gallerte umgeben". One genus only, *Micrococcus*, is included

Eucoccus. A subgenus of the genus *Micrococcus* Cohn created by Migula (1895, p 16) to include those forms in which the cell contents are colorless and free from sulphur granules. The second section is *Thiopolycoccus* Winogradsky.

Eu-Cornilia. A subgenus of the genus *Cornilia* Trevisan established by De Toni and Trevisan (1889, p 998) to include those species which are saprogenic and anaerobic. A variant of *Eucornilia*

Eucornilia. The name applied by Trevisan (1889, p 21) to the first subgenus of *Cornilia*. The second subgenus is *Pleurospora*. Altogether twelve species are listed, of which the first is *Cornilia alvei* Trev.

Eucrenothrix. A section of the genus *Crenothrix* suggested by Hansgurg (1891, p 314) to include the fresh water forms of the genus with *C. Kuhniana* as the type. The other section for marine forms he terms *Phragmidiothrix*.

Eu-Klebsiellaeae. A subtribe of the tribe *Klebsiellaeae* created by De Toni and Trevisan (1889, p 1028) with the following diagnosis "Baculi recti vel curvi, numquam spiraliter torti". It contains three genera *Winogradskya* Trevisan, *Klebsiella* Trevisan, *Dicoccia* Trevisan.

Eumantegazzaea. A subgenus of the genus *Mantegazzaea* created by De Toni and Trevisan (1889, p 942) to include three species which are colorless and do not contain granules of sulfur.

Eumicrospira. The name applied by Meyer (1912, p. 4) to the second section of the genus *Microspira*. He included in the section *Rhodospira* "Bakteriopurpurin enthaltende Species" and in *Eumicrospira* "Alle anderen Species".

Eumonas. Enlows (1920, p 42) states

Eumonas Diesing, 1850 *Systema Helminthum* v 1, Vindobonae, 1850, p 22 Subgenus of *Monas* Muller and Ehrenberg Diesing says *Monas* is synonymous with *Bodo* and *Bacterium* Ehrenberg *Eūmonas* Animalculasolitaria libera, corpus ecaudatum, subglobosum, ovatum, v obconicum, hyalinum v coloratum, haud mutabile, divisione spontanea simplici bipartitum, v indivisum Os terminale truncatum, limbo ciliatum v nudum Flagellum nudum ocellus nullus Includes 20 species *Monas* (*Eumonas*) *crepusculum* Ehrenberg *M* (*Eumonas*) *bicolor* Ehrenberg, *M* (*Eūmonas*) *ochracea* Ehrenberg, *M* (*Eumonas*) *hyalina* Ehrenberg, etc

Eu-Pacina. A subgenus of the genus *Pacima* Trevisan named by De Toni and Trevisan (1889, p 1019) to include those species which conform to the following diagnosis "Baculi recti vel raro levissime curvuli, interdum in filamenta recta vel subrecta consociati, Arthrosporaē magnae, quadriplooctuplo et ultra diametri transversalis baculorum latiores" Eleven species are included It is interesting to note that *Pacima cholerae-asiaticae*, the organism upon which the genus *Pacima* was founded is not included in this subgenus, in other words, the type species of the genus is apparently regarded by De Toni and Trevisan (1889, p 1018) as not typical of the genus A variant of *Eupacima*

Eupacina. The first subgenus of *Pacima* (Trevisan 1889, p 23) The description is "Filamenti normali di *Bacillus*" Nine species are included, but *Pacima cholerae-asiaticae* is placed in the second subgenus *Pseudospira*

Euplanococcus A subgenus of the *Planococcus* created by Migula (1895, p 19) to include those forms in which the cell contents are colorless and free from sulphur granules The second subgenus is *Thiopedia* Winogradsky

Euplanosarcina A subgenus of the genus *Planosarcina* Migula created by Migula (1895, p 20) to include those forms in which the cell contents are colorless and without sulphur granules The other subgenera are *Thiocystis* Winogradsky and *Lamprocystis* Winogradsky

Eupseudomonas. A subgenus of the genus *Pseudomonas* Migula proposed by Migula (1895, p 29) to include those forms in which the cell contents are colorless and without sulphur granules The second subgenus is *Chromatium* Perty

Eusarcina. A subgenus of the genus *Sarcina* Goodsir proposed by Migula (1895, p. 18) to include forms in which the cell contents are colorless and free from sulphur granules The second subgenus is *Thiosarcina* Winogradsky.

Euspirillum. A name first proposed by Trevisan (1889, p 24) for the first subgenus of *Spirillum*. The description is "Spirilli tipici, costantemente con forme a guisa di cavaturaccioli." Twenty-five species are included. The second subgenus is *Pseudospirillum*. This name was also used by Migula (1895, p 33) for the first section under his genus *Spirillum* to include those forms in which the cell contents are colorless differentiating it from the second section, *Thiospirillum* Winogradsky.

Meyer (1912, p 3) included in this section of the genus *Spirillum* all those species which contain no bacteriopurpurin and are motile.

Eu-Spirillum. A subgenus of the genus *Spirillum* proposed by De Toni and Trevisan (1889, p 1009) with the following diagnosis "Spirilli regulariter spirales, rigidi, abbreviata, spiris uniformibus." Twenty species are included, among them *Spirillum undula*, *S. tenue* and *S. serpens*. This is a variant of *Euspirillum*.

Euspirosoma. A subgenus of the genus *Spirosoma* created by Migula (1900, p 955) to include those forms in which the cells are free, that is not embedded in gelatin. The second subgenus was *Myconostoc*.

Euvibrionaeae. A subtribe of bacteria proposed by Trevisan (1879, p 137) with the following description "Somatia libera eiamosa." It was included in the tribe *Vibrionaeae*, and contained the following genera *Mantegazzaea*, *Metallacter*, *Leptotrichia*, *Beggiatoa*, *Crenothrix*, *Vibrio*, *Spirillum*, *Spirochaete*, and *Spiromonas*.

Fenobacter. This name has been used by Beijerinck (1900, p 200) as a generic designation of organisms belonging to the hay bacillus group. Apparently it has not been used by other authors. It may possibly be regarded as a synonym of *Bacillus* although the author emphasizes its relationship to *Aerobacter*. It is rejected as invalid by Erwin F. Smith (1905, p 174).

Ferribacterium. A generic name proposed by Brussoff (1916, p 547) for a species *Ferribacterium duplex* (the type). The summary of the characters which he gives follows:

1 *Ferribacterium duplex* ist ein unbewegliches, dunkel oder hellgelb gefarbtetes *Bacterium*.

2 Es hat meistens die Form eines Doppelstabchens mit abgerundeten Enden, erscheint aber auch in der Form von Stabchen und kurzen Ketten.

3 Die Längen der Stabchen mit dem Gallerthof schwankt zwischen ca. $2\frac{1}{2}$ und 5μ , die Breite ist ca. $1\frac{1}{2}\mu$ gleich und ungefähr dieselbe bei allen Individuen.

4 In jungen Torf-Wasser-Kulturen und in allen, selbst in alten Eisenammonciatrat- und Eisenpepton-bouillonkulturen besteht jedes Individuum meistens aus drei Elementen: dem inneren Stabchen, dem es umgebenden Gallerthof und der

Eisenausscheidung um ihn herum, die Eisenausscheidung kann auch fehlen. Diese Formen färben sich bei der Behandlung mit gelbem Blutlaugensalz und Salzsäure intensive blau.

5 In älteren Torf-Wasser-Kulturen kommen Individuen vor, bei denen die Eisenausscheidung um den Gallerthof herum immer fehlt, dagegen ist dieser von einer dunklen Hülle umgeben, welche für die Lösung des gelben Blutlaugensalzes undurchlässig ist. Deswegen lassen sich diese Individuen mit der Eisenreaktion nicht blau färben.

6 Die dunkle Hülle hat meistens eine ovale Form, zeigt aber manchmal auch unregelmässig zerhackte Konturen. Es kommen auch Fälle vor, wo sie teilweise fehlt, und dann färben sich solche Bakterien leicht blau.

7 Die unter 4, 5, und 6 angeführten Formen kommen in älteren Torf-Wasser-Kulturen nebeneinander vor.

8 Die Kolonien bilden eine Haut an der Oberfläche des Wassers oder der künftlichen Nährösungen.

9 Die Haut an der Oberfläche des Wassers ist schwach regenbogenfarbig oder metallisch glänzend und makroskopisch deutlich sichtbar, wogegen

10 Die Haut auf Eisenammonenitrat- und Eisenpeptonbouillonlösungen immer makroskopisch unsichtbar bleibt und nur durch das Übertragen auf den Objektträger bemerkbar gemacht werden konnte.

11 Mikroskopisch erscheint die Haut immer mehr oder weniger deutlich gelb gefärbt.

Fixococcus. A name proposed by Meyer (1912, p. 5) for the first section of the genus *Micrococcus*. His description is "Die Formen, bei denen keine Geisseln nachgewiesen wurden". The other sections are *Planococcus* (motile) and *Rhodococcus* (with sulphur granules).

Fixosarcina. A name proposed by Meyer (1912, p. 5) for the second section of the genus *Sarcina* with the description, "Die Spezies bei denen Geisseln nicht nachgewiesen wurden".

Fixostreptococcus. A name applied to Meyer (1912, p. 4) to the second section of the genus *Streptococcus*. This includes "Alle Spezies für welche Geisseln nicht bekannt wurden".

Flavobacterium. A name used by Bergey *et al.* (1923, p. 97) for the second genus of the tribe *Chromobacterae* with the description

Rods of medium size, occurring in water and soil, forming a yellow to orange pigment on culture media. Characterized by feeble powers of attacking carbohydrates, occasionally forming acid from hexoses but no gas. Motile or non-motile. Generally Gram-negative.

The type species is *Flavobacterium aquatilis* (Frankland).

Flemmingillus. A name given by Heller (1922, p. 15) to the sixth genus of the subfamily *Clostridoideae*. The definition follows:

Clostridoideae that do not liquefy gelatin. They produce little gas or acid in meat medium. They do not grow very abundantly in milk or change it. They

do not show any marked tendency to split sugars. Slender Gram-negative rods with oval terminal spores. Colonies in deep agar lenticular, may show an areola of fine radiations. Frequently found in moulds, not pathogenic for guinea-pigs.

Type species *F. cochlearius* (*Bacillus cochlearius* Douglas, Fleming and Colebrook) as described by the Committee. The type is highly motile and it split none of the carbohydrates in which it was grown. It was described as bacillus III type C by McIntosh.

Flexnerella. The first subgenus of the genus *Shigella* named by Castellani and Chalmers (1919, p. 937). It differs from the second subgenus (*Shigella*) in that mannitol is fermented. It is subdivided into two groups, the Flexner group and the Pseudodysentery group. Later (1920, p. 606) two species are listed. The type is *Shigella flexneri* Castellani and Chalmers.

Fluoromonas. A generic name proposed by Orla-Jensen (1921, p. 271) to include his earlier described *Dentromonas* and *Liquidomonas*. He says:

As the property of liquefying gelatin also cannot be adopted as a generic character but only as a species character, I think it would be best to group together my earlier genera *Dentromonas* and *Liquidomonas* in a single genus, which can be conveniently termed *Fluoromonas*, as the bacteria of this group are ordinarily fluorescent. I cannot agree to call this genus *Pseudomonas* merely out of regard for so-called priority, as each and all of the bacteria which belong to the order under consideration are really Pseudomonades as well.

No species is named.

Fusiforminae. A subtribe named by Buchanan (1918, p. 44) to include the single genus *Fusiformis*.

Fusiformis. A generic name proposed by Hoelling (1910, p. 240) to include the fusiform bacteria. He concludes that those organisms usually included in the name *Bacillus fusiformis* should not be placed in the genus *Bacillus*. He mentions as species *Fusiformis dentium*, *F. muris*, *F. termatidis*. He describes the morphology of the last named species in detail. He states that the protoplasm of the young organisms is relatively homogeneous, sometimes staining more intensely toward the pointed ends. The larger cells sometimes show an alveolar plasma structure. The nuclei are in general stained dark. The nuclei in some are oval and show a definite structure, others show, especially in mononucleate cells, as a broad band. Evidently the *Bacillus fusiformis* is *Fusiformis termatidis*. The name is accepted by Dobell (1911, p. 486).

Vuillemin (1913) suggested the revival of Trevisan's name *Mante-gazzaea* for this group as one of the "*formogenera conservanda*" with the fusiform bacillus of the medical writers as the type. The name as used by Trevisan, however, was used principally for certain of the sulfur bacteria and not for forms related to this type.

Winslow *et al* (Committee Soc Am Bact, 1917, p 555, 1920, p 199) and Bergey *et al* (1923, p 388) recognize this genus with the following description

Obligate parasites. Cells usually elongate and fusiform, staining somewhat irregularly. Filaments sometimes formed, non-branching. Non-motile. No spores. Growth in laboratory media feeble.

The Type species (?) is *Fusiformis termitidis* Hoelling.

Buchanan (1918, p 43) used the following description

Obligate parasites. Cells usually elongate and fusiform. Gram-negative? Anaerobic. Non-motile. No spores. In some respects approaching the spirochetes in morphology.

The type species is *Fusiformis termitidis* Hoelling (possibly *F. dentium*).

Castellani and Chalmers (1919, p 959) have suggested the name *Bacteroides qv* for this group.

Apparently *Fusiformis* constitutes a valid generic designation with *Fusiformis termitidis*? Hoelling as the type.

Gaffkya A genus of bacteria proposed by Trevisan (1885, p 105) based upon *Micrococcus tetragenus* Gaffky as a type. The description of the genus is "Colonie di 4 cocci globosi, avviluppati da una capsula mucosa, ialina, globulosa, da ultimo liberi. Specie unica." The diagnosis given by De Toni and Trevisan (1889, p 1042) follows

Cocciglobosi vel divisionis tempore ovoidei, in muco matricali nidulantes, quaterni, cystidibus specialibus parvulis, gelatinosis, tenuiculis, demum dilatantibus, solitariis, nunquam in acervos consociatis, obducti. Cystides universales nullae. Coccocorum divisio in duas directiones. Sporae ignotae.

Four species are included. The type species may be taken as *Gaffkya tetragena* (Gaffky) Trevisan. Should this organism prove to be entitled to generic recognition, the genus *Gaffkya* would be valid. The name does not appear to have been used by other authors. It is rejected by Erwin F. Smith (1905, p 174).

Gaffkyaeeae. The second subtribe of the tribe *Ascococceae* proposed by Trevisan according to De Toni and Trevisan (1889, p 1042) with the following diagnosis "Cocci cystidibus specialibus obducti, Cystides

universales nullae" Two genera are included *Chlamydatomus* Trevisan and *Gaffkya* Trevisan

Gaillonella. A genus of algae named by Bory de St Vincent (1823, p 393) Enlows states

Gaillonella Bory de St Vincent, 1823 Dict classique d'Hist Nat, v 4, 1823, p 393 See also idem, v 7, 1825, p 101 Simple cylindrical filaments, articulated, each section including two capsular corpuscles, spheroidal, transparent, even when filled with ferruginous coloring matter, and divided into two equal parts by a "dissépiment" which appears as a line "Nous n'hésitons pas à regarder les Gaillonelles comme de simples végétaux."

Type species (original designation) —*Conserva moniliformis* Muller Places *C nummuloides* here also

No species were described in 1823

In 1824 Agardh created the genus *Meloseira* with 5 species The following year (1825) Bory included two of these species in his genus *Gaillonella* Ehrenberg (1833, p 294) gives the following synonymy and description

Gaillonella Bory de St Vincent 1823, *Meloseira* Agardh 1824, *Gaillonelle* Familie der Stabthierchen, Bacillaria Character generis *Polygastricum*, *anenterum*, *Gymnicum*, *Pseudopodium*°, *loricatum* *Lorica* subglobosa aut oblonga bivalvis divisione, spontanea intra vaginam deciduam peragenda cateniformis, filiformis

He described *Gaillonella lineata* Bory In 1835 Ehrenberg included in the genus an organism which he termed "*Gaillonella ferruginea*," and which he believed might be the same as the *Hygrocrocis ochracea* of the botanists In the following year (1836, p 217) Ehrenberg discussed this organism at greater length including it with several of the species of diatoms of the same genus He terms it the "Eisenockerthierchen," and notes finding its "shell" in many places as a fossil His figure shows his conception of the organism to be a very minute diatom in chains

In his later monumental publication on the infusoria (1838, p 166) the name was changed to the form *Gaillonella*, a form which has been commonly accepted The generic diagnosis is given as follows

Gaillonella Gaillonelle Character Animal e familia Bacillariorum, liberum lorica simplici, bivalvi (silicea), cylindricum globosum aut disciforme, spontanea imperfecta divisione cateniforme

Seven living and three fossil species are described Three of these had been previously listed as species of the genus *Meloseira* by Agardh

Included is the species (p 169) *Gallionella ferruginea*, "rostförmige Dosenkette" with the following description "*G. corpusculis* terminatis, utrinque convexis, ovatis, ferrugineis, filis articulatis sagittatis, subramosis" with the following synonymy *Conferva ochracea* Roth? Catal bot, I, p 165, Tab 5, Fig 2, 1797, Schumacher. Enumeratio plant Zeelandiae, 1803, II, p 105, Dillwynne? Synopsis of brit Conferv, 1809, pl 62 *Lyngbya ochracea* Leiblein Floia, bot Zeit, 1827, p. 260, 280. *Oscillatoria* ? *ochracea* Lyngbye ? Tent Hydrophyt dan, T 26, C—————? Agardh ? Syst Alg, 1824, p 69, *incerta species* *Gallionella ferruginea* Ehl Taf x dieses Werk, 1835

Gallionella ferruginea Bericht der Berl Akad d Wissensch., 1836, 27, Jan, p. 52, 84.

Why Ehrenberg abandoned the earlier specific *ochracea* is not evident

Ralfs in 1843 (p 351) described the same organism in the diatom genus *Melosira* as follows "*Melosira ochracea* Frustules very slender, convex at each end, ovate, not striated, ferruginous, filaments often connected together in a subramose manner" He gives as synonyms the following *Gallionella ferruginea* Ehrenberg, *Conferva ochracea* Dill. t 62 and *Oscillatoria ochracea* Grevil He further states

Filaments are so slender and joints so obscure that I have been unable to determine the form of the frustules, and therefore have taken the specific characters from Ehrenberg Having received from Mr Dillwyer a specimen of his *Conferva ochracea*, which I am able confidently to refer to this species, I have restored the original specific name

Kutzing (1849, p 28) included this organism in his genus *Gloeotila* The description given is as follows

Gloeotila Kutzing, *Trichomata eramosa viridia mucosa, ex cellulis monogonimicis composita* (Paludosae, mundatae) *G. ferruginea* Kg Phyc Germ p 191 *G. ochracea*, trichomatibus brevissimis moniliformibus, articulis ovato-globosis Diam 1-2000" *Gallionella ferruginea* Ehrenb Infus, T 10, f 7 *Melosira minutula* Brebiss Falais p 42, pl 5 In fontibus martialibus (v v) Cf Kg Bacillac, p 56 *Gallionella ferruginea*

Griffith (1853, p 28) pointed out that this organism (*Gallionella ferruginea* Ehrenb) is not a diatom, and that it cannot be retained in Kutzing's genus *Gloeotila* He states that the organism consists of two intertwined filaments, and proposed the name *Didymohelix*

According to Mohsch, Mettenheimer (1856, p 139) inclined to the view that *Gallionella* and *Leptothrix* are identical

According to Lohnis (1913, p 704) this organism was placed in the genus *Gloeosphaera* by Rabenhorst, with the name *G ferruginea* De Toni and Trevisan (1889, p 1007) give as authority Rabenhorst, Alg Mitteleur, n 387 Kützing (1865, p 53) used *Gallionella* as the second subgenus of *Melosira* Later (p 56) he says "Ganz ausgeschlossen muss werden *Gallionella ferruginea* Ehr. welche gar keine Diatomee, sondern eine Confeiva ist" Kütchner (1878, p 250) placed this organism in the genus *Spirulina* with the name *Sp ferruginea* Hansgirg (1888, p 264) uses the designation *Spirochaete ferruginea* (Ehr.) Hansg. De Toni and Trevisan (1889, p 1007) included the species in the genus *Spirillum* because of its curved form De Toni (1891, p 1331) used *Gallionella* Boiy as a valid generic name of a diatom genus Hansgirg (1873, p 184) used the designation *Leptothrix ochracea* to include this organism not recognizing any distinction between *Leptothrix* and *Gallionella* Migula (1897, p 321) discussed this organism

In his "System" Migula named the organism *Chlamydothrix ferruginea* (1900, p 1031) In this genus was included the *Leptothrix ochracea* also

Ellis (1907, p 516) returns to the designation "*Gallionella ferruginea*," He states,

The structure is like that of a hairpin, spirally wound round itself The spiral winding produced a number of loops, the number of which may be few or many The thickness of the threads varies usually from $\frac{1}{4}$ to $\frac{3}{4}$ μ The winding may be loose or tight The structure of the cell is perfectly homogeneous Migula states that an extremely delicate membrane is present, but I have not been able to confirm this observation Multiplication takes place freely by the cutting off of small portions, which then elongate to form a new individual Conidia formation is also quite common, though not universal as in *Leptothrix ochracea* The process of formation of conidia is identical with the same in this organism, as is also the size and shape of conidia The deposit of iron is the same as in other iron bacteria In no stage of its life history was any motility observable

Orla-Jensen (1909, p 331) says

Die Schraubenform des *Leptothrix* sind *Gallionella* und *Spirophyllum* Jedoch kann den mehr oder weniger zufälligen Biegungen und Flechten der *Gallionella*-Faden nicht dieselbe systematische Bedeutung wie den regelmässigen Windungen der starren Spirillen zugeschrieben werden, und wir tun daher besser, *Gallionella* nur als eine *Leptothrix*-Art aufzufassen

The name *Gallionella ferruginea* Ehrenberg was also accepted by Mohs (1910, p 22) who studied the organism at length and has given us the best description He states

Faden geschlangelt, einfach, oder je zwei schraubenförmig dicht um einander gewunden und daher einen Zopf bildend. Hierdurch gewinnt der Faden oft ein kettenartiges Aussehen. Faden in der Jugend farblos, später infolge von Eisenhydrateinlagerung gelbbraunlich bis rostrot. Gewöhnlich sind die Doppelfäden eng verschlungen, stellenweise lösen sie sich aber deutlich in die Einzelfäden auf, ähnlich wie man dies bei *Spirulina* beobachten kann. Ist der schraubig gewundene Faden breiter der vereinigen sich zwei solchen Fäden zu einem Zopf, so entsteht ein kettenartiger Faden, dessen einzelne Glieder früher fälschlich für Sporen gehalten worden sind. Die Einzelfäden zeigen keine Gliederung in Zellen, selbst nicht nach Behandlung mit Farbstoffen und verdünnten Säuren und zeigen auch keine nachweisbare Scheide. Durch hochgradige Inkrustation mit Eisenoxyd können die Fäden und ihr Verlauf ganz verdeckt werden. Breite der Fäden bzw. Zöpfe 1-6 μ . Länge bis 270 μ und darüber. In eisenhaltigen Wässern häufig und weit verbreitet. Lässt sich vorläufig nicht kultivieren.

He states further that Migula first figured the organism accurately, but that he cannot agree with this author in its inclusion in the genus *Chlamydothrix* for no sheath is present. Furthermore there is not the slightest trace of the jointing of the filaments into cells. He also concludes that the *Spirophyllum ferrugineum* of Ellis (1897, p. 21) is probably not specifically distinct, for he has found intermediate forms. He suggests therefore the name *Gallionella ferruginea* var. *lata*. He is also inclined to regard the *Nodofolium ferrugineum* of Ellis (1910, p. 32) as a broad form of *Gallionella ferruginea*. He concludes further, that *Leptothrix* (*Chlamydothrix*) and *Gallionella* are distinct. His pure cultures of *Leptothrix* never gave rise to *Gallionella* forms. Therefore Schwere (1908, p. 53) was in error in his statement "Tout nous port donc a croire avec Hansgirg, qu'il s'agit d'une seule et meme espèce."

Gallionella (or *Gallionella*) since the time of Ehrenberg has been recognized as a valid genus of diatoms or a subgenus of *Melosira* by many writers on algae.

There does not seem to have been sufficient reason for Ehrenberg's change of spelling *Gallionella* to *Gallionella*. The recognition of the fact that the *Gallionella ferruginea* of Ehrenberg was not a diatom, and the creation of the genus *Didymohelix* by Griffiths would appear to be valid. The original specific designation was *ochracea*, which was changed to *ferruginea* by Ehrenberg. It would seem that this organism should therefore be termed *Didymohelix ochracea*, and that *Gallionella* should not be used as a bacterial genus.

Galactobacterium. A casual name given by Guillebeau (1890, p. 43) to an organism associated with inflamed udder in a cow. No description is given.

Galactococcus. A generic name proposed by Guillebeau (1890, p 32) for three species They are described by Enlows (1920, p 44) as follows

G versicolor n sp Cocci of about 1μ diameter Nonmotile Gram positive Long chains in milk, which is rapidly acidified Aerobe, Belongs to the "haufigeren Mastitispilzen" *G fulvus* n sp Cocci of not more than 1μ diameter Nonmotile Gram positive Ochre-yellow on potato *G albus* n sp Cocci about 1μ diameter Nonmotile. Gram-positive White colonies on milk gelatin, which is not at all or only slightly liquefied Dirty white growth on potato All of these species found in milk from an inflamed udder

Apparently the organisms described belong to the genus *Staphylococcus*, of which *Galactococcus* may be regarded as a synonym The type species probably should be designated as *Galactococcus versicolor*, the first species described

Gallionella. A spelling of *Gaillonella* introduced by Ehrenberg (1838, p 166) and commonly used by subsequent writers For discussion, see *Gaillonella*

Glaeogenes. A tribe of Schizophytes named by Cohn (1875). The tribe included many forms now classed among the bacteria as well as forms now included with the blue green algae The tribe is characterized by having its cells free or united into glairy families by an intercellular substance

Gleobacter A casual name used by Fischer (1895, p 135) according to Enlows (1920, p 44) to include capsulated rod-shaped bacteria *Klebsiella* Trevisan is stated to be a synonym

Ghabacteria A name proposed by Billroth (1874, p 5) to designate a growth form of his *Coccobacteria septica* in which the organisms appear as rods embedded in a mass of gelatin It is not a generic name.

Ghiacoccus. A name proposed by Billroth (1874, p 5) to designate a growth form of his *Coccobacteria septica* in which the cells appeared as cocci united by gelatin into irregular colonies He proposed this name to replace an earlier designation *Zoogloea* Cohn With the spelling *Ghiacoccus* it was rejected by E F Smith (1905)

Ghiacoccus. See discussion of *Ghiacoccus* Maggi (1886, p 82) gives *Ghiacoccus aceti* as a variety of his *Micrococcus aceti*

Ghi-Kokkus. A variant spelling of *Ghiacoccus* used by Klebs (1887, p 310)

Ghschrobacterium. An aberrant spelling of *Ghschrobacterium* (g v)

Ghschrobacterium A casual designation introduced by Malerba and Sanna-Salaris (1888, p 196) for an organism found by them to

produce a high degree of viscosity in urine. The same organism was studied by Rothmann (1904, p 491) under the name *Glischrobacterium*. This term has not been used as a generic name. It is rejected by Erwin F. Smith (1895, p 174)

Gloeosphaeria. Rabenhorst (alg Mitteleur No 387) distributed an organism as *Gloeosphaeria ferruginea* (1854, p 43). The name has never been recognized by other writers. The organism is one of the iron bacteria, variously termed *Chlamydothrix ferruginea*, *Gallionella ferruginea* or *Didymohelix ferruginea*. The name may probably be regarded as nonvalid as a bacterial generic designation.

Gloeotila. A name used by Kutzing (1843, p 245) for a genus of algae with the following description

Trichomata parenchymatica viridia, simplicia, monogonimica, libera. Cellulae gelinae delicatissimae, mucosae. Gonidia oblongo-angulata, vel rotundata, hinc flaccida, illinc, turgida (nunquam longitudinaliter divisa)

Six species are recognized. Later this author (1849, p 363) included one species *Gloeotila ferruginea* which is generally regarded as a synonym of *Gallionella ferruginea* or *Didymohelix ferruginea*, one of the bacteria. The description given is "*G. ochracea, trichomatibus brevissimis moniloformibus, articulis ovato-globosis*. Diam 1/1200—*Gallionella ferruginea* Ehrenb." In addition nine other algal species are described. This name is also used by Perty (1852, p 215). *Gloeotila* is a valid genus of algae and should not be used for bacteria.

Glucobacter. See *Glycobacter*

Glycobacter. A generic name proposed by Wollmann (1912, p 613) for a group of amylolytic intestinal bacteria. Two species were described of which the first, *Glycobacter proteolyticus* may be regarded as the type. This organism was isolated from the ileum of a monkey. It is a rod, usually straight, rarely slightly bent, about 1 μ in diameter and 2 to 6 or even 10 μ in length. Cells usually isolated, rarely in pairs, never in chains. The cells stain readily, but take the Gram stain poorly. They do not give the granulose reaction. Spores are readily produced, on agar within twenty-four hours. The spore is subterminal or medium, the cell somewhat inflated. The spores are resistant even to boiling temperatures. The bacilli are very motile.

On sloped agar the cultures are quite transparent and delicate, where colonies are isolated they reach 1 mm in diameter. In sugar gelatin shake cultures, small irregular colonies appear in twenty-four hours, and gas is formed. Gas is formed in glucose broth in the first twenty-four hours. Milk is feebly coagulated after forty-eight hours. Whey

is clear, the casein is slowly digested, the reaction remains acid. In gelatin the colonies are round, liquefied. Egg white is not attacked appreciably. On potatoes a granular layer in twenty-four hours. Starch is rapidly liquefied in the presence of a suitable source of nitrogen. It attacks albumins, but does not seem to break them down completely. Amylase may be readily demonstrated in filtrate from potato broth culture. With glucose, levulose, maltose, saccharose and lactose it gives an acidity corresponding to 1.2 to 1.47 grams of H_2SO_4 per liter, in lactose about 0.49 gram. Small amount of alcohol produced.

The second species *Glycobacter peptolyticus* resembles the first in digestion of starch, but does not cause proteolysis.

The genus *Glycobacter* evidently includes the amylolytic aerobic spore producing organisms usually assigned to the genus *Bacillus*.

Gonium. A genus of algae named by Muller (1773). Ehrenberg assigned to it the species, *Gonium hyalinum*, one of the bacteria. This organism was removed from this genus by Schroter (1886, p. 151) who gave to it the name *Lampropedia hyalina*. According to Robin (1847, p. 110), Busk (1843) regarded *Sarcina ventriculi* as a species of *Gonium*. The genus *Gonium* is a valid genus of algae, but not of bacteria.

Gonococcus. A name first used as a casual designation for the organism which is the specific cause of the disease gonorrhea. The name is usually ascribed to Neisser (1879, p. 497).

Apparently its use as a generic name has been without formal designation as such. Migula (1896, p. 16) gives *Gonococcus gonorrhoeae* Neisseri as a synonym of his *Micrococcus gonorrhoeae*. However, examination of the work of Neisser fails to show this designation, it cannot be validated by quotation in synonymy (See Botanical Code).

Lindau (1898, p. 100) named the species *Gonococcus neisseri*.

This organism was earlier named *Neisseria qv* by Trevisan. *Gonococcus* as a generic designation is a synonym, and must be ascribed to Lindau (?).

Graciloidae. A name given by Castellani and Chalmers (1919, p. 932) to the third tribe of the family *Bacillaceae*. The description (p. 958) is

Bacillaceae growing very slowly and scantily on ordinary and blood media, without endospores or capsules, neither fluorescent nor chromogenic.

Type Genus *Graciloides* Castellani

Graciloides. The name given by Castellani and Chalmers (1919, p 933) to the single genus of the tribe *Gracilordeae*, with characters of the tribe, *qv*, and the type species *Graciloides albofaciens* (Castellani) Castellani and Chalmers

Granulobacillus A genus named by Schattenfroth and Grassberger (1899, p 202) for two species of anaerobic bacteria Enlows (1920, p 45) states that it differs from *Granulobacter* Beijerinck in that it includes both motile and non-motile organisms The species named are *Granulobacillus saccharobutyricus immobilis liquefaciens* and *G. saccharobutyricus mobilis nonliquefaciens* The names are polynomials, hence invalid This invalidates the generic name as well

Granulobacter. A genus of bacteria created by Beijerinck (1893, p 7) He named at this time four species His generic description follows

Obligat- oder temporar anaerobe Gahrungsbakterien (Für die Erklärung des Ausdruckes "temporar anaerobie" verweise ich auf Par 12)—welche bei vollstandiger Anaerobiose sich theilweise oder ganz mit granulose anfullen und dann Clostridiumform annehmen Bei Gegenwart von Sauerstoffspuren entstehen schnell bewegliche Stabchen, welche mit Jod gelb werden Sporen entstehen in den Clostridien, und können einige Secunden oder Minuten auf 95°C bis 100°C in den Nahrflüssigkeiten erhitzt werden, wodurch die Entfernung von verunreinigten Bakterien möglich ist Unter den Gahrungsproduction finden sich immer Kohlensäure und gewöhnlich auch Wasserstoff, während Methan vollständig fehlt

The first species described (which may be considered the type) is *Granulobacter butylicum* This is said possibly to be identical with Gruber's *Bacillus amylobacter I* Other species described are *G saccharobutyricum*, *G lactobutyricum* and *G Polymyxa*

Later (1896, p 1) the generic diagnosis is as follows

Granulobacter: Bacterie de fermentation, anaerobies temporaires ou obligatoires, qui si l'anaerobiose est complete se remplissent soit en partie soit en totalite de granulose et prennent alors la form de clostridiuims S'il y a des traces d'oxygene en presence il se forme des batonnets à motilité tres vive, quise colorent en jaune par l'iode Des spores se forment dans les clostridium et peuvent resister pendant quelques secondes ou quelques minutes, dans les liquides de culture, a une temp de 95-100°C On trouve toujours parmi les produits de fermentation de l'acide carbonique et d'ordinaire aussi de l'hydrogen tandis que le methan fait complètement défaut

The species first named is *G butylicum* The following description given by Stockhausen (1907, p 42) characterizes the genus well

Zur Gattung *Granulobakter* stellt Beijerinck alle jene obligat oder temporär anaerobiotischen Gärungsbakterien zusammen, welche bei vollständiger Anaerobiose sich teilweise oder ganz mit Granulose anfüllen, den Clostridium form annehmen. Bei Gegenwart von Sauerstoffspuren entstehen schnell bewegliche Stäbchen. Die in den Clostridien enthaltenen Sporen halten einige Minuten 95–100° aus. Unter den Gärungsprodukten finden sich immer Kohlensäure, meist auch Wasserstoff. Zu dieser Gattung gehören die Bakterien der Butylalkoholgärung (*Gian butylacum*) der Buttersäuregärung und die Bakterien der Flachsrotte, *Gran pectinovorum* und *Gran urocephalum*.

This name has been used for the most part only by Beijerinck and his pupils. Lindner (1909, p. 515) defines the genus as follows: "Obligat oder temporär anaerobe Gärungsbakterien, welche bei vollständiger Anaerobiose sich teilweise oder ganz mit Granulose anfüllen und dann Clostridium form annehmen."

Granulobacter may probably be regarded as a synonym of *Clostridium*. It is rejected by Erwin F. Smith (1905, p. 174).

Granulobakter. A variant of *Granulobacter*, used by certain German authors as Grassberger and Schattenfroh (1899, p. 702), Stockhausen (1907, p. 42).

Grippestreptokokkus. A casual designation proposed by Seligmann (1912, p. 81) for a streptococcus associated with an epidemic of "Grippe" among infants.

Gummibacillus. According to Migula (1900, p. 577) this is a casual designation used by Loeffler for the organism *Bacillus lodermos* Flugge. It is rejected by Erwin F. Smith (1905, p. 174).

Gymnobacteria. A group name for the "naked" bacteria as distinguished from the Trichobacteria used by Messea (1891) according to Kruse.

Gyrococcus. A generic name proposed by Glaser and Chapman (1912, p. 223) for the species (type) *Gyrococcus flaccidifex*. Enlows (1920, p. 46) summarizes the description:

Cells in free state spherical, becoming slightly oblong just before division. Division in 1 direction only. After division each half may be spherical or may come to an abrupt tip, assuming a more or less heart-shaped appearance. Frequently the two halves are unequal, one-half may be spherical while the other may be more or less heart-shaped, or slightly oblong. If cells remain connected after fission, chains of 3 or 4 are formed. Diameter of cells 0.51 to 0.85 μ . No endospores. Capsule distinct. Motile—progressing in a gyrating manner, but no flagella were stained. Gram negative. Very closely resembles the Pneumococcus, except that it is Gram negative, and motile. Cause of wilt disease or "flacherie" of the gypsy moth.

Haematococcus. This generic designation of the "red snow" alga was first given by Agardh (1828, p 45) to the species *Haematococcus nivalis*. Since that time this name has been frequently used for an algal genus. It is generally now regarded as a synonym of *Sphaerella*, according to Hazen (1896, p 238) and West (1904, p 189). Wille (1903) regards the generic name as valid and as replacing *Sphaerella*.

In a different sense the same generic name has been used a few times in bacteriology. V Babes (1889, p 81) in describing the causal organism of bovine hemoglobinuria used the phrase (p 103) "fui die Krankheit Charakterischen Hamatokokken" and later (p 106) "Neben dem Haematococcus finden sich," etc.

The same organism was earlier discussed by Babes (1888, p 692) and according to several authors (Doflein, 1911, p 815, Eisenberg, 1891, p 271) the name *Haematococcus* was here first used. A careful examination of the article fails to show that the term is here used. Babes regarded the organism as a bacterial form.

The name *Haematococcus* is invalid in bacteriology because of the prior existence of the algal genus of the same name. Furthermore no species name seems to have been used, the term as employed by Babes was apparently only as a casual designation. The first use of the term in a strict generic sense seems to have been by Eisenberg (1891, p 271) with the spelling *Hamatokokkus*, and the species *H. bovis*. This organism is now recognized as a protozoan. The generic name *Haematococcus* is not valid as a protozoan genus because of previous use.

Hamatokokkus. A variant of *Haematococcus* *qv* used by Eisenberg (1891, p 271).

Haemophyleae. The name given by Castellani and Chalmers (1919, p 932) to the second tribe of the family *Bacillaceae*. Growth in laboratory media is poor, the organism is Gram-negative, grows best on blood media. The single genus is *Haemophilus*.

Haemophilus. A generic name (variant of *Hemophilus* *qv*) used by Castellani and Chalmers (1919, p 933) as the single genus of their tribe *Haemophyleae*. The type species is designated as *Haemophilus influenzae* (Pfeiffer) Castellani and Chalmers.

Halobacterium. A generic name proposed by B Fischer (1894, p 19) to include certain organisms living in salt water, including *Halobacterium pellucidum* (p 19), *H. roseum* (p 19), *H. liquefaciens* (p 21), *H. polymorphum* (p 25), *H. aurantiacum* (p 26), *H. rubrofusum* (p 28), *H. purpurum* (p 30). The description of *H. pellucidum* (p 22) which may be regarded as the type follows.

Kleine bis mittelgrosse Kurzstabchen, fast wie Kokken, einzeln oder zu 2 oder in kurzeren Ketten zu 4, 6 selten mehr vereinigt. Manchmal die Kurzstabchen in ähnlicher Weise angeordnet wie bei Sprossverbänden der Bierhefe, alsdann gewöhnlich in Zoogloeamasse eingebettet. Neben den Kurzstabchen oft auch etwas längere, spindelförmige Stäbchen mit hier und da soeben angedeuteter kommaartiger Krümmung, zuweilen aber auch typische Komma- und S-Formen, ferner kurze, ungegliederte, gerade oder gebogene Fadenstücke, theils wie um die Längsachse gehreht und von ungleicher Dicke, theils deutlich schraubenförmig, theils unregelmässig gewunden. Oft erhebliche Unterschiede in der Grösse. Manchmal lebhaft Eigenbewegung. Leicht färbbar. Kolonien in Gelatineplattenrunde, graue, durchscheinende Tropfchen, die tiefgelegenen mit gelblichem Stich, nicht verflüssigend. Bei schwacher Vergrösserung rund oder citronenförmig, schmutziggelb, gleichmässig oder feinkörnig, oft konzentrische, manchmal wie mit kleinen Luftblaschen besetzt, manchmal von gelappter oder hockeriger bezw. rosettenartiger Form. In Stichkulturen ausgesprochenes Oberflächenwachsthum, anfangs durchscheinende, später mehr, trübe, trockene und braunliche, oder auch grauweisse, runzlige, blattartige Auflagerung. Auf Stichkulturen feuchter, grauer, durchscheinender, weiterhin oft runzligfaltiger Belag. Für Mäuse nicht pathogen. Im Oceanwasser weit verbreitet. Wegen der durchscheinenden Beschaffenheit der jüngeren Kulturen weiterhin als *Halbacterium pellucidum* bezeichnet.

This has generally been regarded as a physiological genus, and has not been recognized as valid by subsequent authors. It is rejected by Erwin F. Smith (1905).

Haplobacteriacei. An ordinal name proposed by Fischer (1895, p. 138) coordinate with *Trichobacteriacei*. It includes all those bacteria whose cells are not united into true filaments. He says

Das unterscheidende Merkmal liegt in dem Bau des Vegetationskörpers, der bei den Trichobakterien immer ein unverzweigter auf einzelnen aufgebauten Faden ist, während die Haplobakterien einen einzelligen Vegetationskörper besitzen, der kugelig, stabartig oder schraubig ist.

With the spelling *Haplobacterinae* (q v) this name is used in Coppen Jones (1897, p. 32) translation.

Haplobacterinae. An ordinal name first used with the spelling *Haplobacteriacei* by Fischer (1895, p. 138). Coppen Jones (1897, p. 32) in the translation of Fischer's Vorlesungen used the form *Haplobacterinae*. It was later used by Benecke (1912, p. 1888) for a group coordinate with *Desmobacterinae* to include all non-filamentous bacteria. According to the last author it contains six families, *Coccaceae*, *Bacillaceae*, *Spirillaceae*, *Rhodobacteriaceae*, *Mycobacteriaceae* and *Myrobacteriaceae*.

Haplobakterien. A casual name used by Fischer (1890, p 119) for his group *Haplobacteriaceae* q v

Helicobacterium. A generic designation used by Miller (1880, p 119) for a species *Helicobacterium aerogenes*. The organism is a slender motile rod, single or in chains, and may develop to long, wavy bent filaments, which may resemble "Spirulinen". It forms transparent white or only weakly yellow colored colonies which are exceedingly variable in form. In stab cultures the organism grows throughout the length of the stab with a light yellow color and covers the surface in twenty-four hours with a thin, scarcely visible, bluish, dry layer. On stroke culture there develops a thin broad growth, which in transmitted light shows a flocculent or crystalline appearance. Growth on agar is not characteristic, growth on potato is slow, with a dry surface and yellow brown color. Poor growth in the absence of oxygen. It is again described by Miller (1892, p 370). An *Helicobacterium klebsii* is also described.

Enlows (1920, p 47) states, "*H. aerogenes* (Escherich) Miller."

The genus is made a synonym of *Bacillus* by De Toni and Trevisan (1889, p 952). Apparently the name has not been given generic recognition by subsequent authors. The description is scarcely adequate for identification, probably the genus should be regarded as invalid.

Helicomonaden. A casual name used by Klebs (1879, p 161) for the organism which he believed to be the etiologic factor in syphilis. Enlows summarizes his description (1920, p 47) as follows:

A pleomorphic organism, consisting at times of short rods, arranged in more or less spiral form, and of granules, motile. The granules arise from the rods, which toward the end of a spiral series become shorter and shorter, finally appearing as small round bodies. As to their being micrococci he says "Die Möglichkeit dass dem so sei, lässt sich nicht ableugnen, doch wird es in diesem Falle nicht an der Auffindung weiterer Differenzen fehlen, welche uns gestatten werden, ein kurzes Stäbchen, ein Brachibacterion etwa, von einem Coccus zu unterscheiden." He thinks no spores are formed, and that longitudinal division occurs. The granules (körnchen) also form spiral-like masses. The rod form in culture forms "Bakterienballen."

Helicomonas. Enlows (1920, p 47) states

The name *Helicomonas syphiliticum* for the organism Klebs describes as above has been ascribed to Klebs, but in none of his publications have I found it. This name occurs in many German papers on this subject, and in Lipp Med Diet, Philadelphia, 1910, p 411.

Inasmuch as the description of the species is quite inadequate, the genus name has not come into use. Its validity is doubtful. It is rejected (1905) by Erwin F. Smith.

Helikobacterium. Probably to be regarded as a variant of *Helicobacterium*. It was used by Escherich (1886, p. 2) in the combination *Helikobacterium zopfii*. Enlows (1902, p. 47) states

Characterized by its spiral colonies on gelatin plates. As the gelatin liquefies zoogloae of spindle-shape are formed, which anastomose, covering the entire surface of the gelatin, and consisting of "swarming" bacteria, spirilla, and watch-spring-like threads. In older gelatin cultures round and elliptical forms in varied grouping are found (diplococci, tetrads, chains, etc). He figures what he describes as "spirochäten" occurring in a milk culture of his *Helikobacterium*.

Species. Thinks *Bacterium zopfii* Kurth belongs here and suggests the name *H. zopfii* for it. In one paragraph he writes his genus *Helikobacterium* (Klebs), probably referring to *Helicomonaden* Klebs.

Helikomonas. A variant of *Helicomonas* used by German writers as Escherich (1886, p. 2).

Helobacteria. A name applied by Billroth (1874, p. 27) to the growth form of his *Coccobacteria septica* in which sporulation occurred. The genus is discussed and rejected by Cohn (1875, p. 188) and is rejected by Erwin F. Smith (1905). The name has never been used in a strict generic sense.

Hemophilaeae. A name given by Winslow *et al.* (Committee Soc. Am. Bact., 1920, p. 212) to a tribe with the genus *Hemophilus*. The tribal description is "Minute parasitic forms growing only in presence of hemoglobin, ascitic fluid or other body fluids."

The name should probably be spelled *Hemophileae*. The latter spelling is used by Bergey *et al.* (1923, p. 268).

Hemophileae. A corrected spelling of the tribal name *Hemophilaeae*, suggested by Bergey *et al.* (1923, p. 268). Two genera are included. The description follows:

Minute parasitic forms growing only in the presence of hemoglobin, ascitic fluid or other body fluids, or in the presence of certain growth accessory substances found in sterile, unheated plant tissue (potato). Mon-motile. Gram-negative.

Hemophilinae. A name given by Buchanan (1918, p. 44) to the second tribe of the *Bacteriaceae* with the description "Strict parasites, requiring hemoglobin or at least serum for their growth in media. Gram-negative. Non-motile. Cells may be pleomorphic. Usually very small. No spores."

Two genera are included, *Asterococcus* and *Hemophilus*.

Hemophilus. A name used by Winslow *et al.* (1917, p 561) for the fourth genus of the family *Bacteriaceae* with the following description.

Minute rod-shaped cells, non-motile, without spores, strict parasites, growing best (or only) in the presence of hemoglobin, and in general requiring blood serum or ascitic fluid Gram-negative

The type species is *Hemophilus influenzae* (Pfeiffer)

Buchanan (1918, p 44) used the same definition The "Committee" (1920, p 212) added to the diagnosis "sometimes thread forming and pleomorphic."

Castellani and Chalmers (1919, p 933) use the spelling *Haemophilus* *q v*

Bergey *et al* (1923, p 268) includes the genus as the first in the tribe *Hemophyleae* with the description

Minute rod-shaped cells, sometimes thread forming and pleomorphic Non-motile Strict parasites growing best (or only) in the presence of hemoglobin and in general requiring blood serum, ascitic fluid, or certain growth accessory substances Gram-negative

Henrillus A name given by Heller (1922, p 15) to the fifth genus of the *Clostridioideae* with the following description

Clostridioideae that do not liquefy gelatin They produce acid and gas in meat medium They clot milk readily and attack many sugars, producing much acid Gram-negative or gram-positive slender rods with terminal oval spores Their colonies in deep agar are large, lenticular and opaque Very common in soil, often found in wounds do not invade tissue

Type species *H. tertius* (*Bacillus tertius* Henry) as described by Henry Henry believes that the conception *B. tertius* applies to a group and not to a species His nine strains which may be taken as a type split the monoses, disses, mannose, xylose, starch, dextrin, glycogen, salicin, amygdalin and mannitol

This type was described by von Hübner with the number IX Fleming (Bac Y) Rodella (1902) (Bac III), Robertson (1916 a), McIntosh, Adamson (1919) and the Committee and Weinberg and Séguin also describe it Choukévitch (1913) identifies spherical sporulating organisms with Rodella III The original Rodella III did not clot milk and should perhaps be associated with *Fleminigillus*

Hübnerillus. A name proposed by Heller (1922, p 17) for the ninth genus of her tribe *Clostridioideae* The description is

Clostridioideae that do not liquefy gelatin They do not clot milk Large gram-positive rods which form more or less reluctantly long ellipsoid spores, they may form orgonts Colonies in deep agar, small and lenticular or with fine radia-

tions May be pathogenic for the guinea-pig, producing oedema and gas, or paralysis

Type species *H. sextus* (*Bacillus* VI of von Hibler), as described by von Hibler (résumé by Weinberg and Séguin, pl 212)

Von Hibler describes another species which he terms VII These organisms have much in common and we are probably justified in including them in one genus as *Hiblerillus sextus* and *Hiblerillus septimus* The latter resembles an organism described by Tizzoni and Cattani, according to von Hibler There are probably many organisms in soil which are pathogenic for rabbits and guinea-pigs when given certain conditions favorable to invasion, which rarely invade under natural conditions or which on account of shy growth habit are missed when they invade in the company of other organisms

To this genus may be assigned two organisms isolated from the intestine of the horse and described by Choukevitch (1911), they possess similar fermentative ability the production of acid and no gas in glucose agar A non-pathogenic one *Streptobacillus anaerobicus-rectus* may be termed *H. rectus*, another, *H. megalosporus*, produced a fibrino-purulent peritonitis in a guinea-pig

Hillhousia. A generic name proposed by West and Griffiths (1909, p 398) for a giant sulphur organism The following summary taken from their paper satisfactorily characterizes the genus

Hillhousia mirabilis is a sulfur bacterium of giant proportions, and is much the largest solitary bacterium which has so far been discovered Its average length is about 60 μ and breadth about 36 μ

The organism is a peritrichous bacterium with a large number of short cilia It occurs among decaying organic matter in the mud of shallow fresh water pools

Each individual contains a protoplasmic network in the wide meshes of which large globules of sulfur (probably not pure, but in loose combination with proteid material), are located The network included numerous small granules, a considerable proportion of which consist of some nucleo-proteid None of them are chromatin granules

The cell-wall is firm and has great powers of resistance to reagents It is not homogeneous, and 5 per cent carbolic acid demonstrates its lamellose character

The multiplication of the organism is relatively slow, one division occupying upwards of twenty-four hours

The genus is recognized by Buchanan (1918, p 463) as the third in the family *Achromatiaceae* with the following description

Cells very large, 42 to 86 by 20 to 33 μ , motile by means of peritrichous flagella Cells packed with large globules of oily amorphous sulphur

The type species is *Hillhousia mirabilis* West and Griffiths

It is included by Bergey *et al* (1923, p 412) as the fourth genus of *Achromatiaceae* Borsa (1920, p 231) concludes that this is a synonym of *Achromatum* Schewiakoff

Homococcaceae. A subfamily of the family *Coccaceae* created by Fischer (1897, p. 32) to include those cocci in which the plane of fission are in a definite sequence. The following genera were included: *Sarcina*, *Planosarcina*, *Pediococcus* and *Streptococcus*.

The same name was also used by Fischer in 1903, p. 59.

It should be noted that the name is not derived from one of the constituent genera.

Hyalococcus. A generic name proposed by Schroeter (1886, p. 152) with the following generic description.

Zellen kugelig oder elliptisch, einzeln oder zu 2, seltener in Reihen zu 4 und 6 in einfache, weit abstehende, scharf umgrenzte Kapseln eingeschlossen. Two species were included, *H. pneumoniae* (the pneumococcus of Friedländer) and *H. Bergleyi*.

Hansgirk (1888, p. 266) included this as the fifth genus of his subfamily *Cystococcaceae*. It was listed as a synonym of *Chlamydatomus* by De Toni and Trevisan (1889). It was recognized by Ludwig (1892).

The name was rejected by Erwin F. Smith (1905) and by Vuillemin (1913, p. 521).

The fact that *Hyalococcus pneumoniae* is given first would seem to fix it as the type. It is the pneumobacillus of Friedländer (not Fraenkel's pneumococcus). It would seem that this is a valid generic designation for this group of organisms, in spite of its inappropriateness, providing they are worthy of generic recognition. It would seem to have priority over *Encapsulatus* *q.v.* which has the same type species.

Hydrogenomonas. A generic name proposed by Orla-Jensen (1909, p. 311) in the following paragraph:

Die Oxydation des Wasserstoffes

$H_2 + O = H_2O + 69 \text{ Kal}$

Per Gramm 3.83 Kal

Nach Kaserer wird dieser Prozess von einigen monotrichen Kurzstäbchen ausgeführt, die von ausschliesslich anorganischer Nahrung leben können, jedoch auch auf organischem Substrat gedeihen, weshalb er sie *Bacillus pantotrophus* genannt hat. Nach B. Niklewski wächst dieser Organismus nur im Verein mit anderen Bakterien autotroph. Nach meiner Nomenklatur kann man ihm passend den Gattungsnamen *Hydrogenomonas* geben.

It is included as the third genus of his family *Oxydobacteriaceae*.

Niklewski (1910, p. 113) accepted this generic designation and named two species, *H. vitrea* and *H. flava*.

Winslow *et al.* (Committee Soc Am Bact (1917, p 551, 1920, p. 201)) give the following diagnosis

Monotrichic short rods capable of growing in the absence of organic matter, and securing growth energy by the oxidation of hydrogen (forming water) Kaserer (1905) who first described the organism states that his species will also grow well on a variety of organic substances

The type species is *Hydrogenomonas pantotropha* (Kaserer) Jensen Nikleuski (1910) described two additional species, *H. vitrea* and *H. flava*

The genus is included by Bergey *et al* (1923, p 31) as the first in the tribe *Nitrobacteriaceae*

Hygrocrocis A genus of algae created by Agardh (1824) It has been recognized as such by several algologists To it were referred certain bacteria by various writers For example, one of the iron bacteria (*Gallionella ferruginea*) was placed here by Perty (1852, p 215) It is invalid as a bacterial genus

Valentin in 1836, p 110, described an organism from the intestinal tract of the common cockroach *Blatta orientalis* which he named *Hygrocrocis intestinalis* He gives the following description "Fila simplicia, tenuissima, prolunga, articulata, serpentina, apice recta, moniliformia, articulis globosis

Leidy (1851, p 41) states that while he was able to find many entophytes in the intestinal tract of the cockroach, he was unable to find the organism described by Valentin Chatton and Perard (1913, p 1160) noted the name *Hygrocrocis intestinalis* of Valentin As a result of a comparison of material from *Blatta* and that from the *Julus*, these authors conclude that the *Hygrocrocis intestinalis* of Valentin and the *Arthromitus cristatus* of Leidy are of the same genus They accept *Hygrocrocis* as having priority They describe the form as follows

Ce sont de longs filaments à compartiments subgêaux, cubiques, à croissance indéfinie, immobiles, et qui au surplus sont, d'après Leidy, fixés à la paroi intestinale par bouquets sur une sorte de pied commun globuleux Il y a dans les régions sporulées de ces filaments une endospore par cellule et ces cellules porigènes ne sont point hypertrophiées

Nous ne croyons pas pouvoir attribuer au genre *Hygrocrocis* (*Arthromitus*) le parasite des cobayes, qui s'en distingue par son cloisonnement très étroit et irrégulier, sa croissance limitée, le nombre restreint des endospores, la différenciation des cellules qui les contiennent, et enfin sa condition libre et mobile Ces caractères seront ceux du nouveau genre *Oscillospira*

Hypheothrix A genus of algae created by Kutzing (1843) Some of the iron bacteria have been placed here by various writers De Toni and Trevisan (1889) make it in part a synonym of *Detoniella*

Hyphomycetes. A general name frequently used by mycologists to include the molds. It has been used by Lehmann and Neumann (1896, p 107) to include the mold-bacterial genera *Mycobacterium*, *Corynebacterium* and *Oospora* with the following diagnosis

Chlorophyllfreie Fadenpilze teils immer, teils vorwiegend mit echt verzweigtem Mycel ohne Endosporen, z T mit Bildung conidienartiger Sporen Von dieser sehr grossen, durchaus provisorischen Familie haben wir nur 3 Gattungen berücksichtigt, ausschliesslich solche, die durch ihr dünnes Mycel in Fragmenten als Spaltpilze imponieren konnten und z. T allgemein imponiert hatten

Later these authors substituted the name *Actinomycetes* (q v)

Hyphomicrobium. A generic name proposed by Stutzer and Hartleb (1901, p 76) for a species *H vulgare* The description is summarized by Enlows (1920, p 49) as follows

A nitrifying (?) organism found in soil Related to the bacteria and to the hyphomycetes On nitrate agar, small homogenous rods, with usually pointed ends, 0.6 to 0.8 μ by 1 to 1.5 μ long Stained with phenol fuchsin a darker central body surrounded by a clear zone may be observed Egg-shaped forms in older cultures, which send out threads, some of which show true branching Multiplication also by transverse division Found also in cement which they think was decomposing through the assistance of this organism

Hypnococcus. A pseudo-generic name given by Bettencourt, Kopke, de Renzende and Mendes (1904, p 55) to a "diplo-streptococcus" which they isolated from the meningeal fluid of patients suffering from sleeping sickness and believed this to be the causal organism The name is not a true generic designation

Indiella A genus of fungi created by Brumpt (1906 according to Castellani and Chalmers 1919) It is of bacteriological significance because *Indiella somaliensis* Brumpt was later made a synonym of *Nocardia somaliensis* by Chambers and Christopherson (1916)

Indiellopsis. The following summary is given by Enlows (1920, p 49)

According to Chalmers and Christopherson Ann Trop Med and Parasitol, Liverpool, 1915, pp 240-255 Brumpt classified the cause of certain mycetomas of the hand as *Indiella somaliensis* (1906) In 1913 he renamed this species *Discomyces somaliensis*, and in the same year created a new genus or subgenus for it *Indiellopsis somaliensis* (Brumpt, 1906) Brumpt based his *Indiellopsis* on the fact that the species secreted around itself in the grain a hard sheath, insoluble in potash and eau de javelle, which no other *Nocardia* is known to do

Indolococcus A generic name tentatively proposed by Orla-Jensen (1909, p. 340) in the following paragraph

Da die peptonisierenden Milchsäurekokken reichliche Mengen Pepton, aber sehr wenig weitere Erweisszersetzungserzeugnisse, bilden, während die von den Proteus-Bakterien abgeleiteten Kokken durch Indolproduktion charakterisiert sein müssen, so liesse sich vielleicht zwischen den Gattungen *Peptonococcus* und *Indolococcus* unterscheiden

The genus is not included in his outlined classification

Iodococcus. A corrected spelling of *Jodococcus* q v. (Miller 1892, p. 63) used by Migula (1900, p 218) and others

Jodococcus A generic name used by Miller (1888, p 612) to designate certain cocci found in the mouth which stain blue to violet with iodine.

The first organism named by Miller was *Jodococcus magnus* This is a large coccus or diplococcus of variable size (1892)

Das Reinzuchten gelang zuerst auf einem Nährboden, der aus gleichen Theilen Agar-agar Gelatine und einer, bei Zimmertemperatur eben starr werdenden Lösung von Zahnbeinleim zusammengestellt war Er enthielt ausserdem 1 5 Zucker und 1 5 Stärke

Upon this medium an abundant growth quickly developed. The plate was then covered with a weakly acidified iodine potassium iodid solution The medium was colored bluish, and most bacterial colonies yellowish Some, however, show a violet color, and subcultures from these retain this power The reaction shows up best on agar containing sugar

A second species described is *Jodococcus parvus* A third species, *Jodococcus vaginatus* (1892, p 63) is described as follows

Er erscheint in Ketten sind mit einer Scheide versehen, in welcher die Zellen als flache Scheiben oder in mehr abgerundeten, auch viereckigen Formen erscheinen, die zuweilen grosse Aehnlichkeit mit Tetraden besitzen Die Verbände haben einen Durchmesser von 0 75 Mitunter findet man Ketten, in welchen ein oder mehrere Glieder fehlen oder ausgefallen sind, wieder andere, bei denen die Hülse zersprengt worden ist, die Glieder aber ihre Verbindung mit der Kette nicht völlig gelöst haben, die Reste der zersprengten Scheide sind mit unter leicht nachweisbar, dieselbe nimmt keine violette Färbung an, sondern bleibt farblos oder wird nach längerer Einwirkung des Reagens schwach gelblich Der Zelleninhalt färbt sich mit Jod stets tiefblau bis violett

Migula (1900, 218) has attempted to identify Miller's *Jod. vaginatus*, but without success

The species of Miller do not seem to have been recognized by later authors It is rejected by Erwin F Smith (1905)

The first species described, *Jodococcus magnus* may be regarded as the type. The organisms, however, have not been identified. The position of the genus and its nomenclatural status is certain.

Kalymmabacterium. A name used by Beaurefaire-Aragao and Vienna (1912). See *Calymmatobacterium*.

Karphococcus. A variant of the generic name *Carphococcus* (q.v.).

Karphokokkus. A variant spelling of *Carphococcus* q.v.

Keratophyton. A casual name used by Rosenbach (1908, p. 514) to an organism believed to cause corneal ulcers. Enlows (1920, p. 50) gives the following characterization:

Doubtful as to its position—belongs either with the bacteria or with the "schimmelpilzen." Varied form and size. Some of the rods are so short as to appear like cocci, many are somewhat longer, others again (young cultures) form very long, wave-like, at times branched threads. Here and there are thick, spindle-shaped or irregular drawn-out forms, which sometimes are filled with vacuoles. Many of the rods are similar to the bacilli of chicken cholera.

Kladothrix. A German spelling of *Cladothrix*, q.v.

Klebsiella. A genus named by Trevisan (1885, p. 105) with the following description:

Due stadi di sviluppo vegetativo 1 Bacilli 2 Cocchi. Bacilli (stadio protoplastico tipico) cilindrici, dritti, inarticolati, incolori, biforini, macrobacilli e microbacilli, citoplasma egualmente diffuso, Cocchi (stadio finale) derivati da microbacilli, seriatì in monili, apparati solitari. Bacilli e cocchi costantemente in ogni stadio racchiusi in crasse vagine sacchiformi membranaceo-mucose. Spore nulle. Specie unica *Klebsiella crouposa*. *Bacterium Pneumoniae-crouposae* Zopf (Spaltpilze, 3 ediz. pag. 66, fig. 20).

Later (1887, p. 94) he expanded the generic description as follows:

Tre stadi di sviluppo vegetativo 1 Baculi, 2 Filamenti, 3 Cocchi. Baculi (stadio primigeno tipico) cilindrici, inarticolati, incolori, biforini, macrobaculi e microbaculi. Filamenti (stadio anormale transitorio osservato solamente in culture nella camera umida) risultanti da baculo o cocchi rimasti in congiunzione per incompleta temnogenesi. Cocchi segregati (monococchi od ottoformi) o seriatì in catenelle. Baculi e cocchi normalmente racchiusi in crasse capsule membranaceo-mucose. Spore (osservate nella sola *Klebsiella rhinoscleromatis*) provenienti nei baculi o ignote.

The following species are listed: 1 *Klebsiella rhinoscleromatis* (Bacillus of rhinoscleroma of Frisch), 2 *K. salvaris* (Microbe de la salive of Pasteur), 3 *K. pneumoniae*, and 4 *K. aerobia* (capsulated air bacillus of Babes). Trevisan later (1888, p. 787) discussed the genus *Klebsiella* in detail. Sixteen species were described. The

species previously termed *Klebsiella crouposa* or *K pneumoniae* was renamed *K. Friedlanderi*. According to De Toni and Trevisan (1889, p. 1028) the diagnosis is as follows

Baculi recti, tum cylindracei apicibusque rotundatis, tum ellipsoideo-fusi-formes apicibusque attenuatis, in articulati, hyalini, immobiles vel rarius mobiles, plasma aequaliter diffusum Filamenta flexuosa vel recta, cylindracea vel moniliformia, articulata, a baculis temnogenesi transversali imperfecta adhuc conjunctis constituta Cocci globulosi, e baculorum seu filamentorum transformatione orti Baculi, filamenta, cocci intra capsulas membranaceomucosas, raro firmas, inclusi Sporae (endosporae) in baculis provenientes

The generic description does not fit closely any of the species included by De Toni and Trevisan as they are now understood The genus is divided into two species groups "*Species anthrophobiae*," with fourteen species, and "*Species zoobiae*," with three Two species *Klebsiella rhinoscleromatis* and *K salvarius* are credited to "Trevisan 1887" Of the species named in 1889, *K Friedlanderi* (Friedlanders pneumobacillus) and *K enteritidis* are well known This genus name is rejected by Erwin F Smith (1905)

Vuillemin (1913, p 525) has proposed that Trevisan's *Klebsiella salvarius* (*Diplococcus pneumoniae* of Weichselbaum) be constituted the type of a "*Genus conservandum*," with an emended diagnosis However, the generic name *Diplococcus pneumoniae* Weichselbaum has priority, and *Diplococcus* (qv) should be used if the pneumococcus is to constitute the type of the genus

As indicated by Enlows (1920, p 50) the type species is *Klebsiella crouposa* Trevisan

De Toni and Trevisan (1889, p 1029) give the following synonymy for Trevisan's *Klebsiella Friedlanderi*

Hyalococcus pneumoniae Schroeter (1886)
Pneumococcus Friedlander
Bacterium pneumoniae-crouposae Zopf
Klebsiella crouposa Trevisan 1885
Klebsiella pneumoniae Trevisan 1887

It is evident that the correct name for the type species therefore is that of the Friedlander pneumobacillus Its name is probably either *Klebsiella crouposa* or *Klebsiella pneumoniae* From the standpoint of usage the latter is preferable *Klebsiella* has priority over *Hyalococcus* and *Encapsulatus*, both based on the same type species

Klebsielleae A tribe of bacteria proposed by De Toni and Trevisan (1889, p 1028) with the following diagnosis "*Baculi tegumentis speci-*

albus membranaceogelatinosus (capsulis) involuti” The following subtribes are included *Euklebsielleae* (genera *Winogradskya*, *Klebsiella* and *Dicoccia*) and *Myconostocaceae* (genus *Myconostoc*).

Klebsiellee A variant of *Klebsielleae* used by Trevisan (1889, p. 25) with the description “Baculi, filamentie cocchi racchiusi in capsula membranaceo-gelatinose” It constituted the second tribe of *Baculogene* and contained the three genera *Klebsiella*, *Dicoccia* and *Myconostoc*

Kokkaceen. An aberrant spelling of the family *Coccaceae* used by certain German writers (as Hueppe 1886, p. 140)

Kokken. Baumgarten (1890, p. 127) states “1 Gatt *Kokken*, Bacterien arten umfassen d welche nur kugelige Vegetationsformen bilden ”

Usually used as a casual designation in German.

Kokkobacillus. A generic name used by Biedert (1885, p. 439) for the species *K. zymogenus* It is to be regarded as a variant of *Coccobacillus* (q v) *Kokkobacillus ancidus* is listed by Eisenberg (1891, p. 275) as a synonym of *Bacillus cholerae gallinarum*

Kokkobacteria. A variant spelling of *Coccobacteria* used by Klebs (1887, p. 50)

Kokkobacterium. A variant spelling of *Coccobacterium*, q v The name *Kokkobacterium Lyssae* is used in a review of an article by Rivolta in *Deutsche med Wochenschr* (1886, p. 313).

Kokkothrix A German variant of *Coccothrix* (Unna 1887, p. 29)

Kokkus A spelling of *Coccus* or *coccus* used by certain German authors For discussion see *Coccus*

Kurthia A genus established by Trevisan (1885, p. 92) to include *Kurthia zopfii*, the *Bacterium zopfii* of Kurth The original description follows

Tre stadi di sviluppo vegetativo 1 Filamenti 2 Bacilli 3 Cocchi
Filamenti (stadio protoplastico tipico) cilindrici, dapprima apparentemente inarticolati, dappoi articolati, diritti, uniformi, agglomerati in colonie irregolarmente arrotondate Cocchi (stadio finale dapprima appaiati, indi liberi Spore nulle od ignote Specie unica *Kurthia Zopfii*

It is noted as the fifth genus of *Leptotrichae* by Trevisan (1889, p. 11) with the description “Vagine nulle”

Later the organism named *Bact. Laminariae* of Billet was included The generic diagnosis as given by De Toni and Trevisan (1889, p. 931) is as follows

Filamenta cylindrica, vagina delicatissima evanescente obducta, simplicia, utrinque apicibus conformia, primitus apparenter subcontinua, cito articulata, articulis aetate provecta, vagina evanida, in baculos et coccus soluta. Multiplicatio baculis primitus vivide mobilibus, propere immotis. Arthrospora transformatione cocci singuli ortae. Obs. Arthrospore in parvulos acervos zoogloeicos conglobantur in quibus nova filamenta radiatur exoriuntur.

The genus is rejected by Erwin F. Smith (1905).

Bact. zopfii Kurth may be regarded as the type.

Apparently Wenner and Rettger were not aware of the use of *Kurthia* with the species *K. zopfii* as the type, as this has priority over *Zopfius qv* which has the same specific type.

Kurthiaeae. A tribe created by De Toni and Trevisan (1889, p. 929) with the following diagnosis: "Sporae (arthrospora) in filamentis normalibus obvenientes. Filamenta simplicia, basi ab apice superiori ut plurimum distincta, vaginata vel evaginata."

The following genera are included, *Detoniella*, *Rasmussenia*, and *Kurthia*.

Lactobacillaceae. A family named by Winslow *et al.* (Committee Soc. Am. Bact., 1917, p. 561) with the following description:

Rods, often long and slender, Gram-positive, non-motile, without endospores. Usually produce acid from carbohydrates, as a rule lactic. When gas is formed, it is CO₂ without H₂. The organisms are usually somewhat thermophilic. As a rule microaerophilic, surface growth on media poor.

The genus *Lactobacillus* was included.

Lactobacillaeae. A variant spelling of *Lactobacilleae* used by Bergey *et al.* (1923, p. 241). It is included as the eighth tribe of the family *Bacteriaceae*. The description follows:

Rods, often long and slender. Gram-positive. Non-motile. Without endospores. Usually produce acid from carbohydrates, as a rule lactic. When gas is formed it is CO₂ without H₂. The organisms are usually somewhat thermophilic. As a rule, microaerophilic. Surface growth on media is poor.

Lactobacilleae. A tribe named by Winslow *et al.* (1920, p. 211).

Rods, often long and slender, Gram-positive, non-motile, without endospores. Usually produce acid from carbohydrates, as a rule lactic. When gas is formed, it is CO₂ without H₂. The organisms are usually somewhat thermophilic. As a rule microaerophilic, surface growth on media poor.

It includes the genus *Lactobacillus*.

Lactobacillus. A genus of bacteria proposed by Beijerinck (1901, p. 214) to include all rod-shaped forms "qui produisent l'acide lactique".

lévogyre" It differs from the genus *Aerobacter* in that no hydrogen is produced Included in this genus are those forms which produce lactic acid at temperatures above 30°C and which are responsible for the fermentations of Kefyr, of Koumys and of the Matzoon. The acidification begun by the *Lactococcus* is continued in milk by *Lactobacillus*. These organisms are likewise responsible for the souring of ensilage, and constitute the so-called acidophile flora of the intestines of infants. The species *Lactobacillus fermentum*, *L. delbruckii* and *L. caucasicus* are listed, the two former from fermentation industries, the latter from Kefyr

Bejerinck says (1901, p 213) that *Lactobacillus* and *Lactococcus* "sont certainement des genres phylogénétiques et nul point des 'genres physiologiques' comme par exemple *Photobacter*, ou simplement morphologiques comme les 'genres' *Bacillus*, *Pseudomonas*, *Micrococcus*, etc" Later he diagnoses the genus as follows

Au genre *Lactobacillus* doivent être rapportés les ferments lactiques actifs de forme bacillaire qui produisant l'acide actif levogyre et qui se rapprochent du genre *Aerobacter* dont j'ai donné la diagnose en 1900 Pourtant *Lactobacillus* se distingue de ce genre en ce que la production l'hydrogène, caractère important pour le diagnostic d'*Aerobacter* y manque absolument, ainsi que dans tous les autres ferments lactiques actifs

Bejerinck (1908, p 370) discussed the characters which differentiate *Lactobacillus caucasicus* and *L. longus*

The type species is somewhat difficult to determine Bejerinck (1901) states that the question of species is troublesome, then

Passant sous silence pour le moment les *L. fermentum* et *L. Delbruckii*, producteurs du levain lactique, nous rencontrons en premier lieu le *Lactobacillus caucasicus*, dont les zooglyphes sont bien connues sous le nom de grains de Kefyr

He then describes the last named species It would seem that although *L. fermentum* is first mentioned, *L. caucasicus* may be regarded as the type

Winslow *et al* (Committee Soc Am Bact 1917, p 561) state the generic characters are those of the family *Lactobacillaceae* and the type species is *L. caucasicus* (Kern?) Bejerinck

Buchanan (1918, p 54) states

Rod-shaped organisms, cells frequently quite elongate, non-motile, without spores, Gram-positive in young cultures Produce acid, largely lactic, from carbohydrates When gas is produced, it is CO₂ without hydrogen. For the most part the organisms are thermophilic Microaerophilic

The type species is *Lactobacillus caucasicus* (Kern) Bejerinck

Bergey *et al.* (1923, p 241) include this as the single genus of the tribe *Lactobacillae*

Lactobacter. A name applied by Beijerinck (1900, p 200) to include all lactic acid bacteria, both rods and cocci. It may be regarded as a physiological genus. It has been used only as a casual name. It is rejected by Erwin F. Smith (1905).

Lactobacteriaceae. A family name proposed by Orla-Jensen (1921, p 271) in the following statement

Although it is well-known that the rod as well as the colorless sulphur bacteria may appear in all the forms known in the world of bacteria, and that even a single species of bacteria (for instance, *Crenothrix* and *Azotobacter*) in the first state of development may only divide in one plane but later in more planes, yet the Committee cannot admit that in case of other bacteria there may be a near relationship between sphere-, rod- and screw-forms. Still I entertain a perhaps not unwarranted hope, that my recently published monograph of the lactic acid bacteria may be able to change the opinion of the Committee. Here we have to do with a large group of bacteria consisting of sphere- as well as rod-forms, nevertheless forming a natural family which we could call *Lactobacteriaceae*. This family I have founded, of course, not only on the specially developed power of forming lactic acid (since there exist many different organisms which are able to form, at any rate, small quantities of lactic acid), but I base it upon the fact that the bacteria which we call true lactic acid bacteria have so many other properties in common that there cannot be any doubt about their close relationship. Thus, they are Gram-positive, facultatively anaerobic (without surface growth in stab culture), they make excessively great demands as to nitrogenous nutriment, and, most remarkably, in contradistinction to most other bacteria, they are unable to liberate oxygen from peroxide of hydrogen.

Lactococcus. A genus proposed by Beijerinck (1901, p 213). He emphasizes that this genus and *Lactobacillus* are natural genera. He states.

Les ferments actifs (lactique) appartiennent à deux genres naturels, *Lactobacillus* et *Lactococcus*, qui se distinguent nettement aussi bien par la forme que par leurs qualités physiologiques et qui, d'après ma conception sont certainement des genres phylogénétiques et non point des genres physiologiques, comme par exemple *Photobacter*, ou simplement morphologique comme les genres *Bacillus*, *Pseudomonas*, *Micrococcus*, etc., qui tous contiennent des formes bien éloignées dans le système naturel.

The genus includes those cocci, both diplococci and streptococci which at temperatures below 30°C produce the lactic fermentation of milk, and which commonly produce dextrorotatory lactic acid. The most common species, the form which may be regarded as the type, is *Lactococcus lactis*, the *Bacterium lactis acidum* of Leichmann (1896, p .

777) A second species is *Lactococcus hollandiae* Weigmann. The genus is discussed by Stockhausen (1907 p 164) Beijerinck later (1908, p 641) named a third species, *Lactococcus agglutinans*, a form previously described by Barendrecht as *Leuconostoc agglutinans* (1901, p 623) The genus is also recognized by Kossowicz (1912, p 94). It is probable that this genus is to be regarded as a synonym of *Streptococcus*

The type species is *Lactococcus lactis* Beijerinck If the milk cocci are to be separated from the pathogenic forms of the genus *Streptococcus* this would appear to be a valid genus

Lactosarcina. A generic name used by Beijerinck (1908, p. 359) for sarcinae which produce lactic acid He concluded that the *Sarcina ventriculi* produces sufficient lactic acid in milk to justify its inclusion in this genus

This name has apparently not been used by other writers, and is probably to be regarded as a synonym of *Sarcina* Goodsir

Lampozoidia The subdivision of the protozoa created by Perty (1852, p 179) to include the form now known as bacteria (among others) The definition of the group is as follows

Farblos seltener blau, gelb, rothlich (nicht grün) gefärbt ohne spezifische Organe, kaum mit einer Spur von Differenzirung ihrer Substanz Bewegung scheinbar willkürlich, in Wahrheit automatisch Vermehren sich durch Quertheilung und stellen so Ketten und Fäden dar

Lamprocystaceae A subfamily of the sulphur bacteria named by Migula (1900, p 1043) The diagnosis is as follows "Zellen zu Familien vereinigt Teilung der Zellen zuerst nach 3, dann nach 2 Richtungen des Raumes" This subfamily contains the single genus *Lamprocystis* The subfamily is again recognized by Migula (1904, p 146) by Erwin F Smith (1905, p 163) and by Frost (1911, p 61)

Lamprocysteae A name proposed by Buchanan (1918, p 467) for a tribe of sulphur bacteria containing the single genus *Lamprocystis* The definition is "Cells united into families in which division of the cells occur first in three planes, then in two"

This designation is used by Bergey *et al* (1923, p 297) in the same sense

Lamprocystis. A generic name proposed by Schroeter (1886, p 151) to replace the invalid *Cohnia* Winter (*qv*) He gives the following generic description

Zellen elliptisch, anfangs zu rundlichen Zellhaufen zusammengeballt, später hohle Sacke bildend, in welchen die Zellen einschichtig einer Schleimmasse eingebettet lagern, zuletzt zerreißt die Haut stellenweise und bildet ein Netz Zellinhalt rothlich Von der Spaltalgengattung *Clathrocystis* Henfrey nur durch den Mangel des Phycochroms verschieden

One species is included, *Lamprocystis roseo-persicina*. Winogradsky (1888, p 71) described the genus in somewhat greater detail He states

Ihre Zellen sind nach der Theilung kugelig oder kaum merklich elliptisch, mit einem Durchmesser von 2 μ , vor der Theilung fast doppelt so lang Die Färbung ist nur in der Masse ziemlich intensiv violett Die sehr kleinen, punktförmigen Schwefelkörnchen verleihen der Zoogloea einen braunlichen Ton Die Intensität der S Einlagerung und Oxydation ist viel geringer als bei *Thiocystis*

De Toni and Trevisan (1889, p 1038) give the following diagnosis

Cocci globosi vel subelliptici, numerosissimi, in muco matricali nidulantes, in familias globosas magnas inordinate cumulatı Glebae singulae cystidibus universalibus gelatinosis involutae, initio globoso-ovideae, nitus solidae, dein cavo-saccatae, aetate proventa figuratae, irregulariter clathratae, demum in parvos cumulos libere natantes solutae Cystides speciales nullae Coccorum divisio initio glebarum in tres directiones, dein in duas Sporae ignotae

The generic description most frequently used is that of Migula (1900, p 1043) "Familien anfangs solid, darvor hohlkugelig, netzformig durchbrochen, endlich in kleine, schwarmfähige Gruppen sich auflösend" He includes the genus in the *Rhodobacteriaceae* This description has been used by Smith (1905, p 163) and Frost (1911, p 61). The generic name is also used by Orla-Jensen (1909, p 334) for the eighth genus of the family *Rhodobacteriaceae* of the order *Cephalotrichinae*

Buchanan (1918, p 467) gave the following generic description

Cells ellipsoidal, dividing at first in three planes to form spherical cell masses, later in two planes, forming hollow sacks in which the cells lie embedded in a layer in the walls, finally the membrane ruptures, and the whole mass becomes net like, much as in the algal genus *Clathrocystis* Usually colored intensely violet Small sulphur granules present Capable of swarming

The type species is *Lamprocystis roseo-persicina* (Cohn) Schroeter

Bergey *et al.* (1923, p. 387) follow Buchanan, including this genus in the tribe *Lamprocysteeae*

Lampropedia. A genus of cocci proposed by Schroeter (1886, p. 151) with the following diagnosis

Zellen zu 4 oder mehr zu regelmässigen flachen Tafelchen vereinigt, farblos oder lebhaft (nicht grün) gefärbt Von der durch Phycocchrom gefärbten Spaltalgen Gattung *Merismopedia* Meyen nur durch den Mangel des spangrünen Pigments verschieden

One species is included, *L. hyalina* (Kützing) Schrioter

The genus is included in *Thopedia* (g v) by Winogradsky (1888, p 65) De Toni and Trevisan (1889, p 1049) give *Erythroconis* Oeist (1842) as a synonym Their description of the genus is as follows "*Cocci quaternati in explanato, ambitu parallelogrammico, obvolutas late consociati Coccorum divisio in duas directiones*" They divide the genus into two sections, one containing sulphur granules, and one not containing them, with four species in the first and two in the second section Vuillemin (1913, p 520) regards this genus as a synonym of *Merista* Van Tieghem 1883

Buchanan (1918, p 468) included this as the first genus of the tribe *Thopedieae* with the description

Cells united into tetrads, forming flat tubular masses Contain sulphur granules and bacteriopurpurin

The type species is *Lamproedia hyalina* (Kützing) Schroeter

Most recent writers have followed Winogradsky and Migula in disregarding the genus It is rejected by Erwin F Smith (1905)

Lankoides A name given by Castellani and Chalmers (1919, p 934) to the ninth genus of the family *Ebertheae* Their description is

Ebertheae fermenting glucose partially with the production of acid, but no gas, lactose not fermented or only partially, without gas production Milk clotted

Type species —*Lankoides pyogenes* (Passet, 1902)

Later (1920, p 606) five species are included

Leptomitus A generic name used by Agardh (1824, p 83) to include "algae" of various types which grew in certain solutions The generic description given by Kützing (1843, p 154) reads

Trichoma articulatum, in apicem attenuatum, ramosum, articuloaginati Spermatia lateralia, epispermio pellucido cincta Ausser den angeführten Merkmalen weichen die meisten Arten dieser Gattung auch noch dadurch von *Hygrocrocis* ab, dass ihre Zellen hohl und ihre Zelleninhalt häufig vielkörnig (polygonisch) ist, er wird von Jodinctur, die auch hier bei der Untersuchung gute Dienste leistet, braun gefärbt Die ersten Arten schliessen sich sehr an die letzten von *Hygrocrocis* an

Three species were described at this time. Later (1849, p 154-157) thirty species were included, some of them doubtless being mold mycelia, others yeasts and probably some bacteria.

De Toni and Trevisan (1889, p 933) regarded one species, *L. divergens* as a bacterial form and named it *Leptotrichia rigidula*.

This is not a valid bacterial genus.

Leptonema. A genus of algae named by Rabenhorst (1857). It is invalid because of prior use in botany for other forms (Jussieu, 1824, Hooker 1844) according to De Toni and Trevisan who (1889, p 932) give *Leptonema nvea* Rabenhorst as a synonym of *Leptotrichea nvea* Trevisan.

Leptospira. A generic name proposed by Noguchi (1917, p 755). Enlows (1920, p 51) summarizes as follows:

Type species (monotypy) *L. icterohaemorrhagiae* (Inada and Ido, 1914). Cause of infectious jaundice. Closely wound, 10 to 12 coils within 5μ , slender, cylindrical filaments with gradually tapering ends. Lengths 7 to 14μ , rarely 30 to 40μ , diam 0.25 to 0.3μ . Spiral amplitude, 0.45 to 0.5μ . Spiral depth, 0.3μ regular. One or more gentle wavy curves throughout the entire length. In a free space one or both ends may be semicircularly hooked, while in semisolid media the organism appears serpentine, waved or bent. Flexible. No axial filament present, no chambered structure, no membrane, no crista, no flagellum, no terminal finely spiral filament, terminal or caudal (last 6 or 8 spirals) portion highly motile. Division transverse. Stains reddish violet by Giemsa's solution. Also places here *Spirochaeta biflexa* Wolbach and Binger. Noguchi considers this genus intermediate between the protozoa and bacteria. He later included the cause of yellow fever under this genus *L. icteroides* Noguchi. See J. Exp. med. v 29, 1919.

Bergey *et al* (1923, p 429) include this as the sixth genus of the family *Spirochaetaceae* with the following description:

Parasitic forms. Sharply twisted cylinders with flagelliform tapering ends one extremity being sharply curved into a "hook."

The type species is *Leptospira icterohaemorrhagica* (Inada and Ido) Noguchi.

Leptothrichaceae. See *Leptotrichaceae*. The third family of the order *Desmobacteria* proposed by Hansgirk (1888, p 230). Two subfamilies are included, *Leptothricheae* and *Beggratoeae*. No description is given.

Leptothricheae. A family name first used by Kuetzing (1843, p. 197) with the following diagnosis: "*Trichomata tranquilla tenuissima, continua, (vel obsolete articulata) Cellulae propagatoriae propriae nullae*." As a subfamily this name was also used by Rabenhorst (1865, p 73) with the following diagnosis:

Trichomata aut libera aut adhaerentia, simplicia, plerumque distincte articulata et vaginata, rarissime oscillantia, singula vel fasciculata, plerumque in stratum compactum saepe longe lateque diffusum, tenue (VIX lineam crassum) vel superimpositione ad pollicem usque incrassatum, spongiosum, interdum stratosum zonatum et varie coloratum dense intracata Propagatio adhuc ignota

The following genera are included *Leptothrix* Kuetzing, and *Hyphothrix* Kuetzing

This name is the equivalent of *Leptotricheae* (q v) Robin (1853, p 345) used this family name with the description given by Kuetzing, with the following addition "Filaments tubuleux, déliés, continus, sans articulations, privés de mouvement, remplis d'un contenu (endochrome) confluent, indistinct Cellules propagatrices speciales nulles ou inconnues" It was also used by Hansgug (1888, p 230) as the first subfamily of *Leptothrichaceae* including the single genus *Leptothrix* Kuetzing

Leptothricheen A family name used by Zopf (1884, p 45) with the diagnosis "Sie besitzen Coccen-, Stäbchen-, Fadenformen (welche einen Gegensatz von Basis und Spitze zeigen), und Schraubenformen" He included the genera *Leptothrix*, *Beggiatoa*, *Crenothrix* and *Phragmidiothrix* Hueppe uses the same term (1891, p 30) with the description "Leptothricheen bilden im vegetativen Stadium Stäbchen-welche meist längere Fäden bilden" Baumgarten (1890 p 127) defines the term as a genus (Gattung), following Zopf in his definition, and terming Zopf's genera "Untergattungen" It may be regarded as the equivalent of *Leptothrichaceae* and *Leptotrichaceae* (q v)

Leptothrix. This genus was proposed by Kützing (1843, p 198) for a group of algal species with the following description "Trichomata simplicia tenuissima monogononica, turgida, continua, vel obsolete articulata, in stratum vel caespitosum vel compactum, continuum, plerumque late expansum complicata" Four species are included, *Leptothrix ochracea* (*Lyngbya ochracea* Leibl), *L. aeniginea* *L. lutescens* (*Hygrocrocis ochracea* Ag, *Calothrix lutescens* Meneghini) and *L. fontana* (*Hygrocrocis olivacea* Kg *Bangia tenuis* Kg) The first species described (*L. ochracea*) may be taken as the type Its description is as follows

L. fluctuans, natans, ochraceae, trichomatibus curvatis intricatis, subtilissimis (diam 1/1500-1/1200'''), articulis globosis vel oblongis In Eisenquellen, besonders auf sumpfigen Wiesen und in seichten Gräben

The next reference to *Leptothrix* is for an entirely distinct type Robin (1847, p 345) according to Winter (1884, p 56), De Toni and Trevisan (1889, p 930) and Schroeter (1886, p 171) described an organism from the teeth of a horse, which he termed *Leptothrix buccalis*. An examination of this reference shows the organism to be referred to (pp. 42-46) as "Algu filiforme de la bouche," but the name *Leptothrix buccalis* does not appear. Later Robin (1853, 345) gave the following generic description of *Leptothrix* and specific description of *L. buccalis*:

Genre *Leptothrix*, Kutz, Trichomata tenuissima cramosa, nec concreta. Filaments non rameux, ni engainés, ni cohérents. Trichomatibus rigidulis, linearibus rectis vel inflexis, non moniliformibus, achromaticis, extremitatibus obtusis, basi in stromate amorpho granuloso adhaerentibus. Long 0^{mm},020-0^{mm},100, lat 0^{mm},1005. Hab. In superficie linguae, intervallis dentium, cavo dentium corruptorum, et in succis stomachi et intestini. Filaments assez roides, droits ou courbés, quelquefois coudés brusquement, à angle généralement obtus à bords nets, non moniliformes, extrémités non effilées, larges de 0^{mm},0005 et longs de 0^{mm},020 à 0^{mm},100, incolores, élastiques, réunis généralement, par la base, à une gongue amorphe granuleuse, et formant des faisceaux plus ou moins serrés, à moins qu'ils n'aient été détachés (pl I, fig 1 et 2). Hab., surface de la langue, matières accumulées dans les interstices des dents, ou la cavité des dents cariées, d'où ils passent dans les liquides de l'estomac, ou de l'intestin dans les cas de diarrhée.

Robin also described a *Leptothrix insectorum*, from the rectums of various insects, particularly those living in water. The generic description given to fit this form by Trevisan (1879, p 147) is "Somatia cylindrica, plus minus distincte articulata, tenuis, elongata, filiformia, recta, laxa fasciculata."

Kuetzing in 1847 (p 220) published nine species of *Leptothrix*, all of them water and algal forms. Two years later (1849, p 262) he listed and described a total of thirty-seven species. His generic description reads "*Trichomata tenuissima cramosa nec vaginata, nec concreta*." Eight species are grouped in the first division of the genus under the heading "*Species basi adnatae*." All of them are water algae, usually colored, but in some species colorless. Under the heading "*Species trichomatibus liberis, vel intracatis et stratum formatibus*" the other twenty-nine species are described. The first of these is *L. ochracea*. All species are water forms, and most are greenish in color. The *L. buccalis* is not recognized.

Rabenhorst (1865, p 73) includes this genus among the *Nemato-genae* with the description

Trichomata tenuissima, abbreviata, adhaerentia, segregata vel laevae aggregata, nunquam intricata, saepius oscillantia, indistincte articulata et saepe distincta vaginata, cytoplasmata homogenum, aetate protracta granulosum nonnunquam fasciatum contractum

Billroth (1874, p. 15) used the term to designate one of the growth forms of his pleomorphic *Coccobacteria septica* in which the organism occurred as a slender filament without visible articulations

Cohn (1875) included this genus in his tribe *Nematogenes* to include those organisms which produce slender filaments, in which the articulations are indistinct, colorless, cylindrical, free and unbranched.

Trevisan (1879, p. 147) concluded that the genus *Leptothrix* is strictly algal, and created the new genus *Leptotrichia* (q v) with the single species *L. buccalis*

Luerssen (1879, p. 24) accepts both the free living water forms and the parasitic types as members of this genus. He says

Von der Gattung *Bacillus* nur durch die sehr langen dünnen undeutlich gegliederten Fäden verschieden. Im Wasser auf anderen Algen und Wasserpflanzen u. s. w. bei gewissen Krankheitsprozessen auch am menschlichen Körper beobachtet. *L. buccalis* Rob. Auf dem Epithelium der menschlichen Mundhöhle, im Zahnen

Magnin (1880, p. 90) emphasized that *Leptothrix* differs from *Bacillus* in that the filaments are long, adherent, very slender and indistinctly articulated

Winter (1880, p. 56) includes two species only in his genus, *Leptothrix buccalis* and *L. parasitica*. His generic description is "Fäden sehr lang und dünn, unverzweigten scheinbar ungegliedert, farblos, ohne Bewegung, ohne Körnchen, frei oder verfilzt". He states that the fungi which have been assigned to the genus *Leptothrix* are of very questionable specific value, and that the most of the true species of *Leptothrix* are typical algae containing phycochrome, implying that another designation should be formed for the organisms which are colorless and are grouped with the bacteria.

Grove's statement (1884, p. 34) is essentially a translation of that of Winter

Zopf (1885, p. 95) states "Fäden bescheidet oder unbescheidet. Theilungen sehr weit gehend. Zellen schwefellos"

From this time the term *Leptothrix* is frequently found used as a morphological designation for slender unbranched long rods. For example, Flugge states (1886) that "formerly" the bacilli were subdivided to include the true bacillus and leptothrix or very long rods

Schroter (1886, p 170) used the following generic description: "Faden sehr zart, mit dünnen Scheiden, unbeweglich, Gliederung anfangs undeutlich Vermehrung durch Zeifall des Fadens in Glieder." He includes three species only, *L. ochracea*, *L. parasitica* and *L. buccalis*. It may be noted that this definitely excludes from the genus the more strictly algal forms He furthermore emphasizes a possible relationship of *L. ochracea* to *Gallionella ferruginea*

Hansgurg (1888, p 230) recognized this as the only genus of his subfamily *Leptothrichaceae* He described one new species, *Leptothrix cellaris* and listed two others, *L. ochracea* and *L. Thurethiana*.

De Toni and Trevisan (1889, p 930) removed *L. buccalis* to a new genus *Rasmussenia* to include five species, all of them forms from the mouth and related to *L. buccalis* The genus *Leptotrarcha* was reserved for some of them excluding *L. ochracea* This species together with two others are included in still another genus, *Detoniella*

Billet (1890, p 23) says "Leptothrix, élément filamenteux dont la longueur depasse 10 fois la largeur"

Hueppe (1891, p 30) states "Leptothrix unterscheidet sich von den arthrosporen Bakteriaceen dadurch, dass die Faden einem Gegensatz von Basis und Spitze zeigen"

Leptothrix according to Ludwig (1892) includes bacteria which occur in long filaments, are unbranched, are without spores and are non-motile

The tendency of those interested primarily in medical bacteriology to limit the term *Leptothrix* to filamentous bacteria from the mouth is well shown in the following statement from Macé (1897, p 1020)

Le genre *Leptothrix* n'est peut-etre pas à distinguer due genre *Bacillus* Beaucoup de Bacilles, en effet, possèdent la propriété, dans des conditions particulières de milieu, de croître en très longs filaments dont la composition en batonnets n'est visible que lorsqu'on les traite par des reactifs coagulants ou colorants Or, c'est ce qui arrive pour le seul *Leptothrix* dont la morphologie est un peu connue, celui de la bouche Plusieurs espèces de Bactéries en batonnets, habitant normalement la cavité buccale, peuvent peut-être contribuer, à cause des conditions de milieu, à donner ces touffes de longs filaments, on s'expliquerait alors facilement la divergence des caractères exposes par les divers auteurs, surtout la différence des dimensions

Chester (1901, p 369) gives the following generic description.

Filaments unbranched, non-motile, enclosed in very delicate or rather thick sheaths, either fixed or associated in slimy masses The separation of the filaments and the presence of a sheath is demonstrated by special methods of stain-

ing The contents of the filaments become segmented, forming round or ovoid gonidia, which escape from the sheath and develop into filaments Gonidia non-motile

In this he is followed by Allen J Smith (1902, p. 272). Erwin I Smith (1905) rejects the name as a generic designation.

Petruschky (1913, p. 270) includes this as one of the genera of the trichobacteria which never shows branching, is never wavy, with stiff filaments in which stages of division are to be recognized with difficulty

Ellis (1907, p. 515) uses the *L. ochracea* as the type of the genus, defining the genus as consisting of organisms with sharply defined stiff membranes Following are some of the principal characteristics of the organism Usually associated with *Gallionella ferruginea*, over which it preponderates to the extent of about 20 to 1. Organism consists of a number of straight filaments free at both ends Ends often unsymmetrical Membrane very sharply contoured, internally as well as externally Breadth varies from $1\frac{1}{2}$ to 2μ , but when covered with ferric hydroxide often reaches 3μ and more Length reaches up to 200μ and possibly more Formation of conidia takes place by means of a process of budding, constriction taking place as soon as the requisite length has been obtained Sometimes the constriction is prolonged so that a number of quill-like structures are seen protruding from the organism Eventually these also are abstricted and elongate to form new threads The conidia are oval in shape and measure about 1μ in breadth and about $1\frac{1}{4}\mu$ in length Each has a single membrane Multiplication also takes place by cell-division At various unequal distances along both sides of the membranes, small nodules are formed Each nodule divides into two, the split taking place between the two daughter-nodules As the pairs of nodules are not exactly opposite the pairs on the other side of the membrane the daughter-cells are not symmetrical at the ends The cell- and conidium-contents are perfectly homogeneous In no stage of its life-history is any motility observable

Jordan (1908) includes in the genus those trichobacteria with unbranched filaments

Schwers (1908) used the name *Leptothrix ochracea*

West (1904, p. 334) lists *L. ochracea* as a synonym of *Lyngbya ochracea* among the *Myrophyceae*

Orla-Jensen (1909, p. 331) includes *Leptothrix* as the fifth genus of his *Trichobacteriaceae* in which the cells are unattached, not spiral,

unbranched, without sulphur granules and in filaments. He concludes that *Gallionella* and *Spirophyllum* are growth forms of this genus.

Heim (1911, p. 254) again emphasizes the *L. buccalis* type. He says

Diese in der Mundhöhle anzutreffenden Mikro-organismen sind steife, unverzweigte, fadenähnliche Gebilde, die sich mit Jodlösung gelb färben. Es gibt auch fadenartige Mikroorganismen, die sich mit Jod violett färben, sie sind bei Munderkrankungen wiederholt in kornerartigen Gebilden gefunden worden, die in ihrem Aussehen an Aktinomyzeskörner erinnerten. Genauerer liess sich über sie nicht ermitteln, weil ihre Zuchtung bis jetzt nicht gelungen ist.

The legitimacy of *L. ochracea* as a species name is disputed by Mohs, who reduces this designation to a synonym of *Chlamydothrix*.

In a careful study of the organisms of the teeth Kligler (1915, p. 328) cultivated fifteen strains of *Leptothrix buccalis*, and used this designation. His characterization is as follows:

Morphology. A thick, long, straight, or curved thread with a club-head at one extremity and a tapering end at the other. It is generally 0.8-1 μ thick and upwards of 10 μ in length. **Chromology.** It stains readily with anilin dyes in young cultures. In older cultures it has the appearance of a faintly stained sheath enclosing a number of heavily stained granules. Young cultures are Gram-positive. In older cultures the sheath is decolorized while the granules retain the stain. It is not pleomorphic but the threads fragment very early into short, thick rods. Coccoidal forms are not seen. **Biological properties.** They are anaerobic, facultative-aerobic, non-motile, non-branching threads. They grow at 37°C and practically not at all at 20°C. **Cultural character.** No growth obtained on agar. On glucose-agar-plates they give minute pin-point colonies after 3 to 4 days' incubation. Examined microscopically with a low power, they have a dark center with hairy outgrowth. They grow fairly well in serum-glucose-agar, best in the stab and only sparsely on the surface. The surface colony is raised, round, whitish, lustrous, rarely more than 0.5 mm in diameter. The addition of salivary mucinate to ordinary agar renders the latter a very favorable medium, a thin, grayish, white, spreading growth being obtained in 24 hours. Glucose, sucrose and maltose are fermented, but lactose is not attacked. Gelatin is not liquefied and neither indol nor ammonia is produced.

It is evident from the preceding discussion that the name *Leptothrix* has been used in four different ways:

- 1 *Leptothrix* A genus of algae which may or may not include *L. ochracea*
2. *Leptothrix* A genus of iron bacteria with type *L. ochracea*
- 3 *Leptothrix* A genus of elongate bacteria, principally from the mouth, with *L. buccalis* as the type
- 4 *Leptothrix* Any elongate, rod-shaped organism

It would seem logical to make the first named species, *L. ochracea*, the type. This would limit the genus to the iron bacteria. The mouth forms may well be placed in the genus *Leptotrichia* in accordance with the proposal of Trevisan.

This was done by Buchanan (1918, p. 303) in the description:

Filaments of cylindric colorless cells, with a sheath at first thin and colorless, later thicker, yellow or brown, becoming encrusted with iron oxide. The iron may be dissolved by dilute acid, whereupon the inner cells show up well. Multiplication is through the division and abstriction of cells and motile cylindric swarm cells. Swarm cells sometimes germinate in the sheath giving appearance of branching. Pseudodichotomous branching may occur.

The type species is *Leptothrix ochracea* (Leiblein) Kuetzing.

Castellani and Chalmers (1919, p. 1068) give the following description of the genus and designation of type species.

Filamentous fungi with long, very thin mycelial threads, with no capsule or only a very delicate one, non-branching, non-septate, generally non-cultivable.

Type species *Leptothrix maxima* Miller.

Some five species are listed, all parasitic.

Lieske (1921, p. 2) regards *L. ochracea* as the generic type.

Bergey *et al.* (1923, p. 390) follow Buchanan including the genus as the first in the family *Chlamydobacteriaceae*.

Leptotrichacei. A family name used by Schroeter (1886, p. 170) for Zopf's *Leptotricheen* (*qv*). The following genera are included: *Leptothrix* Kuetzing, *Beggiatoa* Trevisan and *Crenothrix* Cohn.

Leptotrichaceae. The corrected form of *Leptotrichacei* used by an abstracter (Journ. Roy. Mic. Soc., 1890, p. 758) of the work of Hansgirg (1890).

Leptotricheae. A family name given by Kuetzing (1843, p. 198) to a group of algae including the genus *Leptothrix*. Later it was used as a "generic" name by Sternberg (1892, p. 19) with the description:

The vegetative cells present rod-shaped and spiral forms, and grow out into straight, wavy or spiral filaments, these may show a difference between the two extremities of base and apex. Cocci-like reproductive bodies are formed by segmentation of the rod-shaped elements in these filaments. In some of the species the segments are enclosed in a common sheath. Subgenera, *Leptothrix*, *Beggiatoa*, *Crenothrix*, *Phragmidiothrix*.

Leptotricheen. A family name used by Zopf (1885, p. 51) with the following diagnosis:

Sie besitzen Coccen-, Stabchen-, Fadenformen (welche einen Gegensatz von Basis und Spitze zeigen) Letztere gerade oder schraubig Sporenbildung nicht nachgewiesen Genera *Beggiatoa*, *Leptothrix*, *Crenothrix*, *Phragmidiothrix*

Later Baumgarten (1890, p 12) used the same family name as a "genus" with the following description

Leptoturicheen (Zopf) die vegetativen Zellen besitzen Stabchen- und Schraubige Form, durch Verbildung der Einzelzellen entstehen grade, wellige und schraubige Faden, die Faden zeigen bisweilen dadurch, dass das eine Ende sich an einer Stelle des Nährbodens festsetzt, einen Gegensatz von Basis und Spitze An den freien Enden der Faden gliedern sich kugelige Bildungen ab, welche theilweise wohl unzweifelhaft die Bedeutung von Arthrosporen haben, ob diese kugeligen Glieder, wie Zopf annimmt, unter Umständen auch direct theilungsfähig sind, d h also sich wie echte Kokken verhalten, erscheint noch sehr fraglich Bei einigen Arten dieser Gattung (*Crenothrix*arten) bilden sich um die Faden Scheiden, in denen sich Eisensalze allagern können

Untergattungen *Leptothrix*, *Beggiatoa*, *Crenothrix*, *Phragmidiothrix*

Hueppe (1891, p 30) also used this form of spelling

Leptotrichia A genus of bacteria created by Trevisan (1879, p 138) with the description "*Somatia cylindrica, plus minus distincte articulata, tenua elongata, filiformia, recta, laxe fasciculata*" The justification for this name for a group of organisms from the older genus *Leptothrix* is given as follows

Da denominazione generica *Leptothrix* fu tanto istrattata che sarebbe forse miglior partito inatterla fatto da parte, in ogni caso spetta alle Alghe, Percio, e per mutare il meno possibile, modificherei la desinenza

One species, *L buccalis* is given with *Leptothrix buccalis* Robin and Lebert as a synonym Trevisan (1889, p 10) included this as the second genus of his tribe *Leptotrichiee* Fourteen species are listed, the thirteenth in the list being *Leptotrichia buccalis* The synonymy and generic diagnosis are given by De Toni and Trevisan (1889, p 932) as follows.

Leptotrichia Trevisan (1879) Batter ital, p 14, Gen, p 10, *Leptothrix* Kuetz (1843) Phycologia generalis, p 198, ex parte, ein Winter Die Pilze, p 56, Schroet Pilz Schles, p 170 *Leptonema* Rabenh (1857) non Jussieu (1824) nec Hooker (1844), *Ophryothrix* Borzi (1878) N Giorn bat ital, x, p 274, *Thothrix* Winogradsky (1888) Beitr I, p 29 Filamenti cylindrica, vagina tenuissima obducta, articulata, simplica, basi ab apice superiori distincta, inaequilata, ut plurimum propter pulvinulum mucosum affixa, raro libera, immobilia vel rarissime lente oscillantia Multiplicatio filamentorum divisione ad unam directionem in articulos mobiles (baculogonidia) secedentium

Thirteen species of this genus are listed, included under four subgenera, *Leucothrix* Oeisted, *Ophryothrix* Borzi, *Thiothrix* Winogradsky and *Leptotrichiella* Trevisan

It is of interest to note that the original species named (*Leptotrichia buccalis*) is no longer included, but removed to the genus *Rasmussenia*

Winslow *et al.* (Committee Soc Am Bact, 1917, p. 555) accept this genus with the following description

Thick, long, straight or curved threads, frequently clubbed at one end and tapering to the other Gram-positive when young Threads fragment into short, thick rods Anaerobic or facultative Non-motile Filaments sometimes granular, non-branching No aerial hyphae or conidia Parasites or facultative parasites

The type species is *Leptotrichia buccalis* (Robin) Trevisan

Buchanan (1918, p. 405) states "Rod shaped or filamentous cells, non-motile, unbranched, without aerial hyphae or conidia parasites or facultative parasites"

Bergey *et al.* (1923, p. 338) follow the Committee, including this as the second genus of the family *Actinomycetaceae*

In spite of Trevisan's disavowal, *Leptotrichia buccalis* (Robin) Trevisan is the type of the genus

Leptotrichieae A tribal designation used by De Toni and Trevisan (1889, p. 932) with the following description "*Sporae nullae a it saltem huiusque numquam detectae Filamenta simplicia*"

The following genera are included, *Leptotrichia* Trevisan *Phragmidiothrix* Engler and *Beggiator* Trevisan

Leptotrichiee A name given by Trevisan (1889, p. 10) to the second tribe of his suborder *Tricogene* The description is "Semplice" Six genera were included, *Cladothrix*, *Leptotrichia*, *Phragmidiothrix*, *Beggiatoa*, *Kurthia* and *Billetia*

Leptotrichiella The fourth subgenus of the genus *Leptotrichia* according to De Toni and Trevisan (1889, p. 935) The sole description is "Zoobia" The only species is *Leptotrichia amphibola* Trevisan, from the stomach of a dog, the *Streptomesobacterio* of Maggi

Leucocystis. A generic name proposed by Schroeter (1883, p. 196) for the species *Leucocystis cellaris* The organism was found in slimy masses on the walls of wine cellars The original description follows

Die Hauptmasse des Schleimes wird gebildet durch zahllose Schizomyceten, die verschiedenen Gattungen angehoren Die haufigste und zwar uberall mit Regelmassigkeit vorkommende Form ist ein eigenthamlicher *Micrococcus*, den ich als *Leucocystis cellaris* beschreiben will Er bildet kugelige oder kurz ellip-

tische Körper von 1,5–2 μ Länge und etwa 1–1,5 μ Breite, die farblos, stark lichtbrechend sind. Sie werden von einer weiten Gallerthülle umgeben, die den Durchmesser des Coccus um das Mehrfache übertrifft, und dadurch eine Weite von 5 bis 8 μ erreichen kann. Die Coccen mit ihren Hüllen liegen dicht an einander, grosse Klumpen bildend, die Hüllen fließen aber nicht zusammen, sondern zeigen sich bei Zerdrücken der Ballen immer gesondert. Man findet in den verschiedenen Ballen Coccen in den verschiedenen Entwicklungszuständen, aus denen man erkennt, dass sich die Coccen durch fortgesetzte Zweitheilung in allen drei Richtungen des Raumes vermehren. Bis zu einer gewissen Grenze bleiben sie dabei in der Mutterhülle eingeschlossen, man findet daher Hüllen, welche einen, andere, welche zwei, vier und selbst acht Coccen enthalten. Die Hüllen, die mit zahlreichen Coccen gefüllt sind, sind etwas weiter als die, welche nur wenige enthalten, aber im Verhältniss nicht so dick. Durch Anilin-Farbstoffe wurden die Coccen stark tingirt, aber auch die Hüllen färben sich durch dieselben kräftig. Wendet man starke Farbstofflösungen an, so sieht man dann nur die stark gefärbten Hüllen, die selbst das Bild riesiger Coccen vortauschen. Bei vorsichtiger Färbung, indem man sich durch scharfes Aufpressen und Abziehen des Deckglases eine sehr dünne Schicht darstellt, diese über der Flamme massig erwärmt, dann mit nicht zu stark concentrirter Farbstofflösung allmählich färbt, erhält man sehr schöne Bilder, in welchen sich die Coccen lebhaft, die Hüllen schwach gefärbt zeigen. Bei Betrachtung unter Oelimmersion und mit Beleuchtungsapparat sieht man nun, dass häufig die einzelnen Coccen in der Mutterhülle besondere Hüllen umgeben, oft auch ist die Mutterhülle aus mehreren bis zu 5, verschiedenen Schichten gebildet, so dass das Ganze im optischen Durchschnitt fast das Aussehen eines Kartoffelstärkekornes erhält. Wurden die Coccen mit alkalischen Farbstofflösungen gefärbt, so wurden sie durch Zusatz von Säuren wieder vollständig entfärbt.

Diese Coccen haben in ihren einfachsten Entwicklungszuständen eine gewisse Ähnlichkeit mit denen, welche von C. Friedländer bei den an croupöser Pneumonie leidenden Kranken in den Lungen mit Constanz gefunden worden sind. Bekanntlich sind auch diese Coccen mit einer starken Schleimhülle umgeben, und auch bei ihnen sind die ersten Theilungszustände meist noch in dieselbe Hülle eingeschlossen. Weiter reicht indess die Ähnlichkeit nicht. Die vielkernigen Cysten von *Leucocystis* sind bei dem Pneumonie-Coccus (*Leucocystis Pneumoniae*) nicht gefunden, auch ist bei ihm eine Schichtung der Hülle nicht nachgewiesen.

Ich habe den Coccenschleim wochenlang im feuchten Raume unverändert gehalten, eine wirkliche Zucht der Coccen, eine Vermehrung derselben habe ich aber bis jetzt nicht erzielen können. In Alcohol lassen sie sich sehr gut conserviren, sie nehmen auf Zusatz von Ammoniak nach langer Conveirungen ihre ursprüngliche Gestalt wieder an, zeigen auch Hülle wieder unverändert.

Later (1884, p. 299) he concluded the organism to be the same as *Erebonema*. Later (1886, p. 152) he gave the following generic diagnosis: "Zellen kuglig oder elliptisch einzeln oder zu mehreren zusammenstehend, von weiten, mehrschichtigen fest abgegrenzten Gallerthüllen umgeben und in Schleimmassen zusammenfließend."

Hansgurg (1888, p 266) noted the presence of this organism in wine cellars, and gave *Erebonema hercynicum* Ktz as a synonym. The genus is also recognized by De Toni and Trevisan (1889, p. 1041) with the description

Cocci globosi v divisiones tempore globosoovoides, pauci in muco matricali nidulantes, segregati, in familias parvulas, singulas cystidibus universalibus, amplis, crassis, plurilamellosis, gelatinosas, involutas, inordinate consociati. Cystides in membranulam mucosam confluentes. Cystides speciales nullae. Coccorum divisio in tres directiones. Obs Generis *Thiothece* clade affines, a quo cystidibus e stratis pluribus latis efformatis in primis distinguitur.

The only species given is *Leucocystis cellaris* with the variety *cavernarum* Hansgurg.

Migula (1900 p 195) reduced this generic name to a synonym of *Micrococcus*, the species becoming *Micrococcus cellaris*. The genus is rejected by Erwin F Smith (1905). Apparently the name has not been recognized by other authors.

If the organisms described by Schroeter is worthy of generic rank, probably *Leucocystis* is valid as its name.

Leuconostoc A generic name proposed by Van Tieghem (1878, p 198) for the organism studied by Cienkowski and regarded as responsible for the formation of gum in sugar factories. Cienkowski (1878) named his organism *Ascococcus mesenteroides*. Van Tieghem showed that this organism was distinct from Cohn's *Ascococcus* (q v). The genus *Leuconostoc* was created with the following diagnosis:

Cellulae achromaticae minimae globosae in catenas laxas flexuoso-curvedas implicatae, vagina gelatinose cartilaginea lobata crassissima circumdatae consociatae. Vaginae in thallum gelatinose-cartilagineum, subglobosum vel crassissimo membranaceum, irregulariter expansum, extus cerebroideum intus pseudoparenchymaticum aggregatae. Sporae singulae globosae majores, terminales vel interstiales, pachydermaticae, intus homogeneae.

Van Tieghem places *Leuconostoc* among the *Nematogenae* of Cohn's classification, while *Ascococcus* was placed by Cohn in the *Glaeogenae*.

The genus with its species *Leuconostoc mesenteroides* (Cienkowski) Van Tieghem was recognized by Winter (1880) and by Grove (1884, p 16).

It was included by Van Tieghem (1884) in the tribe *Oscillariae* of the family *Nostocaceae*. De Bary (1884) describes the genus as consisting of rosaries of cocci embedded in firm sheaths of mucilage, and with arthrospores. Zopf (1885, p 61) states "Bildet Coccen

und Stabchen Sporenbildung in Coccen" Hueppe (1886, p 144) included *Leuconostoc* in his "Gattung Arthro-Kokkaceen" Schroeter (1886, p 154) likewise emphasizes the production of spores His diagnosis is as follows

Vegetative Zellen kugelig, zu Ketten verbunden Jede Kette in eine besondere Schleimkapsel eingeschlossen und in grosserer Menge zu zarten Schleimmassen verbunden Sporen aus einer Zelle (einem Gliede der Kette) gebildet, welche anschwellt und sich mit einer festen Membran umgiebt Aus jeder Spore geht bei geeigneter Ernährung eine neuer Kette hervor

Maggi (1886, p. 84) recognized the genus

Ludwig (1886, p XXV) named a *Leuconostoc Lagerheimii* from the oak slime

Hansgirk (1888, p 266) placed this genus in his subfamily *Cystococcaceae*, and described a variety, *subterraneum* of a *L Lagerheimii* first noted by Ludwig (1886, p XVII)

This genus is described by Trevisan (1889, p 28), De Toni and Trevisan (1889, p. 1051) and by Sternberg (1892) Ludwig (1892, p 29) noted that the genus contained organisms in which spherical cells occurred in chains and embedded in slime Lehmann and Neumann (1896, p 102) concluded that *Leuconostoc* is only a *Streptococcus* with enormously thickened capsules Macé (1897) used much the same description as Schroeter Migula (1900, p 25) followed Lehmann and Neumann in concluding that *Leuconostoc* is not distinct from *Streptococcus*, and combined the two genera under the latter name The interpretation of Migula has been accepted by most subsequent authors Barendrecht (1901, p 632) described a *Leuconostoc agglutinans*, a lactic acid organism from wort It was later named *Lactococcus agglutinans* by Beijerinck (1908, p 644) The generic name is rejected by Erwin F Smith (1905, p 174)

Winslow and Rogers (1905, p 669) have revived Cohn's *Ascococcus*, accrediting it to Cienkowski and emending it to replace *Leuconostoc* They state

Van Tieghem (1878) a little later worked on the same form (*Ascococcus nesen-teroides* Cienkowski) and substituted for *Ascococcus* the generic name *Leuconostoc* in order to emphasize the resemblance between zoogloea-forming Coccus and the blue green alga, *Nostoc*

It would seem, however, that Cohn's name *Ascococcus* was applied to an entirely distinct type of organism, and that the Winslows' substitution of *Ascococcus* is unfortunate Their characterization of the genus, however, gives the modern conception of this form

Saprophytic, growing vigorously in saccharine solutions. Cells in pairs or in true chains. In presence of certain carbohydrates large, lobed, gelatinous masses of zoogloae are formed. Fermentation powers high, acid being produced in dextrose, lactose and saccharose.

Vuillemin (1913, p. 520) concludes that *Leuconostoc* should be treated as a section of the genus *Streptococcus*.

Buchanan (1917, p. 611) states

Saprophytic, usually growing in cane sugar solutions. Cells in chains or pairs. Forming large zoogloal masses when grown in sugar solutions. Produces acid actively from carbohydrates. Some types at least are Gram-negative.

The type species is *Leuconostoc mesenteroides* (Cienkowski) Van Tieghem.

Winslow *et al.* (Committee American Soc. of Bact. 1920, p. 206) give the following diagnosis

Saprophytes, usually growing in cane sugar solutions. Cells in chains or pairs, united in large zoogloal masses. Some types at least Gram-negative.

Type species, *L. mesenteroides* (Cienkowski) Van Tieghem.

Bergey *et al.* (1923, p. 45) follows the Committee, including the genus as the second in the tribe *Streptococceae*.

Leucothrix. A generic name adopted from Oersted (1844, p. 44) by De Toni and Trevisan (1889, p. 932) to serve as a subgenus of *Leptotrichia* Trevisan. The type species is *Leptotrichia* (*Leucothrix*) *mucor* (Oersted) Trevisan. The subgeneric description given is simply "*Marinae, majores*". *Leucothrix* was previously used by Trevisan (1879, p. 138) as a synonym in part of *Beggiatoa*.

Lineola. A name used by von Baer (1827, p. 748) in the following sentence

Die Reihe für die Thiere des Typus mit vorherrschender Längendimension beginnt mit lebendigen Fäden, Lineola (so mögen die einfachsten Vibrationen heissen), repräsentirt sie unter den Protozoen.

Lipobacter. A generic name proposed by de Kruyff (1906, p. 1) for bacteria which hydrolyze and oxidize fat. These organisms are common in the soil, hydrolyzing fat to glycerine and fatty acids and later oxidizing each of these products.

If these organisms are deserving of generic recognition, this name possibly would be valid, although his species were numbered only and not named.

Liquidobacterium. A generic name proposed by Orla-Jensen (1909, p. 337) for organisms of the *Proteus* type. He says

Die Definition einer *Proteus* Bakterie wird daher Eine sporenfreie, peritriche Stäbchenbakterie, welche die Gelatine verflüssigen kann, und ihr Gattungsname muss in Analogie mit meiner übrigen Nomenklatur *Liquidobacterium* sein Laut dieser Definition müssen wir unter anderem die farbstoffbildenden Bakterien *B. prodigiosum* und *B. violaceum* als *Proteus*-Bakterien betrachten Dies stimmt auch gut denn beide erzeugen Indol und andere stinkende Produkte Ueberdies bildet *Liquidobacterium prodigiosum* ebenso wie *Liquidobacterium vulgare* aus Zucker vornehmlich Bersteinsäure und Ameisensäure Es erinnert durch sein hohes Fettspaltungsvermögen an *Liquidomonas fluorescens* und durch seine Vorliebe für starkehaltige Substanzen an die Kartoffelbacillen

The genus is assigned to the order *Peritrichinae* family *Alkalibacteriaceae*, and is defined as requiring as a rule organic N sources, and as being an aerobic decay bacterium which does not produce spores, and which liquefies gelatin

Since it is first mentioned, probably *Liquidobacterium prodigiosum* (Ehrenberg) Orla-Jensen is to be regarded as the type

Liquidococcus. A generic name proposed by Orla-Jensen (1909, p 332) The genus is included in the family *Luminobacteriaceae* of the order *Cephalotrichinae*

It is defined to contain those cocci which liquefy gelatin and which do not as a rule require organic nitrogen sources for development No species is named or designated

As a generic name *Liquidococcus* is probably best regarded as invalid or as a synonym of *Staphylococcus* or *Micrococcus*

Liquidomonas A genus of bacteria proposed by Orla-Jensen (1909, p 332) It is included in the family *Luminobacteriaceae* of the order *Cephalotrichinae* It includes straight rod-shaped organisms which are active liquifiers, which as a rule do not require organic nitrogen, aerobic and are as a rule fluorescent and sometimes denitrifying

The first mentioned organism (type) is *Liquidomonas fluorescens* Orla-Jensen

Later (1921, p 271) he states

As the property of liquefying gelatin also cannot be adopted as a generic character, I think it would be best to group together my earlier genera *Denitromonas* and *Liquidomonas* in a single genus, which can be conveniently termed *Fluormonas*, as the bacteria of this group are ordinarily fluorescent I cannot agree to call this genus *Pseudomonas* merely out of regard for so-called priority, as each and all of the bacteria which belong to the order under consideration are really *Pseudomonades* as well

And also

Perhaps the Committee is right in not regarding the *Proteus*-bacteria (my genus *Liquidomonas*) as a separate genus, as their whole metabolism indicates that they are to be looked upon as gelatin-liquefying *Coli*- and *Aerogenes*-bacteria.

Probably this genus is to be regarded as a synonym of *Pseudomonas*.

Liquidovibrio. A generic name proposed by Orla-Jensen (1909, p 334). It is included in his family *Luminibacteriaceae* of the order *Cephalotrichinae*.

The generic characters are Cells bent or spiral, usually not requiring an organic source of nitrogen, aerobic, often photogenic. The type species given is *Liquidovibrio comma*.

This name is probably to be regarded as invalid, and a synonym of *Vibrio* or of *Microspira*.

Lophotricha. A group name proposed by Messea (1891) to include bacteria with a cluster of flagella at one pole.

Luminibacteriaceae. A family name proposed by Orla-Jensen (1909, p 333) to include the photogenic and fluorescent bacteria which do not contain sulphur or bacterio-purpurin, cells never in thread-like filaments and with no tendency to branching. The genera included are *Denitromonas*, *Liquidomonas*, *Liquidococcus*, *Solidococcus*, and *Liquidovibrio*.

Macintoshillus. A name given by Heller (1922, p 13) to the third genus of her tribe *Clostridioideae* with the following description:

Clostridioideae that do not liquefy gelatin. They produce acid and gas and no putrefaction in meat media. They do not readily attack milk and they ferment few or no sugars. Gram-negative rods with parallel sides and with terminal spherical spores. Colonies in deep agar are small and irregular but not woolly. Frequently found in wounds. Apparently incapable of invading tissue.

Type species *tetanomorphus* (pseudotetanus bacillus McIntosh (p 32), *Bacillus tetanomorphus* Committee (p 41)), as described by the Committee. Glucose and maltose are fermented.

Macrococcus. A name occasionally used to contrast with micrococcus. It apparently has not been used as a generic name. It has the same meaning as megacoccus which is the preferable form. It has been used principally with the spelling "*Μακροκόκκος*" (*qv*). It is listed among the rejected names by Erwin F. Smith (1945).

Makrokokkus. A name used by Müller (1892 p 73) in his discussion of cocci. He states "Dieselben zeigen sehr verschiedene Dimensionen, von sehr kleinen runden Kokken bis zu auffallend grossen Kokken "*Makrokokken*." It is an aberrant spelling of *Macrococcus* (*qv*). A better form is *Megacoccus*. It is not a generic name.

Malleomyces. According to Trevisan (1879, p 144) this name was applied by Hallier to the organism causing glanders, with the name *Malleomyces equestris*. It is possible that this should displace the much less appropriate *Pfeifferella* as a generic designation for the glanders group.

Mantegazzaea. A genus created by Trevisan (1879, p 137). The description is as follows:

Somatia fusiformia vel cylindrica, distincte articulata, valida abbreviata, recta, segregata. Oss. VI appartengono due specie *Mantegazzaea Cienkowski* e *M. articulata* (*Bacterium articulatum* e *B. triloculare Ehrenb.*, (1831). La prima ricorderebbe alquanto per la forma *Bacterium roseum* e *Bacterium fusiforme* di Warming, la seconda *Bacterium Lineola*. Da *Bacterium* però differiscono perché evidentemente composte di più cellule, né ad alcun altro genere di Vibrionee potrebbero riferirsi. Il nome generico ricorda Paolo Mantegazza, che primo di tutti (1851) investigò e descrisse le spore perduranti delle Batteriacee in germinazione.

Later (1889, p 12) he included it as the fourth genus of the tribe *Bacillae*. Five species are listed. The generic description is "Baculi fusiformi con protoplasma uniformemente diffuso. Spore ignotae."

The synonymy given by De Toni and Trevisan (1889, p 942) is as follows:

Rhabdomonas Cohn (1875) Beitr 1, 3 p 167 (nomen ineptum)

Rhabdochromatium Winogr (1888) Beitr 1 p 100 et in Bot. Centralbl. 1889, N 6, 172

The diagnosis is as follows: "Baculi fusiformes vel subcylindracei, recti, plasmate uniformiter diffuse, divisione transversali sese multiplicantes. Sporae ignotae."

Two subgenera are recognized, *Rhabdochromatium* with three species and *Eumantegazzaea* with three.

This generic name has not been generally recognized by writers on bacterial nomenclature. Vuillemin (1913, p 521) has proposed that this name be retained as one of the "formogenera conservanda" with *Mantegazzaea (Bacillus) hastilis* (Seitz, 1889), the fusiform bacillus of the medical writers as the type species. This last suggestion would be invalid because of the earlier use of the name *Fusiformis*.

The type species (first species named and described) is *Mantegazzaea Cienkowski* Trevisan. It is an organism occurring upon the surface of putrefying meat. Its identification is probably quite uncertain. It is quite improbable that it is related to the so-called "fusiform" mouth bacteria, and its use should not be revived.

Martellillus. A name given by Heller (1922, p. 25) to one of the genera of her tribe *Putrificoideae*

Hardy *Putrificoideae* that in meat medium multiply rapidly at an early stage of incubation, producing a greyish coloration and later a blackish deposit on the meat particles, and after three days' incubation cease to multiply actively. Sporulate early in the development of a culture, later cease to do so but vegetate very slowly. Produce very little gas in meat medium. Digest milk. Attack a few sugars. Heavy deeply gram-positive rods, may vary greatly in size. Spores usually cocoon-shaped, usually median or sub-terminal, do not greatly bulge the sides of the bacillus. Colonies in deep agar lenticular, irregular, or stellate. Common putrefactive organisms that readily invade tissue in company with other organisms, producing a greenish proteolytic gangrene. I have found them in the heart's blood and organs of a woman dying of pernicious vomiting and uraemic poisoning and apparently the only invader in a mouse dying from an otherwise unknown cause. The Committee find *B. bifementans* in acute cases of gas gangrene. Weinberg and Séguin state that such an organism invades guinea-pigs in company with *B. perfringens*. It is possible that *Martellillus* bacilli produce metabolites poisonous to themselves and to animals and that cause early sporulation. They frequently cease active multiplication when the culture has a reaction near pH 7.0.

Type species *M. bifementans* (*Bacillus bifementans-sporogenes* Tisser and Martelly) as defined by Tisser and Martelly.

Mastichomonas. A subgenus of *Monas* used by Diesing (1850, p. 22). Enlows (1920, p. 55) states

Animalcula solitaria libera. Corpus ecaudatum elongatum, subglobosum, ovatum, turgidum, v. planum, haud v. mollitie sua solum mutabile, divisione spontanea simplici perfecta bipartitum v. indivisum. Os terminale. Flagellum simplex. Ocellus nullus. Est Parenema (Euparenema) corpore haud mutabile. Includes under this subgenus 22 species. *Monas Mastichenensis* termo Muller (Ehrenberg). *M. Mastichenensis punctata*. *M. Mastichenensis*. *M. Mastichenensis*, etc.

Megabacteria. Use ascribed to Billroth (1874, p. 16) for one of the growth forms of his *Coccobacteria septica*. See *Megabacterien*.

Megabacterien. A casual name used by Billroth (1874, p. 16) for one of the growth forms of his *Coccobacteria septica*.

Megabacterium. A name ascribed to Billroth (1874, p. 16) with the spelling *Megabacterien* to designate that growth form of his pleomorphic *Coccobacteria septica* in which the cells assumed the shape of large rods. The name has apparently never been used as a generic designation, although Smith (1905) has included this name in his list of invalid bacterial genera.

Megacoccus. A name used by Billroth (1874, p. 16) for a growth form of his pleomorphic *Coccobacteria septica* in which the cells assume

the form of large cocci. The name has apparently never been accepted as a generic designation, although it is included by Smith (1905) (with the spelling *Megacoccus*) in his list of invalid genera.

Megacoccus. Maggi (1886, p. 83) termed one of his bacteria (the fermentum butyricum of Fitz) *Micrococcus* (*Megacoccus*) *butyricus*. The name was listed as invalid by Erwin F. Smith (1905, p. 152).

The incidental use by Maggi is scarcely sufficient to validate the term.

Megalothrix. A generic name proposed by Schwerts (1912, p. 273) to designate a genus of iron bacteria, with the single species *Megalothrix discophora*. The original description follows:

Besonders ist mir seit Jahren eine Fadenbakterie aufgefallen, welche sich von den *Leptothrix ochracea*-Fäden stets leicht unterscheiden liess durch einen zarter abgegrenzten Kanal und durch eine sehr breite, homogene, oder sehr feinkörnige, graue, hellgelbe oder hellorange Scheide, deren Umfang nur langsam nach einem Ende des Fadens abnahm, seltener kam eine Dichotomie vor, selten auch war es mir möglich, durch die dicke Scheide eine deutliche Einteilung des Fadens in langliche Zellen zu sehen.

Diesen eigenartigen Faden beobachtete ich zuerst nur spärlich bei zahlreichen anderen Bakterien, bis ich einen eisenhaltigen Niederschlag auffand, der aus lauter solchen Fäden bestand. Auffallend war in diesem Präparat schon bei geringer Vergrösserung, die Gruppierung der Fäden, deren Büschel perückenförmig an einer Seite scharf abgegrenzt waren. Bei genauer Beobachtung konnte man erkennen, dass jeder Faden für sich an seinem dicken Ende mit einem dunklen Quertrich aufhorte, und dass gewisse dieser Büschel von diesem Strich ab auf einer dünnen Pflanzenwurzel hafteten. Ferner liess sich bei vereinzelter Fäden der Strich wiederfinden, bei anderen zahlreichen Fäden war aber der Strich durch eine runde Scheibe ersetzt, in deren Mitte der Kanal des Fadens deutlich endete. Dazu boten die isolierten Fäden nie ein solches Bild an beiden Enden. Bei Bruchstücken waren gewöhnlich die beiden Enden ohne Haftscheibe, nur selten war diese an einem der Enden vorhanden. Es handelte sich also hier um Fadenbakterien, die mit einer Haftscheibe enden, letztere entspricht dem breitesten Ende des Fadens und stimmt mit ihm in der Grösze überein. Je nach der Stellung des Fadens wird im Gelatineglyzerinpräparat die Scheibe seitlich schrag oder von vorne gesehen, und ist stabchenförmig, oval oder rund. In der Mitte dieser runden Scheibe ist ein runder, heller Hof, welcher dem Ende des Kanals entspricht. An dieser Stelle scheint die Scheibe bis auf das Ausserste verdünnt zu sein. Bei Fäden von einer Länge von durchschnittlich 300μ —wenn nicht gebrochen—einer Breite von $10\text{--}12\mu$ am dicken Ende und einer Kanalweite von $1\text{--}1,5\mu$ hatte die Haftscheibe entsprechend einen Durchmesser von $10\text{--}12\mu$ wovon $1\text{--}1,5\mu$ auf den Hof entfielen, und eine Dicke von $0,3$ bis $0,5\mu$.

Enlows (1920, p. 55) gives the following characterization:

Type species (monotypy) *M. discophora*. Threads 300μ long, and 8 to 12μ wide, which contain longish cells. Dist. no. shed from *Leptothrix* 1 = 41.

session always of a delicate sharply defined canal, and a very wide, homogeneous or very finely granular, gray, bright yellow or bright orange sheath, whose circumference decreases gradually toward one end of the thread, dichotomy rare, seldom possible, because of thickness of sheath, to observe the division of the threads into the long cells

Ellis (1913, p 450) after a study of this organism concluded it to be simply a growth form of *Crenothrix polyspora* Cohn

Melanella. A generic name created by Bory de St Vincent (1824, p 46) It is later described (1826, p 533) as the second genus of the family *Vibronides*. The organisms were described as linear or worm-like, opaque, not coiled in a discoid spiral

Bory described at least three species of organisms, *Melanella monadina*, *M flexuosa* and *M Spirillum* As a generic name this term seems to have been used by no other writers. Ehrenberg (1828, p 17) gave *Melanella atoma* as a synonym of *Vibrio lineola* and *M erythraea* as a synonym of *V rugula* Attempts have been made by various authors to identify his types, particularly Vuillemin (1913, p 521) This author concludes that good generic characters and recognition of type species are impossible, and that the name should be rejected

Trevisan (1879, p 148) regarded *Melanella* as a synonym of *Vibrio*, listing *Melanella flexuosa* Bory and *M erythraea* Hempr et Ehrenb (1828) as synonyms of *Vibrio Rugula* The name is rejected by Erwin F Smith (1905)

Melococcus A generic name used by Amradzibi (1907, p 309) for the species *Melococcus ostrjanini*, an organism isolated from the alimentary tract of the bee moth *Galleria melonella* The organism in many ways resembled a *Streptococcus* culturally Growth was always slow Gelatin was not liquefied The organism stains well with fuchsin and by Gram's method No gas was formed from dextrose

It would appear that the description of the organism is inadequate, and the nomenclatural status of the generic name uncertain

Meloseira (*Melosira*) A genus of diatoms created by Agardh (1824) The organisms usually known as *Gallionella ferruginea* was incorrectly placed in this genus by Ralis (1843, p 351) as *Melosira ochracea* This is one of the genera of diatoms commonly recognized today, and the name is invalid as a bacterial designation

Meningococcus. A casual name frequently applied to the causal organism of epidemic cerebro-spinal meningitis in man.

According to Enlows (1920, p 56) the designation was first used by Foà and Bordono-Uffreduzzi (1888, p 67) The first use in a generic

sense, that is, with a species name, apparently is that of Coats and Forbes (1911, p 242) who used the combination *Meningococcus intracellularis*. This would seem to validate the name, even though the authors were not aware that they were introducing a new generic name. However, it must be dated from its first use in binomial nomenclature, and must yield priority to *Neisseria*.

Meningokokkus A variant of *Meningococcus* used by German writers as a casual name.

In binomial nomenclature apparently it was first used by Huebner (1896, p 6) in the form *Meningokokkus intracellularis*. *Neisseria* has priority.

Merismopedia. A genus of blue green algae created by Meyen, (1839, p 67). Kützting (1849, p 471) described the genus as follows: "Phycema parvulum non afficum, quadratum, planum, gonidius quaternus, solidis (*Aquaticae*)". Rabenhorst (1865, p 56) gives the following description: "Cellulae globosae, tempore divisionis oblongae, tegumentis subcrassis confluentibus, 4, 8, 16, 32, 64, 128 in familias tabulatas unistratas consociatae, thallum quadratum planum libere natans formantes". One species included, *M. littoralis*, is a form that is probably to be reckoned among the bacteria.

Merismopedia is a valid algal genus, its use in bacterial nomenclature is therefore quite invalid. However, certain writers have placed in this genus bacteria which resemble *Merismopedia* in cell shape and grouping. The bacterial equivalent is *Merista* Van Tieghem.

Robin (1853, p 331) placed *Sarcina ventriculi* Goodsir in this genus.

Zopf (1885, p 51) recognized this as a bacterial genus with the diagnosis: "Tafelcoccen. Theilungen nach zwei Richtungen des Raumes, zur Bildung von Zellflächen in Tafelform führend, deren Glieder Später ausser Verband treten". The only species assigned to the genus was the *Merismopedia Gonorrhoeae*, the newly described gonococcus. As a bacterial genus, *Merismopedia* was also recognized by Billet (1890, p 24) with the description: "Merismopedia, zoogléé tabulaire, à éléments unis par 4". Sternberg (1892, p 17) likewise included this genus in his classification of the cocci. Lehmann and Neumann (1896, p 102) conclude that all forms included in *Merismopedia* should be placed in *Micrococcus*. The name is rejected by Erwin F. Smith (1905).

Caspari (1874, p 104) named an organism *Merismopedium Reitenbachii*, which was included among the bacteria as *Lampromedia Reitenbachii* by De Toni and Trevisan (1889, p 1048). He says "Es ist

daher gegen richtige Grammatik diesen Gattungsnamen zu schreiben; *Merismopedia* wie Meyen (Weigmanns Archi V 67) "

Merismopedium. A correction of the spelling of *Merismopedi* *q v* proposed by Caspary (1874, p 104) and used in the new species *Merismopedium reitenbachii*

Merista. Cunningham (1839, p 47) listed *Merista laevigata* Banks and Soland Mss 1769 as a synonym of *Myrsine Urviller* one of the spermatophytes It is regarded as a synonym of *Rapanea* Aubl in Engler and Prantl's Pflanzenfamilien As a bacterial generic name this was used by Van Tieghem (1884, p 1114) with the following description "Deux directions de cloisonnement Thalle membraneux, se dissociant en tétrades canéés à mesure qu'il s'accroît" Hueppe (1886, p 144) included this type in his "Gattung Arthro-Kokkaceen" as an "Untergattung"

Prazmowski (1888, p 235) renamed the *Micrococcus ureae* of Cohn, *Merista ureae*. Apparently this is the first use of the term in a strict generic sense, that is, as a binomial for bacteria

Billet (1890, p 24) includes the genus in his "Formes zoogoniques" with the descriptions, "zoogone tabulaire, à éléments unis 4 par 4" Lehmann and Neumann (1896, p 102) concluded that the genus should be merged in *Micrococcus* Vuillemin (1913, p 525) concluded that this genus has the same reason for existence as has *Sarcina* or *Streptococcus*, and that the characteristic tetrad grouping is worthy of generic recognition He had proposed therefore that this be constituted a "*Genus conservandum*" with *Merista tetragena* (*Micrococcus tetragenus* Gaffky, *Gaffkya tetragena* Trevisan) as the type If *Mic tetragenus* is to constitute the type of a genus, it would seem that *Gaffkya* Trevisan has priority

The type species evidently is *Merista ureae* (Cohn) Prazmowski

Whether the previous use of *Merista* for a genus of phanerogams invalidates its use for bacteria is somewhat problematical Apparently Banks and Soland never published their name, it was never proposed Quotation in synonymy does not validate a name It would seem therefore that *Merista* Prazmowski is valid if its type requires separate generic recognition

Merizomyria. A genus of algae ascribed to Kützinger (1843) Rabenhorst (1865, p 224) lists six species, giving the following diagnosis

Trichomata flagelliformia simplicia rarius pseudoramosa, arcuissime vaginata, basi cellula perdurante instructa, sursum moniliformia, in apicem capillarem cuspidata, in mucosae matricis nidulantea plerumque in stratum difforme coacervata

This name is of significance in bacteriology because according to De Toni and Trevisan (1889, p 926) *Sphaerotilus thermalis* Kuetz is a synonym of *Merizomyria aponina* var *fasciculata* Kuetz

Mesobacteria. According to Enlows (1920, p 57) used by Billroth (1874, p 6) See *Mesobacterien*

Mesobacterien A name used by Billroth (1874, p 16) to designate a growth form of *Coccobacteria septica* in which the cells are rod shaped and of moderate size The name has apparently not been used as a generic name, although Smith (1905) has included this in his list of invalid genera, with the spelling *Mesobacterium*

Mesobacterium. Listed by Erwin F Smith (1905) as invalid. Probably refers to *Mesobacterien q v*

Mesococcus A name used by Billroth (1874, p 6) to designate a growth form of his *Coccobacteria septica* in which the cells were spherical and of moderate size The name has been spelled *Mesococcus* by later writers This term has apparently never been used as a generic designation

Mesococcus A variant spelling of *Mesococcus (q v)* It is listed by Smith (1905) as an invalid genus

Metabacterium. A genus of bacteria proposed by Chatton and Perard (1913, p 1232) for the species *Metabacterium polyspora* from the coecum of the cavy The authors give the following description

Elle se rattache aux bactéries sporées du type *B Butschli* Mais elle en diffère nettement par ce caractère jusqu'ici unique, croyons-nous, chez les Bactéries, de pouvoir former un nombre de spores supérieur à deux, et variable de une à huit, par élément Ces éléments sont toujours isolés Ils sont en forme de fuseau tronqué aux deux extrémités et mesurent de 10 à 25 μ de long sur 5 μ de large en moyenne Nous ne les avons jamais vus mobiles Ils présentent une mince membrane d'enveloppe à l'intérieur de laquelle on distingue une masse cytoplasmique peu dense, le plus souvent réduite à des trabécules La majeure partie du contenu est condensé en masses chromatiques, qui, lorsqu'elles sont au nombre de deux, sont situées chacune à l'un des pôles de l'élément Il est très rare de n'en observer qu'une, qui d'ailleurs est également polaire Il y a par contre très souvent plus de deux masses chromatiques, trois, quatre, cinq, jusqu'à huit, ces nombres étant d'autant plus fréquents qu'ils sont moins élevés Dans ces cas, deux des masses sont toujours polaires, les autres étant situées en des points quelconques de l'élément Les unes et les autres sont toujours au contact de la pellicule d'enveloppe Chez quelques bacilles à deux masses polaires, nous avons observé ces deux masses, ce gros noyau chromatique circonvoilé, que Schaudinn, Swellengrebel, Dobell ont décrit chez diverses bactéries sporées *B Butschli*, *B maximus-buccalis*, *B flexilis* et *B spirogyra* Nous ne pouvons nous prononcer sur la signification précise de cette structure que nous n'avons pu retrouver chez des éléments ayant plus de deux

masses chromatiques Quant à ces masses memes, elles ne nous paraissent pas avoir la valeur de vrais noyaux, bien que, chez certains individus, leur structure habituellement dense fasse place à une structure vésiculaire Nous n'avons d'ailleurs jamais vu ces masses se diviser (à moins que l'on doive interpréter comme une figure de division le noyau chromatique de la figure 3), et il semble qu'elles procèdent toutes de condensations isolées de substance chromatique Chacune d'eiles est l'ébauche d'une future spore Pour former les spores, ces masses s'accroissent et régularisent leur forme en celle d'un cylindre arrondi aux deux bouts, cylindre d'abord court, qui s'al longe progressivement sans s'épaissir notablement, jusqu'à mesurer de 10 à 15 μ de long sur 2 μ de large environ Il se forme autant de spores qu'il y avait de masses chromatiques, mais on voit souvent une ou plusieurs spores avorter en cours de développement La spore mure ne se colore plus en totalité, sa paroi devenue impénétrable ne prend plus au fer qu'une teinte gris bleu Au Ziehl, qui colore intensément en rouge les spores d'*Oscillospira*, celles de notre bacille ne prennent qu'une teinte rose pale Il persiste toujours au pole libre de la spore une calotte chromatique Le nombre de ces spores varie de une à huit par élément Elles sont d'autant plus longues et plus larges qu'elles sont moins nombreuses, sans que les écarts dépassent 5 μ en longueur et 1 μ en largeur Les spores étant longues chevauchent les unes sur les autres, lorsqu'elles sont plusieurs, comme chez les formes dispersées du B Dubosea de Joyeux La sporulation absorbe la plus grande partie du contenu de l'élément ne laissant qu'un insignifiant reliquat dans l'enveloppe qui se moule sur les spores Nous n'avons jamais vu les formes végétatives en division Nous n'avons pas constaté en particulier cette scissiparité complète ou abortive que Schaudinn et Dobell décrivent chez leurs bacteries dispersées avant la sporulation et que Schaudinn avait interprétée comme l'ébauche d'une zygose Nous n'oserions pas affirmer cependant que notre espèce ne se divise jamais Mais il est certain qu'elle presente par rapport aux autres bacteries une reproduction scissipare réduite compensée par une reproduction sporogénétique qui atteint chez elle un degré de perfection beaucoup plus élevé que chez les autres représentants du groupe L'absence ou la rareté de la reproduction scissipare n'en a pas moins cette conséquence que jamais notre bacille ne se présente en culture dense dans le caecum des cobayes Il parait d'ailleurs dépourvu de tout pouvoir pathogène

According to Vuillemin (1913, p 520) this should be regarded as a *genus conservandum*

Buchanan (1918, p 39) give the following generic description

Rod-shaped, known only in the sporulating state, from the caecum of a guinea-pig Sporogenous cell becomes ellipsoidal with one to eight endospores within a single cell

The type species is *Metabacterium polyspora* Chatton and Perard

He includes it as the fourth genus of his family *Bacillaceae*. Enlows (1920, p. 57) gives the following description

Type species (monotypy) *M polyspora* n sp Found in the caecum of the guinea-pig Characterized by its ability to form from 1 to 8 spores in a single

cell A truncated spindle-shaped organism, always single, 10 to 25μ long by 5μ wide Nonmotile Cytoplasm condensed into chromatic masses, from which the spores develop, which (when limited to 2) are usually bipolar The vegetative forms have never been observed to divide Nonpathogenic

Metacoccaceae. A subfamily of the family *Coccaceae* proposed by Winslow and Rogers (1905, p 669) to include the genera *Micrococcus*, *Sarcina* and *Rhodococcus* The diagnosis given is

Facultative parasites or saprophytes Thrive best under aerobic conditions Grow well on artificial media, producing abundant surface growths Planes of fission often at right angles, cells aggregated in groups, packets or zoogloea masses

In later descriptions (1906, p 485, 1908, p 258) the Winslows have added "Generally decolorize by Gram Pigment yellow or red "

It was pointed out by Buchanan (1915, p 9) that the name *Metacoccaceae* fails to conform to the rules of botanical nomenclature in that it does not carry the right ending and is not formed from the name of one of the constituent genera Possibly this name should be replaced by the tribal designation *Micrococceae* used by Trevisan

Metacoccus. A name stated by Conn (1909, p 12) to have been recognized in "recent classification of the spherical forms" The writer has been unable to find the classification to which Conn refers It may be regarded as a *nomen nudum*

Metallacter. A generic name proposed by Perty (1852, p 180) for a rod shaped bacterium occurring in filaments or chains The type was *M Bacillus*, the *Vibrio Bacillus* of Muller and of Ehrenberg The name was used as a synonym of *V Bacillus* Muller by Rabenhorst (1865, p 72)

Concerning Perty's organism, Cohn (1875, p 188) has the following statement:

Perty hatte, wie ich erst nachtraglich erfahren, unsere Bacillusarten schon im Jahre 1852 als eine selbständige Gattung von den eigentlichen Bac'erien unter dem Namen *Metallacter* abgetrennt (von einem griechischen Worte, welches sich "verändernd" bedeutet), weil dieselben in steife oder wenig biegsame, unbewegliche, ungemein verlängerte *Hygrocrocus* (oder *Leptothrix*)—fäden unter gewissen Umständen sich umwandeln, und ebenso hatte er äusserst kleine, bewegliche cylindrische Fäden beobachtet, welche an einem, oder seltener an beiden Enden eine, manchmal auch zwei elliptische Körperchen (wohl Sporen) einschliessen, obwohl er dieselben öfters mit *Bacillus* (*Metallacter Bacillus* Perty) zusammen fand, denen sie sehr gleichen, hielt er sie doch für eine selbstständige Gattung und Art (*Sporonema gracile* Perty)

Cohn created the genus *Bacillus* which replaced *Metallacter* Trevisan (1879, p. 137) recognized the genus *Metallacter* in the following description and statement

Metallacter Perty, 1852, *Bacteridium* Davaine (Compt rend de l' Acad des sc de Paris, Tom LVII) 1863, *Bacillus* Cohn (Beitr I, Heft 2, p 174) 1872, *Streptobacteria* Billroth, 1874 *Somatia cylindrica*, plus minus distincte

Oss Perty fondo il suo genere *Metallacter* sulla stessa specie (*Vibrio Bacillus* O F Muller) pella quale, vent' anni piu tardi, Cohn propose il suo *Bacillus* Il nome generico *Metallacter*, non contrario ai caratteri, ne ad alcuna legge di nomenclatura, anzi accennate la proprieta di mutarsi, in date condizioni, in somazj filiformi, straordinariamente allungati, leptotricheiformi, rigidi, immobili (1), quindi nome sotto ogni aspetto ineccezionabile, deve avere la preferenza. Resterebbe a vedere, non avendo il libro alla mano, se *Bactrella* di Morien (in Bullet des scienc natur, n XXVII, 1830) sia identico con *Metallacter*, nel qual caso la denominazione proposta da Morren dovrebbe essere conservata

As species he lists *Metallacter ulna*, *M bacillus* (*B subtilis*), *M. amylobacter*, *M tremulus*, and *M anthracis*

Later Trevisan (1882, p 151) again emphasized the genus *Metallacter* as follows

Per essa Klebs mise innanzi l'appellativo *Bacillus typhosus*, pero, sia pel diritto di priorita che il nome generico *Metallacter* (Perty 1852) ha sull'altro *Bacillus* (Cohn 1872), sia pel diritto di priorita che la piu appropriata denominazione specifica *Ileotyphi* ha sull' altra *typhosus* avrebbersi piu correttamente a chiamare *Metallacter Ileotyphi*

The next year he again used this name (1883, p 149) Two years later, however, he abandoned *Metallacter* for *Bacillus* (1885, p 94)

De Toni and Trevisan (1889, p 945) give *Metallacter* as a synonym of *Bacillus*

Migula (1879, p 7) has the following to say of Perty's genus

Sein *Metallacter* ist ein *Bacterium*, welches sich durch die Eigenschaft lange geteilte Faden bilden zu konnen, zeichnet "Bakterienahnliche Einzelwesen verlangern sich durch fortgesetzte Teilung zu steifen oder wenig biegsamen Faden, welche unter gewissen Umstanden nach einiger Zeit die Bewegung verlieren, ungemein wachsen und einer *Hygrocroctis* ahnlich werden, indem sie aus langeren verfilzten, flockige farblose oder grauliche Massen darstellenden Faden bestehen " Als Art rechnet er hierzu *M Bacillus* (= *Vibrio Bacillus*), ferner als Formen dieser Art *V subtilis* und *Bacterium Catenula* Du]

The name is rejected by Erwin F Smith (1905)

Vuillemin (1913, p 519) states that *Metallacter* is a synonym of *Serratia* and so disappears

With the exception of Trevisan, it does not appear that as a generic name *Metallacter* has been used since Perty's description. It is generally regarded as invalid. If it were possible to fix exactly the species *Metallacter bacillus* it probably would be necessary to recognize the genus. Unless it can be shown that it is the same as Cohn's *Bacillus subtilis* it will scarcely replace *Bacillus*. Trevisan's use of *Bacillus subtilis* as a synonym is scarcely adequate.

Metchnikovillus. A name proposed by Heller (1922, p. 29) for a genus of the tribe *Putrificoideae*. The description is

Highly proteolytic *Putrificoideae* that readily blacken meat. They do not produce in it abundant amino-acid crystals, but digest meat, serum, egg and casein rapidly, forming more alkali or less fatty acid than do the organisms of the succeeding group. They split few sugars. Gram-positive or weakly Gram-positive rods, vegetative forms uniform and considerably smaller than sporangia. Sporulate readily in ordinary media, forming oval spores which are usually subterminal, though in some strains median spores predominate. Multiplication is exceedingly active, forty-eight hour colonies in deep agar are large and woolly. Frequently a few colonies are larger than the others but they do not give rise in a following generation to a preponderating number of large colonies. Ubiquitous. Common intestinal organisms, abundant in soil, very common in infected wounds. Not capable of invading in pure culture in moderate doses, may invade in company with other organisms or alone when given in large doses.

Type species *M. sporogenes* (*Bacillus sporogenes* type A of Metchnikoff) as described by the Committee (p. 36). Klein described as *B. enteritidis-sporogenes* a mixed culture which contained a non-proteolytic organism, a pure culture of which was described by von Hibler as *B. enteritidis-sporogenes* Klein (von Hibler IV). This tissue invading pathogen, thought by some to have been *B. welchii*, is most referable to the genus *Arloingillus*. The strain was derived from patients with enteritis and it was apparently contaminated with a proteolytic organism of the genus *Metchnikovillus*. Metchnikoff described two types, A and B, of intestinal anaerobes which he thought were similar to the organism of Klein, and which he termed *B. sporogenes*. His descriptions permit of no identifications. Choukevitch, working in Metchnikoff's laboratory described *B. sporogenes* A and B more carefully. Type B should be referred to genus *Martellillus*. Weinberg made the identification of type A more exact, and the Committee discuss it at some length. They allow two species, *B. sporogenes* type A of Metchnikoff, and *B. parasporogenes* McIntosh, which are different in colony formation and serologically. Donaldson describes the "reading" bacillus, to be assigned to this genus. Superficial acquaintance with many strains of proteolytic anaerobes leads me to suggest that *Metchnikovillus* may be defined as a genus of many species, Dr. K. F. Meyer is also of this opinion. Henry (p. 361) believes it likely that the conception "sporogenes" refers to a group of organisms.

Methanomonas. A generic name proposed by Orla-Jensen (1909, p. 311). This is the first genus of the family *Oxydobacteriaceae* of the

order *Cephalotrichinae* The organisms of this genus are described as autotrophic, monotrichic, aerobic short rods capable of oxidizing methane

Winslow *et al.* (Committee Soc Am Bact 1917, p 551, 1920, p. 201) and Bergey *et al* (1923, p 32) describe the genus as follows

Monotrichic short rods capable of growing in the absence of organic matter and securing growth energy by the oxidation of methane (forming carbon dioxide and hydrogen) The type species is *Methanomonas methanica* (Sohngen) Jensen

Meyerillus. A generic name proposed by Heller (1922, p 22) for the fourteenth genus of her tribe *Clostridiodeae* Her diagnosis is

Clostridiodeae that produce H_2S on blood broth and liquefy gelatin In meat medium they produce gas, but little or no pink coloration, they show no marked proteolytic action Do not readily attack milk Large Gram-positive rods with little or no inclination to form spores Attack a few sugars Colonies in deep agar large, opaque and lenticular Typically attack connective tissue more than the muscle

Type species *M sadowa* nov sp To be described in a future paper This organism was at first taken for *B Welchii* It does not sporulate It is one of the four guinea-pig invaders isolated from a case of human gas gangrene I am inclined to place the sporulating bacillus L of Adamson (1919) in this genus but have not handled that organism Perhaps it is premature to decide whether sporulation may be used as a generic character Adamson finds bacillus L very slightly proteolytic on milk and not so on other media *M sadowa* does not grow on milk

Microbacillus According to Schamberg (1902, p 293), Sabouraud named an organism *Microbacillus seborrhoeae* The original references have not been available

Microbacteria. According to Enlows (1902, p 58) used by Billroth See *Microbacterien*

It was used by Cohn (1872, p 167) to designate his second tribe of bacteria

Die zweite Tribus der Bakterien bezeichne ich als *Microbacteria* oder *Stäbchenbakterien*, sie stimmen mit den Kugelbakterien in der Kleinheit ihrer Zellen und deren zeitweiser Vereinigung zu Gallert- oder Schleimmassen überein, unterscheiden sich jedoch, abgesehen von ihrer physiologischen Thätigkeit, durch die kurz-cylindrische Gestalt und die spontane Bewegung der Zellen

This name was also used by Hansgirk (1888, p 264) to designate the second family of his order *Eubacteria* Two genera were recognized, *Bacillus* and *Chromobacillus*

Microbacterien. A name used by Billroth (1874, p 16) to designate a growth form of his *Coccobacteria septica* in which the cells are rod-shaped and small

Microbacteries. Maggi (1886, p 84) used this designation for his third order of bacteria, with the single genus *Bacterium*. No description was given

Microbacterium. A form of the name *Microbacterien* (*q v*) It is listed by Smith (1905) with the invalid bacterial genera, although the writer has not been able to find an instance of its use as a generic name previous to this time Klein (1885, p 37) uses "*Bacterium* (*Microbacterium* Cohn)" as a heading Apparently Klein has confused this generic name with Cohn's *Microbacteria q v*

In a strict generic sense the name was first used by Orla-Jensen (1919, p. 179) in the following statement

Of the other Gram-positive, rod-shaped lactic acid bacteria (Table XXXII) which we have encountered in our investigations, the majority (Nos 3 to 10) are considerable smaller than the rod forms hitherto described, and have so many peculiar qualities in common that it will be only natural to collect them in a genus, which may suitably be named *Microbacterium*

The microbacteria do not, for the most part, curdle milk, and are on the whole weak acid formers and produce dextro-lactic acid, with the single exception of No 10, which forms inactive lactic acid When sown out in high agar tubes, they grow only in the upper part, and in stab cultures, they exhibit more or less pronounced surface growth No 7 even forms a highly curled surface layer Nos 3-6 on the other hand, give only surface growth with favourable sources of nitrogen, and even then not always to any perceptible degree The best nitrogen source for these bacteria is casein peptone, yeast extract, on the other hand, is as a rule very unfavorable With the exception of No 1, they split up hydrogen peroxide and reduce nitrate to nitrite In biological respects, the micro-bacteria thus greatly resemble the tetracocci, and there is also a gradual transition to forms which are no longer acid formers, but which liquefy gelatin to a slight degree, and can break down amino-acids The true microbacteria never ferment pentoses, and of alcohols, at the outside a little mannite

The microbacteria fall again into several well distinguished species Nos 3, 4, 5, and 6, for instance, are closely allied forms, and as they very often occur in milk, we will call them *Microbacterium lacticum* No 7 we will call *Microbacterium mesentericum*, from its very characteristic surface growth, and Nos 8 and 9, which have constantly exhibited a powerful yellow surface growth, *Microbacterium flavum* No 10 probably also constitutes a distinct species

Later (1921, p 272) he states

Microbacterium is to be understood as merely a provisional collective name for Gram-positive rods of size a little smaller than the ordinary bacteria. In biologi-

cal respects some of these rods (*Bacillus acidophilus*) are closely related to true lactic acid bacteria, whereas others approach the Tetracocci or the bacilli

The first mentioned species *Microbacterium lacticum* Orla-Jensen is apparently to be regarded as the type.

Micrococceae. A tribal name used by De Toni and Trevisan (1889, p 1067) with the following diagnosis

Cocci nudi, nec cystidibus, nec capsulis, nec vaginis obducti, liberi, nunquam moniliformiter concatenati, raro gemini vel in turmas aut zoogloas consociati Endosporeae microsomae in coccis normalibus obvenientes

It is included in the family *Coccigenae* The following genera are included *Neisseria*, *Staphylococcus*, *Micrococcus*

Buchanan (1917, p 613) recognized the tribe with the following description

Saprophytes or facultative parasites Usually grow best under aerobic conditions Grow well on artificial media, producing abundant surface growths Planes of fission often at right angles, cells aggregated in groups, packets or zoogloea masses Generally Gram-negative Usually pigmented yellow, orange or red

He included the genera *Micrococcus*, *Rhodococcus* and *Sarcina*

Winslow *et al* (Committee Soc Am Bact, 1917, p 559, 1920, p 207) have used practically the same description

It is also accepted by Castellani and Chalmers (1919, p 925) with the addition of the genus *Nigracoccus*, and by Bergey *et al* (1923, p 57) as the third tribe of the *Coccaceae*

Micrococcoglia A name used by Billroth (1874, p 14) to designate a growth form of his *Coccobacteria septica* in which the spherical cells were arranged in small zoogloea masses. The name has apparently not been used to designate a genus

Micrococcus An aberrant form of spelling of *Micrococcus* used by Billroth (1874, p 6) to designate that growth form or stage in the development of his *Coccobacteria septica* in which the organism consisted of very small spherical cells

Micrococcus. This name was first used by Haller (1866) to indicate a growth form of a mold in accordance with his conception of pleomorphism He did not use the name in a generic sense, and the genus should therefore not be ascribed to him, as has frequently been done by various authors Cohn (1872, p 151) adopted the name and defined it as a genus containing very small spherical or oval organisms,

with colorless or light colored cells, without motility, remaining after cell division united into rosary-like threads, many celled families or slimy masses. The first organism to which the name was given was *Micrococcus prodigiosus* (Ehr.) Cohn. This organism is now generally removed from this genus because of its shape. The next organism named was *M. luteus* (Schroet) Cohn. This form may well be considered the type of the genus as it has been adequately described.

The name *Micrococcus* was used by Billroth (1874, p. 6) to designate a growth form of his pleomorphic *Coccobacteria septica*, in which the cells consisted of very minute spheres. He did not use the name in a generic sense. In 1875 Cohn split his former genus into two, creating the new genus *Streptococcus*. His revised definition of *Micrococcus* included those organisms with spherical cells, united into glairy families, amorphous, cells without phycochrome, very small. Trevisan (1879 p. 126) gives the following generic description:

Micrococcus Halberl. em. Cohn (*Zoogalactina* Sette 1824, *Serratia* Bizio 1827, *Microzyma* Bechamp 1868, *Microsporon* Klebs 1872, *Microsphaera* Cohn 1872) Cellulae globosae vel globoso-ovoideae. Multiplicatio divisione cellularum vegetativa in omnes directiones.

It was included under *Eubacteriae*. In all, sixteen species were listed, many of them forms now included among the rod-shaped organisms. Among forms listed which may be included among the true cocci are *M. luteus*, *M. lactis*, *M. septicus*. Luerksen (1879, p. 171) followed the original description of Cohn, as did also Magnin, (1880, p. 72). The latter divided the genus into three groups of species: *Micrococcus chromogenes*, *M. zymogenes* and *M. pathogenes*. Winter (1880, p. 42) gave the following generic diagnosis:

Zellen farblos oder blass gefärbt, kugelig oder eiförmig-elliptisch, unbeweglich, nur in einer Richtung sich theilend. Die Tochterzellen trennen sich entweder bald von einander, oder bleiben zu zwei oder mehr kettenartig verbunden, oder bilden Zoogloea. Sporenbildung nicht sicher bekannt.

Was ich oben über die Unterscheidung der Arten gesagt habe, gilt ganz besonders von *Micrococcus*. Die angenommenen Arten zeigen in Gestalt und Grösse keine oder sehr geringe Unterschiede und es bleibt uns nur die chemische Thätigkeit als Mittel zur Auseinanderhaltung der Arten, die daher etwas ausführlicher behandelt worden ist.

Ogston (1883, p. 27) used the name *Micrococcus*, but without species designation. He included the chain forms (streptococci) as well as a new type which he termed staphylococci. Van Tieghem (1884), and Grove (1884, p. 6) all followed Cohn more or less closely, though

increasing the number of genera of spherical bacteria Zopf (1885, p 51) gives five genera of cocci, *Streptococcus*, *Merismopedia*, *Micrococcus*, *Ascococcus* and *Sarcina* *Micrococcus* is defined as made up of cocci, not in packets, not producing large quantities of capsular material, and with the cells irregularly arranged in grape like clusters He concluded that the generic name *Staphylococcus* of Rosenbach is a synonym Cornil and Babes (1886) use *Micrococcus* as a group name and include under it as subdivisions, *Streptococcus*, *Staphylococcus*, *Ascococcus* and *Sarcina* Flugge (1886) defined the genus as having cells oval or spherical, non-motile and with the cells separating as soon as division is complete Schroeter (1886, p 143) recognizes nine genera of cocci *Micrococcus* includes those whose cells are spherical single or arranged irregularly, and in which the cells or colonies are free or embedded in slime without definite gelatinous membranes

Maggi (1886, p 81) included this as the first genus of his order *Sphaerobactéries*

Hansgig (1888, p 267) included this as the only genus of the subfamily *Eucoccaceae*

De Toni and Trevisan (1889, p 1076) include some twenty-two genera among the cocci, one of them *Micrococcus* with a large number of species

Various authors before 1895 defined *Micrococcus* as containing those organisms with irregularly grouped spherical cells Among them were Billet (1890, p 24), Hueppe (1892, p 30), Sternberg (1892, p 17), Ludwig (1892) and Clements (1894, p 26) Migula 1892, p 236 says "*Micrococcus* Cohn Die zellen teilen sich nach 2 Richtungen des Raumes Bewegungsorgane fehlen (Z B *Micrococcus candidans* Flugge)"

Migula (1894) emended the description of *Micrococcus*, creating new genera for the motile forms The genus is given the following diagnosis (1895, p 16)

Single cells spherical, often united in twos and fours and then sometimes flattened on surfaces in contact Division in two directions of space Non-motile Endospore formation not certainly recognized, probably lacking

He notes that some four hundred species had been described

Fischer (1897, p 32) on the other hand, emphasized that *Micrococcus* consists of cells which divide irregularly, and not in definite sequence and which are non-motile Lehmann and Neumann (1896, p 148) used the following description

Die Zellen teilen sich unregelmässig nach verschiedenen Richtungen und liegen heirauf bald einzeln, bald zu 2 oder 4, endlich und zwar vorherrschend in regellosen klumpigen Haufen Hierher rechnen wir alle Kokken, die nicht unzweifelhatte Streptokokken oder Sarcinen sind

Many authors have followed the diagnoses of Migula and of Fischer Among these are Migula (1897, p 27, 1900, p 46 and 1904, p 144), Macé (1897, p 335), Hewlett (1898), Chester (1897, p 62), Baumgarten (1900, p 127), Chester (1901, p 71), Matzuschita (1902), A J Smith (1902, p 268), Kendall (1902), Klocker (1903, p 331), Fischer (1903, p 59), E F Smith (1905, p 160), Jensen (1909, p 340), Ellis (1909, p 4), Frost (1911, p 57), Meyer (1912, p 5), Schneider (1912, p 23) and Lohnis (1913) Migula (1900, p 46) gives the following list of synonyms *Lan. propedia* Schroter = *Microhaloa* Kutz exp., *Cohnia* Winter, *Hyalococcus* Schroter, *Leucocystis* Schroter, *Ascococcus* (Billroth) Cohn, *Monas* Ehrenberg exp, *Bacteridium* Schroter, *Diplococcus* Aut exp, *Staphylococcus* Aut, *Merismopedia*, exp, *Merista* Hueppe exp., *Gonococcus* Neisser

The Winslows (1905, p 669) have again emended the diagnosis of the genus In its most recent form it is (1908, p 258),

Facultative parasites or saprophytes Cells in plates or irregular masses (never in long chains or packets) Generally decolorize by Gram Growth in agar abundant, with formation of yellow pigment Dextrose slightly acid, lactose broth generally neutral Gelatin frequently liquefied Nitrates may or may not be reduced

They include as valid species, *Micrococcus flavus*, *M citreus*, *M luteus* and *M candicans* It will be noted that the species *M luteus* previously noted as a suitable type species is included

Vuillemin (1913, p 520) contends that *Micrococcus* should be termed a form genus only, and relegated to the bacteriologists who have vulgarized the term

Buchanan (1917, p 613) characterized the genus as follows

Saprophytes or facultative parasites Cells in plates or in irregular masses (never in long chains or packets) Generally Gram-negative Growth on agar abundant, with formation usually of yellow pigment Dextrose broth as a rule slightly acid and lactose broth neutral Gelatin frequently liquefied

The type species is *Micrococcus luteus* (Schroeter) Cohn

Winslow *et al* (Committee Soc Am Bact, 1917, p 561, 1920, p. 208) have used practically the same diagnosis and the same type designation

It is probable that *Micrococcus* Cohn should be recognized as a genus with the type *Micrococcus luteus* (Schroter) Cohn. This type designation is menaced by the fact that Cohn's first species described was *Micrococcus prodigiosus*. This form, however, is not a typical *Micrococcus* in that the cells are not spherical. Under the circumstances, the designation of the second species described, *M. luteus* is probably justified and valid.

Bergey *et al.* (1923, p. 58) has followed the Committee, including *Micrococcus* as the first genus of the tribe *Micrococceae*.

Microhaloa. A genus of blue green algae created by Kutzing (1843, p. 169) with the following description "Phycoma globosum, grumosum, ex cellulis in glomerulos coadunatis compositum." Two species are included, *M. botryoides* and *M. rupestris*. Its only relationship to bacterial nomenclature is the reference to it of the organism *Lamprocystis roseopersicina* under the name *Microhaloa aeruginosa* Kutzing.

The name is rejected by Erwin F. Smith (1905).

Micromyces. A generic name first applied to a fungus parasitic upon algae by Dangeard (1888, p. 55) with the species *M. zygonum*. Saccardo gives the following generic diagnosis:

Sphaerulae plasmaticae intracellulae Algae parasiticae, membrana aculeata verrucosa. Cellulae maturitate evacuantur et plasma, membranam tangens, sporangium compositum efficit, sporangium 4-cellulare, cellulae 100 zoosporas uncinatas gignentes.

The name *Micromyces* was independently proposed by Gruber as a generic name for a new organism which was isolated from man and showed great resemblances to *Actinomyces* (1891, p. 648). The species was named *Micromyces Hoffmanni*. This name as applied to one of the thread bacteria is invalid because of the previous use of *Micromyces* Dangeard for a distinct group. It is also a synonym of *Actinomyces*. It is rejected by Merrill and Wade (1919, p. 63).

Microphyta. A name included in the list of those rejected by Erwin F. Smith (1905). No reference to its use as a generic name has been found.

Microsiphonales. An order of fungi named by Vuillemin (1912). Castellani and Chalmers (1919, p. 1040) give the following description:

Definition. *Hyphales* with the mycelium composed of fine bacilli-form hyphae, usually 1 micron or less in diameter. Usually Gram-positive, when young, and without distinct nuclei. Parasitic on man, animals, and plants, or saprophytic.

Remarks. This is a most interesting order, as its members are nearly always mistaken at first sight, by persons unaccustomed to them, for bacilli. Hence the

descriptions of bacilli as the cause of so many fungal diseases due to these organisms, among which could be classified the diphtheria organisms, the tubercular and leprotic bacilli

It is included in Vuillemin's Subclass *Hyphales* It includes the two families *Nocardiaceae* and *Mycobacteriaceae*

Microsiphones. A group of the higher or filamentous bacteria recognized by Vuillemin (1913, p 526) In this group the organisms consist of branched filaments which break up into bacilliform elements. The filaments may either be uniform in diameter or inflated at the tips. Sometimes the filamentous forms predominate over the bacilli, sometimes they are rare and the rods divide indefinitely in the manner of the true bacteria The rods are non-motile, and less fixed in habit than with the true bacteria The following genera and type species are recognized

Nocardia Trevisan 1889

N Foersteri

Pasteuria Metschnikoff 1888

P ramosa

Sclerothrix Metschnikoff 1888

S tuberculosis

Corynebacterium Lehmann and Neumann

C diphtheriae

Microsphaera. A genus of pyrenomycetous fungi described by Leveille (1851, p 381). Cohn (1872a, p 237) named an organism which he observed in the lymph of smallpox vaccine *Microsphaera vaccinae* Later (1872, p 150) he corrected the error, using instead the generic name *Micrococcus* *Microsphaera* as a bacterial genus is invalid Cohn's description of the genus is as follows

Microsphaera (Kugelbacterie) Familie der Schizomyceten, Gruppe der *Bacteriaceae* Zellen farblos, sehr klein, kuglig oder spharoidisch, in der Regel unbewegt, durch Theilung oder Aschnurung zu 2, 4-8 oder mehrgliedrigen rosenkranzformigen Ketten vermehrt, durch Verschiebung zu unregelmässigen Gruppierungen, in Folge fortgeschrittener Vermehrung zu zahlreichen Zellhaufchen oder Colonien entwickelt, oder durch schleimige Intercellular substanz zu zoogloea artigen Massen verbunden, vielleicht Dauerzellen bildend

Microspira A bacterial genus named by Schroter (1886, p 168) with the following description

Vegetative Zellen schwach gekrümmt, meist nur mit $\frac{1}{2}$ Windung (Comma-Form), lebhaft bewegt Die kurzen Formen können, nachdem sie zur Ruhe gekommen, zu schraubigen Spirochaete-artigen

Faden auswachsen und Sporen bilden, welche durch Theilung der Gheder entstehen (Arthrosporen)

The type species was *M comma*, the organism of Asiatic cholera. Other species were *M Finkleri* and *M buccalis*.

Schroter was evidently in ignorance of the previous publication a year earlier of the generic name *Pacina* (Trevisan 1885, p 83) for the same group, which rendered his genus name invalid

Attention was called to the priority of *Pacina* by De Toni and Trevisan (1889, p 1018) Lehmann and Neumann (1896, p 107) rejected the name as an invalid synonym of *Vibrio*

The name *Microspira* was also adopted by Migula (1894, p 237) for a genus of his family *Spirillaceae*. He used the following description "Zellen starr mit 1, seltener 2-3 polaren, wellig gebogenen Geisseln". The comma bacillus was again the type

Migula says (1884, p 237) *Microspira* Schroter em Migula Zellen starr, mit 1, seltner 2-3 polaren wellig gebogenen Bewegungsorganen (Z B *Microspira Komma* Schroter)

In this characterization of the genus Migula was followed by Chester (1899, p 63), Migula (1900, p 31), (1900, p 960), Chester (1901, p 332), Kendall (1902), Smith (1902, p 270), Migula (1904, p 145) Ellis (1909, p 6), Frost (1911, p 59), Meyer (1912, p 4), Schneider (1912, p 23) and others

He was not followed by Fischer (1897) who used the genus name *Vibrio*, nor by E F Smith (1905, p 161) who likewise used *Vibrio*, nor by Blanchard (1906, p 1) and others. Vuillemin (1913, p 521) believes the generic name *Microspira* would be preferable to *Vibrio* in the sense of Cohn providing the relative lengths of the cells would serve to differentiate the genus from *Spirillum*

For additional discussion of priority of *Pacina*, *Microspira* and *Vibrio* see the latter

Microspironema. Stiles and Pfender proposed (1905, p 326) this generic name for the organism causing syphilis (to replace *Spiro-nema*) in the following statement

The parasite which Schaudinn found in syphilitic patients was first classified by him as a member of the genus *Spirochaeta* Cohn, 1872, this name was simply an emended spelling for *Spirochaeta* Ehrenberg 1834

Vuillemin (1905, June 5, pp 1567-1568) removed Schaudinn's organism from the genus *Spirochaeta* and proposed for it the genus *Spiro-nema*, a change in classification which Schaudinn has accepted. The generic name *Spiro-nema* is pre-occupied in Zoology, having been used by Meek in 1864 for a genus of Mollusks. It is accordingly not available for the protozoon of syphilis and under article

34 of the International Code of Zoological Nomenclature, we herewith propose the generic name *Microspironema* as substitute, with *Microspironema pallidum* (Schaudinn, 1905) Stiles and Pfender, 1905, as type species

This was proposed later, however, than *Treponema* of Schaudinn (1905, p 1728) and is hence invalid It has not been used by subsequent writers It may be regarded as a synonym of *Treponema*

Microspora. A name listed by Erwin F Smith (1905) in his genera to be rejected The original reference for this has not been found Perhaps it is a variant spelling of *Microsporon* (q v)

Enlows (1920, p 60) ascribes the term to Beijerinck and states "Variant of *Microspira* Schroter *M tyrosinatica*"

Microsporon. A generic name first used by Klebs (1871, p 33) for the bacterial species *M septicum* This is regarded as a very polymorphic organism which was responsible for suppuration in wounds, etc. His descriptions of the organism are misleading for he did not have pure cultures His theory of pleomorphism was soon discarded and the generic name generally regarded as *nomen nudum* It has not been used by subsequent authors in the original sense *M septicum* is given as a synonym of *Streptococcus pyogenes* by De Toni and Trevisan (1889, p 1062) This genus is generally accepted (sometimes with spelling *Microsporum*) for a genus of molds parasitic upon man

Conn (1909, p 12) describes *Microsporon* (apparently as a valid genus) as "Bacilli with one flagellum at each end" This is evidently a misstatement

Microsporon may be regarded as invalid as a bacterial generic name

Microzoa. Listed by Erwin F Smith (1905) as a non-valid generic designation for bacteria

Microzymba. A generic name used by Bechamp (1868, p 1160) for the parasite of the silk worm, *Microzymba bombycis* Schioeter (1889, p 150) gives an earlier reference to Bechamp (1867, p 42) but an examination of this fails to show the use of the word *Microzymba* Trevisan (1879, p 142) also credits the name *Microzymba* du lait to Bechamp This name has never been used as a generic designation by other authors

The *Microzymba bombycis* is made a synonym of *Streptococcus bombycis* by Schioeter (1886, p 150) The name is rejected by Erwin F. Smith (1905) Enlows (1920, p 60) lists two species, *Microzymba cretae* and *M bombycis* as described by Bechamp (1867, p 231)

Microzymba. A name used by various earlier authors as equivalent to bacteria (see Burdon-Sanderson 1871, p 323) A casual name

Mikrobakterien. A family name used by Flugge (1886) to include the non-filamentous bacteria. It is an aberrant spelling of *Microbacterien* (*q v*)

Mikrococcus. A variant spelling of *Micrococcus* as used by certain German authors, as Unna (1889, p 393).

Mikrokokkus. A spelling of *Micrococcus* used by Hueppe (1886, p 145) to designate an "Untergattung" of his *Arthrokokkaceen*. Other German writers have also made use of this name. While it is appropriate that *Mikrokokkus* be used as a casual designation, it should not be used in combination with a species name as a specific designation as it is not a Latin word.

Mikromyces. A variant spelling of *Micromyces* *q v.* used by Gruber (1891)

Mikrospironema. A variant German spelling of *Microspironema* *q v* (as Gonder 1914, p 690)

Mikrozymas. A variant German spelling of *Microzyma* *q v* (as Frenzel 1884, p 49)

Modderula. A genus of bacteria described by Frenzel (1897, p 802). The organism was observed first in dishes in which "Modder" was kept in the laboratory. "Modder" is the name given by the North German fisherman to the slime in the bottom of certain of the slow flowing waters of that region. The specific name given to the organism is *Modderula hartwigi*. The organism is ellipsoid, sometimes longer or almost spherical. It varies considerably in size, but is among the largest organisms ever included with the bacteria. In length it is 25 to 30 μ , in diameter 9 to 30 μ . The cell shows a double contoured colorless membrane. The interior of the cell shows a relatively compact mass of strongly refractive globules 1 to 2 μ in diameter. They have much the same appearance as the sulfur granules of *Beggiatoa*, and are interpreted in the same manner. The organism is observed to be slowly motile, but the organ of motion was not determined. Frenzel concluded that the closest affinities of the organism were with the bacteria.

Lauterborn (1898, p 95) pointed out that the organism described by Frenzel was not a new species, but a form which in 1893 had been shown by Lauterborn to Dr W Schewiakoff, who described it (1893) as *Achromatium* (*q v*). A careful comparison of the descriptions led to the conclusion that the organisms are identical, *Modderula* therefore lapses into synonymy. According to Bersa (1920, p 231) it is also a synonym of *Hullhousia* West and Griffiths.

Monas. A genus of protozoa created by Mueller (1786, p 1) The description reads "Vermis inconspicuus, pellucidus, punctiformis" As originally described this genus included 31 species of which a number were eventually separated as bacteria Ehrenberg (1838) separated off certain of Muller's species to form his genus *Bacterium* Ehrenberg (1848) in a study of the organism producing red spots in bread etc, named the causal organism *Monas prodigiosa* Since that time this generic name *Monas* has rarely been used by authors for bacteria. An exception is to be found in the use by Klebs (1875, p 472) of the name *Monas pulmonale* for a coccus which he found in human lungs *Monas* is still recognized as a valid protozoan genus It is invalid as a bacterial designation

Stiles (1905, p 30) gives an account of the type of the protozoan genus, which regards as *Monas mica* Muller

Monobacillus. A name used by Maggi (1886, p 174) to designate a growth form of *Bacillus*

Monobacteria A name applied by Billroth (1874, p 16) to a growth form of his polymorphic *Coccobacteria septica* in which the organism assumed a rod shape and occurred singly It apparently has never been used as a valid generic designation It is rejected by Erwin F Smith (1905)

Monococcus. A name applied by Billroth (1874, p 5) to a growth form of his *Coccobacteria septica* which consisted of isolated spherical cells It has not been used as a valid generic name It is rejected (with the spelling *Monococcus*) by Erwin F Smith (1905)

Monococcus. A variant of *Monococcus* Billroth listed by Smith (1905) The name was used by Miguel (1891) for those cocci whose cells tend to remain isolated

Monotricha. The first group of the *Gymnobacteria* named by Messee according to Kruse It includes bacteria with a single polar flagellum

Monovibrio. A casual name used by Maggi (1886, p 176) to designate a growth form of *Vibrio*

Multifermentans. A name given by Heller (1922, p 16) to the sixth genus of the tribe *Clostridioideae* The following is the description.

Clostridioideae that do not liquefy gelatin Produce gas and acid in meat medium Clot milk readily, without stormy fermentation Rather small Gram-positive rods with oval central or subterminal spores Actively saccharolytic Found occasionally in wounds, not tissue invaders

Type species *M. tenalbus* (*Bacillus multifermentans-tenalbus* Stoddard) as described by Stoddard (1915b). This organism, of which Dr Stoddard was so kind as to send me a culture does not fit into any other of the genera here defined

Stoddard's organism ferments glycerol, maltose, lactose, raffinose, glucose, sucrose, inulin, and salicin Mannitol and dulcitol are not fermented

Mycacanthococcus. A generic name proposed by Hansgirk (1890, p 26) for a species *M cellaris* Hansgirk

The description given by Hansgirk is as follows

Zellen kugelig in vegetativem Zustande mit glatter, in encystirtem Zustande mit kurzen stachel-oder warzenartigen Auswuchsen (ähnlich wie bei der Algengattung *Trochiscia* Ktz (*Acanthococcus* Lagrh) versehener Membran

Veget Zellen ohne Hülle meist 3 bis 6, seltener bis 8, oder nur 1 bis 2 μ mit der Hülle 5 bis 15 μ breit, einzeln oder zu 2 bis 8 in fast kugeligen, meist 25 bis 24 μ breiten Familien vereinigt, mit in der Regel völlig homogenem, farblosem, sehr stark lichtbrechendem dichtem, plasmatischem Inhalte, welcher erst zur Zeit der Sporenbildung feinkörnig wird Zu dieser Zeit tritt das Feinkörnigwerden des Plasmas zuerst blos an der Peripherie des ganzen Zellplasmas gleich auf, erst später ist der ganze plasmatische Inhalt, zuerst gleich fein, später grober gekörnt

Vermehrung erfolgt durch veget Zweitheilung der Zellen innerhalb der Mutterzellkapsel in 2 bis 8 Tochterzellen, welche bald wieder jede je eine Separatkapsel ausscheiden Charakteristisch für diese Bacteriengattung ist auch die tetraedrische Anordnung der Tochterzellen in vierzellige Familien, welche wieder meist traubenförmig neben einander gruppirt sind

Während junge Zellen zuerst noch von keiner Specialhülle umgeben sind, besitzen ältere vegetative Zellen eine deutliche knorpelartige, nicht zerfließende, meist ziemlich weit (ähnlich wie bei der Algengattung *Capsulococcus* Bennet) abstehende Specialhülle (Cyste), welche an der Oberfläche glatt, im Dauerzustande aber stumpfwärzig verdickt und weniger durchsichtig ist, als im veget Zustande

Die acanthococcusartigen Dauerzellen sind im Querprofil rundlichachteckig, 6 bis 8 μ dick (Zelllumen etwa 5 μ breit), an der Oberfläche der etwa $\frac{1}{2}$ dicken, fast farblosen membran mit 8 sehr kurzen, stumpfen, farblosen Emergenzen versehen, ihr Inhalt ist stark lichtbrechend, feingekörnt, farblos Ausser diesen Dauerzellen, welche zuerst noch von einer oftters zweischichtigen, farblosen Gallerthülle umgeben sind, mit welcher sie bis 10 μ breit werden, habe ich in grösseren (seltener auch in kleineren) vegetativen Zellen noch stark lichtbrechende und scharf contourirte Kugeln mit ganz homogenem Inhalte beobachtet, deren Durchmesser 1 bis 2 μ in sehr grossen Zellen bis 6 μ betrug, deren Bedeutung mir jedoch noch unbekannt geblieben ist

Diese seltene Kellerbakterien-Gattung kommt meist im Gallertlager anderer unterirdischer Bacterien zerstreut vor, so auf feuchten Mauern in alten Weinkellern auf der Altstadt in Prag, dann in der Pleissenburg und in der Centralhalle in Leipzig

The genus has apparently not been noted by subsequent authors It is valid if the organism described is deserving of generic rank

Mycobacillus. A generic name used by Chantemesse, Matruchot, and Grimberg (1917, p 652) for a species *Mycobacillus synovialis*. Enlows (1920, p 62) has summarized the description as follows

Intermediate between the "micromycetes" and true bacilli, hence the name. Young cultures actively motile Rod-like organism Spores by "enkystemment partiel" The older filamentous growths lose their motility Certain portions of the filaments are Gram-positive, others Gram-negative Isolated from cerebral ventricle in case of acute arthritis

Mycobacteriaceae. A family name proposed by Chester (1897, p. 37) with the genus *Mycobacterium* Lehmann and Neumann as a type. It has been used by Chester (1897 and 1899) with the following description

Cells short or long cylindrical or clavate-cuneate in form without a sheath surrounding the chains of individuals Without endospores, with possibly the formation of gonidia-like bodies With true dichotomous branching (this forms an important group standing between the true *Bacteriaceae* and the true fungi on the one hand and the *Chlamydobacteriaceae* on the other)

The genera *Mycobacterium* and *Corynebacterium* are included

The definition was somewhat altered by Chester (1901, p. 349) later to include the *Actinomyces* forms The diagnosis reads

Cells either short or long, cylindrical-clavate-cuneate in form, which at times may show true branching or as long branched mycelial-like filaments Filaments not surrounded by a sheath as in *Chlamydobacteriaceae* Without endospores, but with the formation of gonidia-like bodies due to a segmentation of the cells Division at right angles to the axis of a rod or filament

The family is recognized by A. J. Smith (1902, p. 270) Schneider (1912, p. 25) Benecke (1912) and by Miehle (1908, p. 131) The last named author evidently is unaware of the previous use of this name by Chester, for he proposes it as new. He places it in his scheme of classification following *Coccaceae* and *Bacteriaceae* and before *Spirillaceae* He included

Einzellige Individuen von sehr unregelmässiger Stäbchenform, auch in normalen Zustand sprossende Verzweigung und zwar regelmässig oder nur gelegentlich, keine Myzelbildung nach Art der Fadenpilze, zuweilen fadige Entwicklung der Individuen, keine Endosporen, keine Bewegung

Winslow *et al* (Committee Soc. Am. Bact., 1917, p. 553) included this family under the *Eubacteriales* with the description

Cells usually elongated, frequently filamentous and with a decided tendency to the development of branches, in some genera giving rise to the formation of a definite branched mycelium Cells frequently show swellings, clubbed or irregular shapes Endospores not produced, but conidia developed in some genera Usually Gram-positive Non-motile Many species are parasitic in animals or

plants Complex proteins usually required As a rule strongly aerobic (except for some species of *Actinomyces* and the genera *Fusiformis* and *Leptotrichia*), and oxidative Growth on culture media often slow, some genera show mold-like colonies

In 1920 the Committee included this family in the order *Actinomycetales* with the description

Parasitic forms Rod shaped, frequently irregular in form but rarely filamentous and with only slight and occasional branching Often stain unevenly (showing variations in staining reaction within the cell) No conidia

Castellani and Chalmers (1919, p 1040) note that this family (credited to Miehle 1909) may be placed as the second family of *Microsiphonales* Vuillemin with the definition, "Microsiphonales without a mycelium," and including the two genera *Mycobacterium* and *Corynebacterium*

These authors, however, themselves place the family as the fifth in the order *Eubacteriales*, and credit the name to Chester 1901 They follow Chester also in reducing *Corynebacterium* to synonymy with *Mycobacterium*

Bergey *et al* (1923, p 372) follow the Committee, including the family as the second in the order *Actinomycetales*

Mycobacteriaceae. A tribe named by Buchanan (1918, p 55) with the following description

Rod-shaped organisms occasionally showing a tendency to branching, acid-fast, Gram-positive, non-motile, without spores

A single genus is included, *Mycobacterium* Lehmann and Neumann

Mycobacterium. A generic name proposed by Lehmann and Neumann (1896, p 363) with the organism of tuberculosis as a type, *Mycobacterium tuberculosis* It is included in their family of the *Hyphomycetes* In the translation of the second edition (1901, p 128) the description reads

Rods stain with usual staining solutions with difficulty or generally not at all Stain by the tubercle bacillus method, i e , it is acid resisting Clubbed swelling of the ends in cultures rare, in tissues somewhat more often

In this edition is found the following statement regarding the priority of *Sclerothrix*:

Since we proposed this name in the first edition, we have seen that Metschnikoff (Virchow's Archiv, 113, p 70, 1888), who first recognized the peculiar position

of the tubercle bacillus as opposed to the other then known bacteria, in a work "Regarding the Phagocytic Role of the Tubercular Giant Cell," has said "If one considers that in the perfected stage the tubercle bacteria have grown into (although short) threads, and also differ from other analogous forms (except the lepra bacteria) in the possession of a very dense envelope, then perhaps it will be easier to accept the designation *Sclerothrix* for the genus, and *Sclerothrix Kochii* for the species of the tubercle bacterium" We should have immediately accepted these names if we had known of them, but believe that according to the rules of botanical nomenclature our names should now stand, since Metschnikoff only made a conditional proposal, did not accurately define his new genus, and never made any use of the new name himself, while we have ourselves already established a name

Neither Lehmann and Neumann or Metschnikoff seem to know of the prior use of *Sclerothrix* (*q v*) as an algal genus by Kutzing (1849, p 37)

Chester (1898) redefined *Mycobacterium* as follows

Cells in their ordinary form short cylindrical rods, often bent and irregularly swollen, clavate or cuneate At times Y shaped forms or longer filaments with true branchings May produce short coccoid elements, perhaps gonidia

The genus with this definition includes the *Corynebacterium* of Lehmann and Neumann.

The genus is listed as a synonym of *Bacterium* by Migula, (1900, p 492)

Chester (1901, p 349) again used this genus to include the acid fast and the diphtheroid bacteria The genus is also used by A J Smith (1902, p 270) and by Kendall (1902)

The genus *Mycobacterium* was also recognized by Miehle (1908, p 131) who emended the description somewhat He included in it those forms with rod-shaped cells, usually somewhat bent, with variable width, and length After division the cells usually bend to one side, and grow side by side, forming a characteristic very firm colony which is not easily broken to pieces, and is easily separated from the medium, usually forming a membrane on liquid media Budded branching occasional Usually acid fast, non-motile, slow growth, no liquefaction of gelatin

Vuillemin (1913, p. 527) rejects the name as a valid genus because of the priority of both *Sclerothrix* and *Coccothrix* It has already been noted that the former name is invalid, but it is possible that if the acid fast bacteria are to be recognized as a separate genus the generic designation *Coccothrix* should be used, which would make *Mycobacterium* an invalid synonym

Winslow *et al* (Committee Soc. Am. Bact. 1917, p. 554) have used the definition

Slender rods which are stained with difficulty, but when once stained are acid-fast. Cells sometimes show swollen, clavate or cuneate forms, and occasionally even branched filaments. Non-motile, Gram-positive. No endospores. Growth on media slow. Aerobic. Several species pathogenic to animals.

The type species is *Mycobacterium tuberculosis* (Koch) Lehmann and Neumann.

Buchanan (1918, p. 55) gave the following definition and discussion

Slender rods which stain with difficulty, but when once stained, are acid-fast. Clubbed, swollen, clavate or cuneate cells occur, even filaments with branches. Non-motile. Without spores. Gram-positive.

The type species is *Mycobacterium tuberculosis* (Koch) Lehmann and Neumann.

Vuillemin (1913, p. 527) contends that *Coccothrix* resembles too closely *Coccolithum* Wallroth. It does not seem that the resemblance is close enough ever to cause confusion, and the genus name *Coccothrix* is apparently valid. The Committee on Classification of the Society of American Bacteriologists, however, has recommended the use of the generic name *Mycobacterium* as better known and less likely to result in confusion.

Castellani and Chalmers (1919, p. 962) indicate as the type species *Mycobacterium leprae* (Hausen 1874), and include the diphtheria and glanders types (*Corynebacterium*) in the genus.

Bergey *et al* (1923, p. 372) follow the Committee, and include the genus as the first in the family *Mycobacteriaceae*.

The validity of *Mycobacterium* as a valid generic name is menaced apparently only by the older *Coccothrix* Lutz, *q v*. Perhaps inasmuch as the latter name has apparently never been used, it may be passed by in the interest of stability.

Mycococcaceae. A family of bacteria proposed by Hansgirk (1888, p. 266) as the only family of his order *Sphaerobacteria*. Two sub-families are included, *Cystococcaceae* and *Eucoccaceae*.

Mycoderma. A generic name used by Persoon (1822, p. 96) with the species *Mycoderma ollare*, *M. mesentericum*, etc., the latter for the film which forms on beer or wine which is exposed to the air. The individual organisms which compose this scum were not studied. We know that this scum may be of yeasts or of bacteria, frequently a mixture of the two. It is not surprising therefore in the subsequent literature to find authors diverging in the use of the term *Mycoderma*, one group regarding it as a genus of yeasts, the other as a genus of bacteria.

Desmazieres (1826, p 59) named the layer which develops on beer, etc. *Mycoderma cerevisiae*, regarding the organisms, however as *animalcula monadina*. According to Lafar he did not connect the organisms with the chemical changes taking place in the solution.

Kutzing (1837, p. 386) discusses the use by Desmazieres of *Mycoderma*, and later (1843, p 149) substituted for the acetic acid form the generic name *Ulvina*.

According to Hansen (1894, (1911), p 465) the designation *Mycoderma aceti* was used by Thomson (1852). This latter author added yeast and a little bread to a sugar solution. In the course of three months a satisfactory vinegar had developed, and in the bottom of the vessel there was a considerable gelatinous deposit which he identified as the *Ulvina aceti* of Kutzing. He states that this organism is also termed *Mycoderma aceti*, though Hansen was unable to determine his authority for the statement.

Pasteur (1868, p 106) used the name *Mycoderma aceti* for the organism present in acetic fermentation.

This name was also used by Hansen (1879, p 230) as a generic bacterial designation, for the species *M. Pasteurianum*.

The name has come into general use for certain of the yeast like fungi. However, its use for an acetic organism was suggested by Winslow *et al* (1917, p 551) with the following description:

Cells rod-shaped, frequently in chains, non-motile. Cells grow usually on the surface of alcoholic solutions, securing growth energy by the oxidation of alcohol to acetic acid. Also capable of utilizing certain other carbonaceous compounds, as sugar and acetic acid. Elongated, filamentous, club-shaped, swollen and even branched cells common and quite characteristic.

The type species is *Mycoderma aceti* (Thompson?) Committee.

Buchanan (1918, p 45) used the following description:

Cells rod-shaped, frequently in chains, non-motile usually, without spores. Obligate aerobes, growing usually as a film on the surface of alcoholic solutions, transforming the alcohol to acetic acid. Involution forms often developed and quite characteristic.

The type species is *Mycoderma aceti* Thompson?

There is some doubt as to the appropriateness of *Mycoderma* as the name of this genus. It is possible that it should be reserved for the yeasts.

Enlows (1920, p 62) calls attention to the fact that the type species "(first in order of arrangement, and subsequent designation by many authors)" is *M. ollare*.

The Committee (1920) abandoned *Mycoderma* as a generic bacterial name, substituting *Acetobacter q v.*

The name is rejected by Orla-Jensen (1921, p 270)

It is probable that *Mycoderma* is invalid as a designation for a bacterial genus, and should be reserved for certain pellicle forming yeasts

Mycomonas. A generic name proposed by Orla-Jensen (1909, p. 329) to replace *Mycobacterium* Lehmann and Neumann to make the designation conform to his principles of nomenclature It is to be regarded as a synonym of *Mycobacterium* and *Coccothrix*, and invalid

Myconostoc. A genus of bacteria described by Cohn (1875, p 183) to include a species *Myconostoc gregarium* This organism was observed in water as gelatinous masses containing cylindrical colorless cells, frequently curved or spiral Found in water kept in glass with decaying algae Water became dark through iron sulphid

An der Oberfläche sammelten sich farblosen Schleimtropfen, diese waren gebildet von isolirten, oder haufenweise am einander hangenden kleinen Gallertkugeln von 10–17 μ Durchmesser und daruber Diese Kugeln, nach aussen ziemlich scharf abgegrenzt, haufig elliptisch in die Lange gezogen, schlossen in einer durchsichtigen Gallert einen farblosen *Leptothrix* artigen Faden ein, welche in knauelartigen aber lockeren Windungen ins Innere eingelagert war Ob jede Kugel immer nur einen oder auch mehrere solcher Faden einschliesst, lasst sich nicht leicht ermitteln, obwohl ich das eistere als Regel vermuthe, unmittelbar vermag man nur die kogenartigen Schlingen in der Peripherie, und die durch einander geschlungenen Wundungen im Innern zu unterscheiden Die farblosen Faden selbst etwa von der Starke des *Bacillus ulna* oder *Spirillum volutans*, enthalten stark lichtbrechende Kornchen, Gliederung ist nicht erkennbar Die Vermehrung geschieht, ahnlich wie bei *Ascococcus* vermittelst Querfurchung der Gallertkugel, die, vermuthlich in Folge bedeutenderer Streckung des eingelagerten Fadens, sich erst elliptisch in die Lange dehut, dann in der Mette sich in zwei Halbkugeln durchfurcht, welche sich nach kurzer zeit von einander trennen Der Gedanke lag nahe, dass es Spirillen seien welche hier im Gallert eingeschlossen sind

The genus was included in the tribe *Nematogenae*

Strangely enough, there appears to be but one subsequent record of the discovery or observation of this organism The genus however, was generally recognized during the first two decades following its description

Trevisan (1879, p 139) gives the following generic description "*Somatia cylindrica, tenerrima, implicata, convoluta, muco obvoluta, in globulos perparvos congesta* " This was included in his subtribe *Myconostocaceae*

The same description was used by Magnin (1880, p 96) *Myconostoc* according to Van Tieghem (1884) consisted of spiral cells embedded in gelatin

Winter (1880, p 60) gave the following description

Faden sehr dünn, farblos, ungegliedert, aber beim Eintrocknen in kurze cylindrische Glieder zerfallend, verschiedenartig gebogen und durch einander geschlungen, von Gallerte umgeben, die rundliche Kugeln von 10–17 μ (und mehr) Durchmesser bildet Vermehrung durch Einschnürung und Zweitheilung dieser Gallertkugel

Schroeter (1886, p 169) gives the following diagnosis

Zellen in Form dünner, gebogener und gewundener Faden, welche in eine rundliche Gallertcyste eingeschlossen sind Die Faden zerfallen in kürzere Stücke Die Cysten vermehren sich durch fortgesetzte Zweitheilung

Hansgirk (1888, p. 265) includes the genus as the sole genus of his family *Myconostocaceae*, and notes finding the organism among algae in Bohemia, and frequently among his algal cultures

The genus is accepted by De Toni and Trevisan (1889, p 1034), Trevisan (1889, p 26), Billet (1890, p 24) and Ludwig (1892)

Migula (1900, p 960) made *Myconostoc* the second subgenus under his genus *Spirosoma*

The genus is rejected by Erwin F Smith (1905) and by most recent writers It is proposed by Vuillemin (1913, p 526) as one of his "Formogenera conservanda"

If the organism described by Cohn is worthy of generic recognition, *Myconostoc* is a valid bacterial genus

Myconostocaceae. The second family of the order *Eubacteria* proposed by Hansgirk (1888, p 265) in the following statement

Diese noch von Schroter mit der Familie der Bacteriaceen vereinigte Spaltpilzgruppe, welche der Familie der Nostocéen unter den Spaltalgen entspricht ist aus ähnlichen Gründen wie die *Crenothrichaceae* zu einer den übrigen Familien der Bacterien gleichwerthigen Familie zu erklären

A single genus, *Myconostoc*, is recognized

Myconostocaceae. A subtribe created by Trevisan (1879, p. 139) to include the genus *Myconostoc* He gave the following characterization "Somatia substantia intersomatium in colonias mucosas consociata, eramosa" This is the second subtribe included within his tribe *Vibrionaeae*

De Toni and Trevisan (1889, p 1934) again used the term for a subgenus with the following description "Baculi spiralter torti, capsulis inclusi"

Mycophyceae. A name probably of ordinal rank proposed by Hansgug (1890, p 31) to include those bacteria which are intermediate between the algae (the *Schizophyceae*) and the *Eubacteria*. The families *Crenotrichaceae*, *Leptotrichaceae*, *Myconostocaceae* and *Mycococcaceae* are included

Mycotetraedron. A generic name proposed by Hansgug (1890, p 28) for a species *M cellare* n sp. He gives the following description

Zellen einzeln, rundlich viereckig, die Ecken tetraedrisch gestellt 5 bis 6 μ breit, mit farblosem, homogenem oder stellenweise spärlich feingekorntem, fast wasserhellem Inhalte und dünner farbloser an der Oberfläche glatter Membran, and den Ecken abgerundet und daselbst mit je einem, 1 5 bis 2 μ langen, geraden, farblosen kegelförmigen Stachel versehen

Die Vermehrung und weitere Entwicklung dieser morphologisch der Algen-Gattung *Tetraedron* Ktz (*Polyedrum* Nag.) ähnlichen Kellerbakterie ist mir noch unbekannt, ich habe sie bisher bloß unter andern Kellerbakterien aus einem alten Weinkeller auf der Prager Altstadt in wenigen Exemplaren beobachtet

The genus apparently has not been recognized by subsequent writers. If the species described is sufficiently distinct, the generic name would appear to be valid.

Mycothamnion. A name of a genus of algae used by Kützting (1843, p 120) with the following description "Trichomata (inarticulata?) achromatica ramosa, rami apice in spermatia intumescens". One species, *M fodinarum* Kuetz. is a synonym of the bacterial species *Leucocystis cellaris* of Schroeter (1886) according to De Toni and Trevisan (1889, p 1041).

Mycotheca. A generic name used by Hansgug (1888, p 266) with the following description of the genus and the single species *M cellaris* Hansgug

Genus *Mycotheca* Hansg. Zellen länglich cylindrisch, einzeln oder in zweibis vierzelligen Familien, von einer meist mehrschichtigen deutlich begrenzten farblosen Gallerthülle umgeben, unbeweglich, einzeln oder zu einer dicken, formlosen Masse vereinigt. Vermehrung durch fortgesetzte vegetative Zweitheilung der Zellen bloß in einer Richtung des Raumes (der Quere nach), wobei die Tochterzellen sich bald von einander trennen. Sporenbildung unbekannt.

M cellaris Hansg. Zellen cylindrisch, gerade oder leicht in der Mitte gekrümmt, meist 1 bis 1 5, seltener 2 bis 3 μ dick, drei- bis sechs-, seltener nur zweimal so lang, stark lichtbrechend, farblos, einzeln, selten zu zwei bis vier, von deutlich begrenzten, 4 bis 5 μ dicken, oft geschichteten hyalinen Gallerthüllen umgeben, dicht zu

gelblichen bis gelbraunlichen Gallertmassen gehaut oder im schleimigen Lager anderer Kellerbakterien verstreut

An wenig feuchten Mauern in alten Weinkellern, meist in Gesellschaft des *Bacillus subtilis* var *cellaris*, so auf der Altstadt und in einem Weinkeller auf der Neustadt

De Toni and Trevisan (1889, p 1033) make the species *Mycotheca cellaris* Hansgig a synonym of *Klebsiella cellaris* Trevisan

The name has apparently not been used by other authors It is rejected by Erwin F Smith (1905) If the organism described is worthy of generic differentiation, it would appear that the genus name is valid

Mycothrix. A name applied to chains of spherical organisms by Itzigsohn (1867) Cohn (1872, p 147) makes the following statement "Itzigsohn und Haller haben für die Rosenkranzketten der Kugelbakterien den Namen *Mycothrix* vorgeschlagen, ich bezeichne sie hier als Torulaform"

Apparently the term has never been used in a strict generic sense The nearest approach seems to have been the ascription by De Toni and Trevisan (1889, p 1059) to Friedberger of the term "*Mycothrix* bei Pleuropneumonie des Pferdes" (1873, p 91) An examination of the latter reference fails to show any such use It is invalid

Myxobacillus. A generic name used by Gonnermann (1907, p 877) (with the spelling *Myxobazillus*) for a species of bacterium, *M betae* isolated from sugar beet juice The organism is a slender rod, 0.3 by 2.3-4.5 μ Easily stained Often granules No definite capsule Grows at 35°C, but not killed at 60° Non-motile It forms spores in twenty-four hours which are oval, of a diameter greater than that of the mother cell. Before spore formation the organism occurs in chains like anthrax bacillus, the filament falls apart into segments each of which produces a spore It grows well on ordinary media In sucrose solutions gas and acid are not formed The solution becomes opalescent and viscous Apparently this is a synonym of *Bacillus*

Myxobacter. A generic name proposed by Thaxter (1892, p 493) for a species of the *Myxobacteriaceae*, *M aureus* The following generic description was given "Rods, forming large rounded cysts, one or more free within a gelatinous matrix raised above the substratum" Thaxter later noted that this generic name is invalid because of the prior use of *Polyangrium* q v for this form This name has accordingly been abandoned in later writings on this group

Myxobacteriaceae. The name of a family proposed by Thaxter (1892, p 394) to include the so-called *Myxobacteria*. The name comes from the genus (now abandoned) *Myxobacter* Thaxter.

The family diagnosis as given by E F Smith is as follows (1905, p 164)

Motile, rod-like organisms, multiplying by fission secreting a gelatinous base, and forming pseudoplasmodium-like aggregations before passing into a more or less highly developed cyst-producing, resting state in which the rods may become encysted in groups without modification, or may be converted into spore-masses. The vegetative rods, which vary little in size and form in the different genera and species, are typically elongate, sometimes reaching 15μ in length. Cell division follows an elongation and nearly medium constriction of the rods which, except at the moment of division, are always separate, never united in chains. A slow, sliding locomotion and a Beggiatoa-like, bending motion is characteristic of the active rods. Organs of motion have not been detected. In all species, with one exception, the rods, when seen in masses, are more or less distinctly reddish. A distinct, firm, hyaline gelatinous base is secreted by the colony as it extends itself, over which the individuals may move or in which they become imbedded.

The vegetative period, in artificial cultures, usually lasts about a week, or even two weeks, but in nature the production of cysts must be more rapid. Common in moist situations on dung, decaying wood, fungi, lichens, etc. According to Bauer they grow best at 30°C .

In forms like *Myxococcus* in which the rods are somewhat scattered, the first preparation for spore production as seen under the microscope consists in the appearance of groups of rods moving with a circular tendency, in which the more central individuals soon become converted into the spores. The formation of a cystophore, when it occurs, results from the basal constriction of a papillate mass of rods which projects from the surface of the colony. In the encysted condition there are two classes—one in which the individuals thus encysted show little or no modification from the rod-like vegetative state, the other in which they are converted into definite spores. They, however seem to run into one another.

This family name is also used by Clements (1909, p 8) and by Benecke (1912).

Buchanan (1918, p 541) included this as the single family of the order *Myxobacteriales*, with characters those of the order, *qv*. Three genera were included *Chondromyces*, *Polyangium* and *Myxococcus*. Bergey *et al* (1923, p 412) follow Buchanan.

Myxobacteriales. An ordinal name proposed by Buchanan (1917, p 163). The description (1918, p. 541) follows

Motile, rod-like organisms, multiplying by fission, secreting a gelatinous base, and forming a pseudoplasmodium-like aggregation before passing into a more or less highly developed cyst-producing, resting state in which the rods may become encysted in groups without modification, or may be converted into spore masses.

There is one family only, the *Myxobacteriaceae*.

It is included as the fifth order of the *Schizomycetes*

Winslow *et al* (Committee Soc Am Bact, 1917, p 549, 1920, p. 196) recognized the order with the following diagnosis. "Cells united during the vegetative stage into a pseudoplasmodium which passes over into a highly-developed cyst-producing resting stage"

It is also recognized by Castellani and Chalmers (1919, p 924) as the third order of the *Schizomycetacea*, and by Bergey *et al* (1923, p 412) as the fifth order of the *Schizomycetes*

Myxobacterieae. A class name proposed by Heller (1921, p 549) for the second class of the phylum *Bacteria* She states "Bacteria which join to form a complex fruiting body"

Myxobacterium. Under the heading "A New Myxobacterium" an abstract was printed of a paper by Faull (1915, p 469) Apparently the word is not used binomially The organism evidently is one of the *Myxobacterales* (*q v*)

Myxobactrales. A name proposed by Clements (1909, p 8) for the second order of the *Schizomycetes* His description is "Cells rod-like, motile, fission in one plane, cells secreting a gelatinous base and forming pseudoplasmodia, then passing into cysts, or spore masses which are often stalked (cystophore)"

One family, *Myxobacteriaceae*, is included It may be regarded as variant of the spelling *Myxobacterales*, *q v*

Myxobacillus. A variant of *Myxobacillus*, *q v* used by Gonnermann (1908 p. 877)

Myxobotrys. A generic name of a fungus proposed by Zukal (1896, p 346) for the species *M variabilis* The generic description is as follows

Die Sporen in dichten Kopfchen auf der blasigen Erwerterung an den Enden eines einfachen oder schwach verzweigten Sporentragers, oder ohne Trager direct auf einem dunnen Hypothallus oder dem Substrate aufsitzend

It was later shown that this was a redescription of an old genus, one of the *Myxobacteriaceae*, *Chondromyces*, (*q v*) The name is therefore invalid

Myxobotrysaceae. A family created by Zukal (1896, p 436) to include his mold genus *Myxobotrys*, with the following description. "Die Sporen entstehen durch Sprossung des Plasmodiums, ohne vorhergehende Sonderung des Plasma in bestimmte Portionen" When it was shown that the genus *Myxobotrys* was invalid, and the organism did not belong to the fungi, this family name was invalidated

Myxococcus. A generic name proposed by Thaxter (1892, p 403) as a genus of the *Myxobacteriaceae*. The diagnosis given is "Rods slender, curved, swarming together after a vegetative period to form definite more or less encysted sessile masses of coccus-like spores." In the possession of the coccus-like spores when encysted it differs from the other genera of the family, *Polyangium* and *Chondromyces*.

Eleven species have been described, *Myxococcus rubescens* Thaxter, *M. virescens* Thaxter, *M. coralloides* Thaxter, *M. stipitatus* Thaxter, *M. currhosus* Thaxter, *M. cruentus* Thaxter, *M. pyriformis* A L Smith, *M. disciformis* Thaxter, *M. macrosporus* Zukal, *M. digitatus* Quehl, and *M. clavatus* Quehl.

The generic name *Myxococcus* has been accepted and used by Saccardo (1899, p 460) by Zukal (1897, p 550) and by Quehl (1906, p 16).

With the spelling *Myxokokkus* this name has been also used independently by Gonnermann (1907, p 877) for a species of bacterium *M. betae* isolated from sugar beet syrup. In this sense the name is invalid.

Buchanan (1918, p 542) gave the following description:

Rods slender, curved, swarming together after a vegetative period to form definite more or less encysted sessile masses of coccus-like spores.

The type species is *Myxococcus rubescens* Thaxter.

Bergey *et al* (1923, p 413) included this as the first genus of the family *Myxobacteriaceae*.

Myxokokkus. A generic name used by Gonnermann (1907, p 883) for a species *Myxokokkus betae* Gonnermann isolated from a gummy fermentation of sugar beet sap. The organism is a streptococcus. It grows on all ordinary media except potato, but usually not luxuriantly. It forms on dextrose agar no gum, but only small, at first transparent, discrete colonies with slight gas formation. On ten per cent sucrose agar and gelatin it produces abundant gum and vigorous gas. The organism quickly dies on this medium. It is possible that in nature development is dependent upon growth with a gum producing bacillus. The genus is probably invalid because of the priority of *Myxococcus* Thaxter, and because of the inadequate characterization. Possibly it is a synonym of *Leuconostoc* *q v*.

Neissereae. A tribe of bacteria named by Winslow *et al* (1920, p 205) as a tribe of the family *Coccaceae*. One genus only, *Neisseria* is included. The diagnosis given is "Strict parasites, failing to grow or growing very poorly on artificial media. Cells normally in pairs. Gram-negative. Growth fairly abundant on serum media."

Bergey *et al* (1923, p 41) includes this as the first tribe of the family *Coccaceae*

It should be noted that the spelling should be *Neisserieae*

Neisseria. A generic name proposed by Trevisan (1885, p 105) with the gonococcus of Neisser as the type The generic description is as follows "Colonie di 4 cocchi subovoidei, consociati a due in un unico strato ad avviluppato da densa mucosa, ialina, spianata a guisa di membrana quadrata, da ultimo libero" The single species listed is *Neisseria gonorrhoeae* Trevisan (1889, p 32) includes this as the first genus of his tribe *Micrococceae* Fourteen species are included, the first being *N gonorrhoeae* De Toni and Trevisan (1889, p 1067) give the following diagnosis

Cocci primitus globosi indivisi, aetate provecata in coccos duos biscoetiformiter geminos latere fratrem versus plus minus complanato, utrinque ad polos isthmis filamentosis tenuissimis insimul nexos, scissi, nunquam in turmas racemiformiter consociati Endosporeae microsomae, in coccis normalibus obvenientes

Twenty-four species are included in the genus The name *Neisseria* has rarely been used, usually being displaced by the term *Diplococcus* *Neisseria*, however, has priority over this name if the gonococcus is the organism which should be taken as the type of the genus, or is to be included within it. It is rejected by Erwin F Smith (1905) For discussion see *Diplococcus* Vuillemin (1913, p 525) has proposed that *Neisseria* with the type species *N gonorrhoeae* be made a *Genus conservandum*

Buchanan (1917, p 612) recognized the genus with the following description.

Parasites, growing poorly in most laboratory media Cells usually in pairs, flattened, coffee-bean shaped Gram-negative As a rule showing considerable fermentative power in carbohydrate media

The type species is *Neisseria gonorrhoeae* Trevisan

It was included as the fourth genus of the tribe *Streptococceae*

Winslow *et al* (Committee Soc Am Bact, 1917, p 557) used the following diagnosis

Strict parasites, failing to grow or growing very poorly on artificial media Cells normally in pairs of flattened cells Gram-negative Fermentative powers low Growth fairly abundant on serum media, usually whitish or yellowish

The type species is *Neisseria gonorrhoeae* Trevisan

The committee (1920, p. 205) made it the sole genus of the tribe *Neissereae*

Castellani and Chalmers (1919, p. 925) include the genus under *Streptococceae* with the diagnosis

Streptococceae growing best, and often only, aerobically, without pigment formation, usually present in pairs without a true capsule, and in exudates usually intracellular, and readily decolourized by Gram's method of staining

Primary cultures grow poorly on usual laboratory media, but best on media containing glucose or blood serum No lysis with bile Ferment carbohydrates usually with but slight acid production

Type species *Neisseria gonorrhoeae* (Bumm, 1885)

The genus *Neisseria* is recognized by Bergey *et al* (1923, p. 41) as valid, with the type *Neisseria gonorrhoeae* Trevisan

Nematogenes. A tribe of *Schizophyta* named by Cohn (1875) It included those forms in which the cells are disposed in filaments Several bacterial genera were included

Nevskia. A bacterial genus created by Famintzin (1892, p. 484) for the species *Nevskia ramosa* According to Migula (1900, p. 1038) the generic description is as follows

Stäbchenförmige Zellen, welche einseitig Gallerte aussondern und dadurch gestielt erscheinen Bei der Teilung des Stäbchens teilt sich auch der Stiel, so dass sich allmählich eine in der Form von *Chaetophora endiviaefolia* erinnernde gelappte Gallertkolonie bildet, die an den Enden der Lappen die Zellen trägt

The species *N. ramosa* is described as follows

Zellen etwa 12 μ lang, 2-6 mal so lang als breit, kugelförmige, stark lichtbrechende, sporenartige Gebilde enthaltend, die jedoch in Alkohol löslich sind Die umschliessende Gallerte lässt sich in 1-proz. Kalilauge nach vorhergegangener Behandlung mit Alkohol leicht lösen Teilung senkrecht zur Längsachse Unter bestimmten, nicht näher bekannten Verhältnissen verlassen die Zellen die Gallerte und werden frei, sie stellen dann den Anfang neuer Kolonien dar Im allgemeinen scheint diese Art Ähnlichkeit mit *Bacterium pediculatum* Koch und Hosaeus zu haben

Von Famintzin in Aquarien des botanischen Laboratoriums gefunden

Erwin F. Smith (1905) places this in his list of doubtful genera It is apparently valid if the organism described is worthy of generic separation

Newsia. A spelling of *Nevskia* used by Migula (1900, p. 1038) and Erwin F. Smith (1905, p. 162)

Nicolaerillus. A name applied by Heller (1922, p. 24) to genus of the tribe *Putrificoidae* The description follows:

Putrificoideae that in meat medium produce gas and various color changes, yellowish, pink, grey or mauve, depending on the medium, the particles of meat are gradually suffused with a black pigment, and bleach at the top. The meat is softened but the particles do not greatly diminish in size. Do not attack sugars. Gram-negative (weak methyl violet) rods that form terminal spherical spores. Colonies in deep agar diverse. Common in soil, found in horse feces, may multiply in wounds, but do not normally invade tissue. Produce a characteristic neuro-toxin.

Type-species *N. tetani* (*Bacillus tetani* Nicolaier), to be described in a future paper.

The definite designation by Buchanan (1918, p. 38) of *Plectridium tetani* (Nicolaier) Fischer as the type of this genus would seem to reduce the genus *Nicolaerillus* to a synonym of *Plectridium*. However, the author (Heller) insists that this is not the *Plectridium* of Fischer.

Nigrococcus. A name applied by Castellani and Chalmers (1919, p. 925) to a genus of the tribe *Micrococceae*. The description (1919, p. 932) follows:

Micrococceae saprophytic, rarely parasitic, producing black or bluish-black pigment.

Type *Nigrococcus nigrescens* (Castellani, 1911).

Nitribacillus. A generic name suggested by Beijerinck (1914, p. 1163) to replace *Nitrobacter* Winogradsky. Later (1914, p. 110) he states:

In meiner Mitteilung in der Akademie der Wissenschaften zu Amsterdam, Bd 23, pag. 1163, 28 März (10 April) 1914, habe ich das Nitratreferment, wegen seiner Beweglichkeit *Nitribacillus* genannt. Doch scheint es mir gegenwärtig, dass der von Winogradsky eingeführte Name *Nitrobacter* bleiben kann.

Nitrobacter. A name proposed by Winogradsky (1892a, p. 87) (1892b, p. 127) to include those organisms which transform nitrous into nitric acid. No species name was appended, hence from the standpoint of strict nomenclature perhaps the name is invalid as a generic designation. Winogradsky makes the following statement:

S'il fallait donner des noms à tous ces organismes, je proposerais de conserver celui de nitrobactéries à tout le group de microbes, transformant l'ammoniaque en salpêtre. Le ferment nitrifique porterait le nom de *Nitrobacter*.

The organism described is a small rod, about 0.5 μ in diameter, about 2 to 3 times as long as broad, non-motile, embedded in delicate gelatinous masses. It stains poorly. It grows only on media containing nitrites and little or no organic matter. It transforms the nitrites to nitrates.

The organism is termed *Bacterium nitrobacter* by Migula (1900, p 514)

The name is rejected by Erwin F Smith (1905, p 162).

Beijerinck (1914, p 109) in a paper on nitrification states that the organisms present include

Nitrobacter oligotrophum—*Nitrobacter polytrophum*, das Nitratferment selbst
1) Im nitratierenden, meist unbeweglichen Zustand als *N oligotrophum*, im saprophytischen oft beweglichen nicht nitratierenden Zustand, als *N polytrophum*, zu bezeichnen Die beiden Formen müssen als physiologische Arten bezeichnet werden und verhalten sich zu einander als Modifikationen und nicht als Mutationen Der Uebergang findet wie der Pfeil andeutet, nur in eine Richtung statt

Just what nomenclatural recognition should be given these names of Beijerinck is somewhat conjectural Probably *Nitrobacter oligotrophum* may be regarded as the first use of the generic name *Nitrobacter* in a binomial

Winslow *et al* (1907, p 552) recognized the genus with the following description

Cells rod-shaped, non-motile, not growing readily on organic media or in the presence of ammonia Cells capable of securing growth energy by the oxidation of nitrites to nitrates

Winogradsky named no species, although he described one It might be termed *Nitrobacter Winogradskyi* and made the type species

The genus was placed in the family *Nitrobacteriaceae* Substantially the same diagnosis was used by Buchanan (1918, p 180) the Committee (1920, p 202), and Bergey *et al* (1923, p 34)

The genus is here considered valid, and the type species *Nitrobacter oligotrophum* Beijerinck (*Nitrobacter Winogradskyi* Committee)

Nitrobacteraceae. A variant spelling of *Nitrobacteriaceae* *qv* used by Castellani and Chalmers (1919, p 932), by Winslow *et al* (1920, p 201) and by Bergey *et al* (1923, p 31)

Nitrobacteriaceae. A family name proposed by Buchanan (1917, p 349) for the fourth family of the *Eubacteriales* The diagnosis (1918, p 180)

Cells spherical or rod-shaped, motile or non-motile, not growing on ordinary laboratory media in the presence of organic matter Securing growth energy primarily by the oxidation of ammonia to nitrites or of nitrites to nitrates

Three genera were included, *Nitrosomonas*, *Nitrobacter* and *Nitrosococcus*.

Winslow *et al* (Committee Soc Am Bact, 1917, p 551) used the following description, including this family as the first in the *Eubacteriales*

Organisms usually rod-shaped (sometimes spherical in *Nitrosomonas* and possibly in *Azotobacter*) Cells motile or non-motile, when motile with polar, never peritrichous, flagella Endospores never formed Obligate aerobes, capable of securing growth energy by the direct oxidation of carbon, hydrogen or nitrogen or of simple compounds of these Non-parasitic (usually water or earth forms)

The Committee (1920, p 200) divided the family into two tribes, *Nitrobacteriae* and *Azotobacteriae* and revised the diagnosis as follows.

Organisms usually rod-shaped (sometimes nearly spherical in *Nitrosomonas*, and possibly in *Azotobacter*) Cells motile or non-motile Branched involution forms in *Rhizobium* and *Acetobacter* Endospores never formed Obligate aërobies, capable of securing growth energy by the direct oxidation of carbon, hydrogen or nitrogen or of simple compounds of these Non-parasitic (except in genus *Rhizobium*)—usually water or earth forms

Bergey *et al* (1923, p 30) has included this family as the first in the order *Eubacteriales*

Nitrobacterieae. A name used by Castellani and Chalmers (1919, p 932) to designate the first tribe of the family *Bacillaceae* with the description "Growth in ordinary laboratory media entirely or almost entirely absent" *Nitrobacterium qv* is designated as the type genus

Winslow *et al* (Committee Soc Am Bact, 1920, p 201) used this designation (with the spelling *Nitrobacteriae*) for the first tribe of the family *Nitrobacteriaceae*, with the description "Organisms deriving their life energy from oxidation of simple compounds of carbon and nitrogen (or of alcohol)"

Bergey *et al* (1923, p 31) have followed the committee

Nitrobacterium. A generic name proposed by Castellani and Chalmers (1919, p 933) for the species *Nitrobacterium nitrobacter* (Winogradsky) Castellani and Chalmers No description is given It is the single genus of the tribe *Nitrobacterieae qv*

Nitromicrobium. A name proposed by Stutzer and Hartleb (1899, p 197) for *Nitromicrobium germinans*, an organism changing nitrites into nitrates Cells $0.76-1 \times 1-2.5\mu$, ellipsoidal one end thicker than the other Multiplication resembles the budding of yeasts Growth takes place on nitrate agar, and on agar with soil extract and ammonium phosphate, likewise on nitrate agar Organic nitrogen unsuited, and carbon compounds also interfere The authors believe

Winogradsky's *Nitrobacter* to have been an impure culture of *Nitro-microbium* and *Hyphomicrobium*. Believe that this constitutes a distinct type of organism.

Nitromikrobium. A variant spelling of *Nitromicrobium*, *qv* used by Behrens (1899, p. 244) in a review.

Nitromonas. A designation for organisms oxidizing ammonia to nitrites used by Winogradsky (1890, p. 258). He later (1892, p. 127) changed the name to *Nitrosomonas* (*qv*). The name was again proposed by Orla-Jensen (1909, p. 334) as the fourth genus of his family *Oxydobacteriaceae* to include the bacteria which oxidize nitrites to nitrates, a substitute for *Nitrobacter* Winogradsky.

Apparently Winogradsky named no species, and this name can be regarded merely as a casual designation. The name of Orla-Jensen is a synonym of *Nitrobacter*.

Nitrosobacterium. A generic name used by Rullmann (1897, p. 288) for a species *Nitrosobacterium novae formae*. This organism grew well in ordinary culture media as a short rod, in inorganic media as simple or branched filaments. The organism possessed the power of oxidizing nitrites. The organism has apparently not been found by subsequent investigators. It is questionable whether this organism belongs with the true nitrite bacteria.

Hartleb and Stutzer (1897, p. 621) believe the organism described by Rullmann to be merely a special growth stage of their "saltpeter fungus".

Nitrosobakterium. A variant spelling of *Nitrosobacterium* used by Rullmann (1897, p. 228) and other writers.

Nitrosococcus. A name proposed by Winogradsky (1892, p. 127) for the "new world" spherical organisms which oxidize ammonia to nitrites. He says "Les microbes nitreux du nouveau monde feraient le genre *Nitrosococcus*". Apparently no specific name was given. It was later (1900, p. 194) named *Micrococcus nitrosus* by Migula. It is rejected by Erwin F. Smith (1905, p. 162).

It was recognized by Buchanan (1918, p. 180) with the following description and type designation.

Cells spherical. Not growing readily on organic laboratory media. Oxidizes ammonia to nitrites.

Winogradsky termed the single species "les microbes nitreux de nouveau monde" but without a species name. It may be termed *Nitrosococcus americanus* and regarded as the type.

Apparently the correct type designation would be *Nitrosococcus nitrosus* (Migula)

Winslow *et al* (Committee Soc Am Bact (1917, 1920)) merged this genus with *Nitrosomonas*, *q v*

Nitrosomonas. A generic name proposed by Winogradsky (1892, p. 127) to replace the earlier *Nitromonas* He says

Les ferments nitreux du vieux monde ferarent partie du genre *Nitrosomonas* (au lieu de *Nitromonas*, que j'ai propose avant) avec deux especes *Nitrosomonas europaea*, *Nitrosomonas javanensis* (d'autres encore, peut être) et des varietes locales

The organisms oxidize ammonia to nitrites They are rod-shaped forms, being differentiated in this respect from the new world genus *Nitrosococcus*

Migula (1900, p 954) included this genus in his genus *Pseudomonas* The name was rejected by Erwin F Smith (1905, p 162) Orla-Jensen (1909, p 334) includes this as the fifth genus of his family *Oxydobacteriaceae*.

Winslow *et al* (1917, p 552) included *Nitrosococcus* in this genus, and used the following diagnosis

Cells rod-shaped or spherical, motile or non-motile, if motile with polar flagella Capable of securing growth energy by the oxidation of ammonia to nitrites Growth on media containing organic substances scanty or absent

The type species is *Nitrosomonas europaea* Winogradsky

Buchanan (1918, p 180) gave the following generic description

Cells rod-shaped Motile Not growing readily on organic media Oxidizing ammonia to nitrates

The type species is *Nitrosomonas europaea* Winogradsky

Bergey *et al* (1923, p 35) followed the Committee, including this as the fourth genus of the tribe *Nitrobactereae*

Nocardia A generic name proposed by Trevisan (1889, p 9) to create a valid name to replace the invalid terms *Streptothrix* Cohn, *Actinomyces* Harz and *Discomyces* Rivolta It is included as the third genus of the tribe *Cladotrichee* The generic description given is "Cladotrix senza vagine" Five species are listed, *N farcinica* and *N actinomyces* being the first two The generic description given by De Toni and Trevisan (1889, p 927) is as follows

Filamenta tenuissima, evaginata, articulata, Cladotricis more pseudoramosa, nunc e nucleo firmo radialiter expansa, nunc varie coalita Arthrospora in

filamentis normalibus obvenientes, transformatione cocci singuli ortae
Cladothrix sine vaginis

Savageau and Radais (1892, p 242) revived the old name *Oospora* for the group Blanchard (1896, p 853) discusses this proposition in the following paragraph

Sauvageau et Radais rapportent au genre *Oospora* Wallroth 1833, toute une série de microphytes de très petite taille, zu'on range communément parmi les Bactéries tels sont les genres *Streptothrix* Cohn 1874, *Actinomyces* Harz 1877, et *Micromyces* Max Gruber 1891 Ils ont, en effet, une grande ressemblance morphologique avec les *Oospora* vulgaires, mais ils s'en distinguent par leurs dimensions beaucoup plus restreintes Trevisan à créé pour eux, avec raison, croyons nous, le genre *Nocardia* (1889)

This author recognized six species Wright (1904, p 396) suggested the separation of the genus *Nocardia* from *Actinomyces* because of differences in morphology both within the tissues and on culture media He says

I do not accept the prevalent belief, based on the work of Bostroem, Gasperini, and others, that the specific infectious agent of actinomycosis is to be found among certain branching microorganisms, widely disseminated in the outer world, which differ profoundly from *Actinomyces bovis* in having spore-like reproductive elements I think that these should be grouped together as a separate genus with the name of *Nocardia*, and that those cases of undoubted infection by them should be called nocardiosis and not actinomycosis The term actinomycosis should be used only for those inflammatory processes the lesions of which contain the characteristic granules or "drusen" That a *Nocardia* ever forms these characteristic structures in lesions produced by it in man or cattle has not been convincingly shown

He further says (p 385)

To members of this group the generic names *Streptothrix*, *Nocardia*, *Cladothrix*, and *Oospora* already have been given by various writers Of these, in accordance with well-recognized principles of nomenclature, "*Nocardia*" is the only one that is tenable for the group, if it be made into a separate genus, because the other names have been previously used for plant genera to which these microorganisms clearly do not belong, and they are therefore untenable

Against the status of the term "*Nocardia*" as a generic name for these microorganisms, it may be urged that its original definition was faulty, chiefly in that the branching of the filaments was described as false, but this objection has no weight in view of the fact that Blanchard has published an accurate description of a number of species of this group and adopted "*Nocardia*" as the generic name

The name *Nocardia* is rejected by Erwin F Smith (1905, p 162)

Jordan (1908) accepts Wright's conclusions, and separates *Nocardia* with the definition "Filaments with true branches, with reproductive elements or spores observed" from *Actinomyces* in which no such spores are formed

Pinoy (1913, p 929) proposed that *Nocardia* should replace the old name *Actinomyces* and that a new name, *Cohnistreptothrix* be created with Cohn's *Streptothrix Foersteri* as the type

Vuillemin (1913, p 526) concluded that the name *Nocardia* should replace all of these terms, and lists it as a "*genus conservandum*" in his group *Microsiphones*

Winslow *et al* (Committee Soc Am Bact, 1917, p 554) gave the following definition

Branched filaments, resembling a mycelium, readily breaking up into segments, usually saprophytic soil forms Differs primarily from *Actinomyces* in the development of *aerial* hyphae and conidia Usually aerobic Many are pigment formers Colonies as a rule mold-like on culture media

Merrill and Wade (1919, p 64) reject the name in the following statement

This name was adopted by De Toni and Trevisan to cover the entire group Blanchard used it for a time in its original application and Wright (1894) adopted it for nonpathogenic strains only As many other authors used it in one sense or another, of late it has gained much prestige Vuillemin, and Chalmers and Christopherson have recently adopted it for the entire genus

The validity of this name we deny on the grounds indicated in the discussion that follows

Castellani and Chalmers (1919, p 1040) recognize the genus as the type of their family *Nocardaceae*

They give the following definition, designation of type, and reasons for adoption of name

Nocardaceae growing aerobically, usually easy of culture, and producing arthrospores

Type Species *Nocardia bovis* (Harz, 1877)

Nomenclature Bollinger's ray fungus (*Nocardia bovis*) belongs to a genus of which the correct name is *Nocardia* Toni and Trevisan, 1889, a term derived from Nocard, the celebrated French parasitologist, who was the first investigator to clearly recognize this fungus in France We state that it is the correct name for the following reasons

- 1 It is the oldest name, against which no objections can be raised

- 2 It has been formally adopted by the Botanical Section of the First International Congress of Pathology

This genus will be regarded as a synonym of *Actinomyces*, see for reasons.

Nocardiaceae A family (first) of the *Microsiphonales* named by Castellani and Chalmers (1919, p 1040) It is defined as "*Microsiphonales* with a mycelium" It contains two genera *Nocardia* and *Cohnistrepthothrix*

Nodofolium. A generic name proposed by Ellis (1907, p 502) for one of the iron bacteria The species named is *Nodofolium ferrugineum* It is defined as a flat band with regular constrictions Loops and arches developed Reproduction is by the formation of conidia on the outside of the filament

Molisch regards this as a synonym of *Leptothrix*

Nosema. A generic name proposed by Naegeli (1857, p 760) for the species *Nosema bombycis*, which was at first taken for a bacterium Later it was shown to be a protozoan It is therefore a protozoan genus.

Novillus. The name of the fifteenth genus of the *Clostridioideae* given by Heller (1922, p 22) with the following description

Clostridioideae that in blood broth produce H_2S demonstrable by lead-acetate-paper test, they liquefy gelatin Produce gas and on continued incubation produce a pink coloration in meat, but this color rapidly fades Autoagglutinate with extreme readiness Slowly attack milk and a few carbohydrates, but do not form much acid Heavy rods of apparently shy growth habit on most media, form a few oval spores that may or may not bulge the sides of the bacilli Colonies usually large and opaque, may form slender projections or even long fine woolly filaments Frequently yellowish Typically toxic tissue invaders which frequently cause formation of a thick gelatinous oedema that does not lose its gelatinous consistency on section May also produce gas and a black haemorrhagic condition of the muscle if rapid invasion by large numbers of bacilli takes place One strain causes considerable tissue destruction Pathogenic for guinea-pigs, mice, man, hogs, horses, cattle

Type species *N. maligni* (*Bacillus oedematis-maligni* II, Novy) as described by Novy

Octopsis A genus of bacteria proposed by Trevisan (1885, p 102) to include *Octopsis cholerae-gallinarum* of fowl cholera, *O. equorum* of horse typhoid, *O. xanthogenicum* of yellow fever, and *O. lyssae* of rabies The name has not been used by subsequent authors and was abandoned by Trevisan himself However, a strict application of the rules of priority would make this a valid name and reduce to synonymy the later name *Pasteurella* It is possible that it might be rejected as invalid upon the basis of an insufficient description, as being "universally regarded as non-valid"

It is rejected by Erwin F Smith (1905, p 162)

Oenobacillus. A generic name used by Forti (1896) for an organism described by him as *Oenobacillus Abbae*. It was a polymorphic bacillus isolated from red wine. The original description is not available. The name has been used by Kossowicz (1911, p 113)

In young cultures it appears as a diplococcus, later as rods, rounded at the ends and in pairs. Gelatin is not liquefied. Produces undesirable changes in wine. Non-motile. Grows better in oxygen.

Omelianskillus. A name proposed by Heller (1922, p 5) as the second genus of the *Clostridiaceae*, with the description

Clostridiaceae that do not grow well in media containing much protein, and may derive all their nitrogen from mineral salts. Split cellulose or hemicellulose. Do not contain starch. Long slender bacilli with spherical spores. Colonies may be produced on potato slants, they are minute, yellowish and transparent. Agents of plant putrefaction, found everywhere.

Type species *O. hydrogenicus* (the ferment of cellulose which produces hydrogen, of Omeliansky) as described by Omeliansky (1895 and 1904b). In his later work he admits that the cultures studied by him were not pure. Another type, *O. methanicus*, similar to the above, is said to be the agent of methane formation in cellulose fermentation. This species may be assigned to the same genus.

Oospora. A genus created by Wallroth (1831, p 182) for a group of eleven species of fungi with the following description: "Sporidia subglobosae oviformia intricata pellucida, primum concatenata, hypharum articulatarum simplicem teneram decumbentem mentientia, articulisque inter se facile secedentibus fragilia."

The genus has in general been recognized by mycologists. It is of interest in bacteriology because in 1892 Sauvageau and Radias (1892, p 246) concluded that the *Streptothrix* of Cohn and the *Actinomyces* of Harz belonged in this genus.

In this conception they were followed by Lehmann and Neumann (1896, p 375) and by Chester (1897, p 37). Berestnew (1899, p 390) concluded that the genera were distinct, and in this was followed by Lehmann and Neumann in later editions, and by Chester (1901). It is rejected by Musgrave and Clegg (1908, p 458).

It is probable that the trichomycetes should not be given the name *Oospora*. It is a valid mold genus, but an invalid bacterial genus.

Ophidobacteries. A name used by Maggi (1886, p 176) for his fifth order of bacteria. He says "Bacteries a corps filiforme, ondule ou flexueux a mouvements serpentiformes." The genera included are *Vibrio* and *Tyrophthrix*.

Ophidomonas. A genus created by Ehrenberg (1838, p 43) for one of his cryptomonads. The illustration (1838, fig 11) (Migula 1897, p 5) shows an olive green or brownish colored individual. It is probable that the organism was one of the red sulphur bacteria, for Ehrenberg named one of his organisms *Ophidomonas sanguinea*. Ehrenberg differentiated from *Spirillum* in that the latter frequently showed incomplete cell divisions, the former always divided completely.

The first species named, *Ophidomonas jenensis* Ehrenberg (1838, p 43) is identified by Cohn (1872, p 183) with his *Spirillum volutans*. Cohn concludes that Ehrenberg's basis of differentiation of *Ophidomonas* and *Spirillum* was inadequate. Cohn later (1875, p 170) noted an observation of *Ophidomonas sanguinea*. He says

Seit dem wir bei dem alten *Spirillum volutans* Flummergeisseln aufgefunden, besteckt zwischen *Spirillum* und *Ophidomonas* überhaupt kein Unterschied, vorausgesetzt, dass auch bei den kleineren Spirillen Bewegungsorgane noch nachtraglich erkannt werden sollten. Wir haben daher nur die Wahl, entweder *Ophidomonas* als selbständige Gattung zu streichen, und unsere Art etwa unter dem Namen *Spirillum sanguineum* gesissemassen als das Mummuth unter die Bacterien enzureihen oder umgekehrt die mit Flummergeisseln nachweislich ausgestatteten Schraubenfaden (*volutans*, *jenensis* und *sanguinea*), unter *Ophidomonas* zusammenzufassen, und den Namen *Spirillum* ausschliesslich für die kleineren Species (*tenue*, *Undula*) so lange beizubehalten als an ihnen noch keine Geisseln entdeckt sind. Sollte dies geschehen, so wurde umgekehrt der Name *Spirillum* zu löschen sein.

The name *Ophidomonas* was used by Hansgirg (1888, p 264) for the second section of the genus *Spirillum*. He says "*Ophidomonas* (Ehrb.) Hansg. Zellen rosen- bis blutroth, violett oder rothlichbraun, meist mit Zahlreichen, stark lichtbrechenden Schwefelkörnchen in plasmatischen Inhalte." One species is included, *Spirillum sanguineum* (Ehrb.) Cohn.

Winogradsky (1888) created the genus *Thiospirillum*, with *Thiospirillum* (*Ophidomonas*) *sanguineum* as the type.

De Toni and Trevisan (1889, p 1006) regard *Ophidomonas* as a synonym of *Spirillum*.

Migula (1895) followed Hansgirg in making *Ophidomonas* a subgenus of *Spirillum*, defining it to include those forms having sulphur granules.

Later, however (1900, p 1049), he accepted the generic designation *Thiospirillum* of Winogradsky for the red sulphur bacteria with spiral cells, making *Ophidomonas* a synonym, with the type *Thiospirillum* (*Ophidomonas*) *sanguineum*.

This name was placed in a list of rejected genera by Erwin F. Smith (1895, p. 162)

Apparently Winogradsky was in error in creating the genus *Thiospirillum* when the valid name *Ophidomonas* was in existence. It would seem that the latter name is a valid generic designation of the red sulphur bacteria with spiral cells. The type species is *Ophidomonas jenensis* Ehrenberg.

Ophryothrix. A generic name given by Borzi (1878, p. 274) to an organism which he named *Ophryothrix Thuretiana* and regarded as one of the algae. This is made as synonym of *Leptothrix Thuretiana* by Hansgirk and of *Leptotrichia Thuretiana* by Trevisan (1889).

It is probably invalid as a bacterial generic name, although its status is not entirely clear.

Oscillospira. A generic name proposed by Chatton and Pérard (1913, p. 1160) for the bacterial species *Oscillospira Guillemondi* observed in the coecal contents of the cavy. The description is as follows:

Oscillospira Guillemondi se présente sous forme de filaments robustes, assez bien calibrés, d'un diamètre moyen de 5μ et dont la longueur n'excède jamais 100μ . Ils sont arrondis aux deux extrémités. Examinés vivants ils montrent un cloisonnement bien apparent, les cloisons sont très rapprochées, et les compartiments qu'elles limitent, ceux où se forment les spores excepté, ont de 1 à 2μ de longueur. Ils sont donc extrêmement étroits. L'épaisseur des cloisons est d'ailleurs variable. Les plus épaisses sont biconcaves. Le cytoplasme est homogène ou finement granuleux sans pigment ni grains de soufre, ni inclusions d'aucune sorte. Ces filaments se multiplient par simple scission transversale entre deux compartiments, généralement de part et d'autre des grosses cloisons biconcaves qui équivalent aux disjoncteurs des autres Oscillariées. Les filaments sporulés ne sont jamais nombreux. Chacun d'eux forme le plus souvent une, rarement deux, endospores. La spore est contenue dans un des compartiments considérablement allongé, mais très peu élargi, qu'elle ne remplit d'ailleurs pas entièrement. Elle est ellipsoïdale, mesure en moyenne 4μ de long sur 2.5μ de large. Elle est orientée suivant l'axe du filament ou très peu inclinée sur cet axe. Après coloration, le contenu des compartiments est sans différenciation chromatique centrale. Par contre on y distingue une ou deux vacuoles. *Oscillospira* est très mobile. Les filaments se déplacent suivant une ligne hélicoïdale à tours très lâches, telle que les filaments les plus longs n'ont jamais plus d'un tour et demi de spire. Nous n'avons pas coloré les cils. La mobilité n'est d'ailleurs pas constante. Chez certains cobayes tous les individus se montrent immobiles. La structure de cet organisme en fait incontestablement une cyanophycée. L'absence de pigment et de corps chromatique central la rapproche des *Beggiatoa*. Mais dans les conditions où nous rencontrons *Oscillospira* elle n'élabore pas de soufre. C'est là d'ailleurs chez les *Beggiatoa* même un caractère purement physiologique, subordonné aux conditions de milieu.

The organism was found quite constantly present in the intestinal contents of cavies at the Pasteur Institute in Paris

Oxydobacteriaceae The first family of the order *Cephalotrichinae* named by Orla-Jensen (1909, p 329) It includes those obligate aerobic bacteria which oxidize carbon, nitrogen or hydrogen compounds without the production of noteworthy amounts of unoxidized split products None of the organisms produce thread-like filaments or produce a mycelial growth

Pacinia. A genus of bacteria named by Trevisan (1885, p 83) to include the organism which causes Asiatic cholera. The original generic description is

Tre stadi di sviluppo vegetativo 1 Bacilli, 2 Filamento, 3 Cocchi Bacilli (stadio protoplastico tipico) cilindrici, più o meno ricurvi, inarticolati, incolori, biforini, lunghi (macrobacilli) e corti dio transitorio irregolarmente flessuoso-ondulato-angolosi risultanti da più bacilli runasti in conguunzione per incompleta divisione trasversale Cocchi (stadio finale) derivati da microbacilli, dapprima in brevi monifi, indi liberi Spore provenienti tauto nei microbacilli quanto nei cocchi

The first species named is *Pacinia cholerae asiaticae* As synonyms he notes *Vibrio cholerae* Pacini (1854) and *Bacillus cholerae asiaticae* Trevisan Other species described are *Pacinia Finkleri* of cholera mostras, *P Lewisii* from saliva, and *P Cecii* from cholera stools Later (1889, p 23) this genus was included as the tenth of the tribe *Bacillae*, with the description "Baculi cilindrici, non mai spirali, con protoplasma uniformemente diffuso Spore macrosome provenienti in una estremita intumescete a guisa di capocchia, di rado in entrambe le estremita, di baculi normali immutati" Sixteen species were listed

De Toni and Trevisan (1889, p 1015) give as synonyms *Dispora* Kern and *Microspira* Schroeter The generic diagnosis given is

Baculi cylindrici, nunquam spiraliter torti, plasmate uniformiter diffuso foeti Sporae (arthrosporae) macrosomae, ex uno apice (raro utroque) capituli ad instar intumescetes, in baculis normalibus immutatis ortae

The genus is divided into two sections, *Eu-Pacinia* with 11 species and *Pseudospira* with 8 species Among the species of the first section are *P putrifica*, *P Loeffleri* (diphtheria bacillus) and *P caucasica* The second section includes *Pacinia cholerae-asiaticae* as its type It corresponds quite exactly to the *Microspira* of Schroeter and Migula

The genus *Pacinia* has not been recognized by subsequent authors. It is rejected by Smith (1905, p 162) although he states that were it not for the availability of the generic designation *Vibrio* for the genus of which the comma bacillus is the type, *Pacinia* would have priority over *Microspira* Schroeter

The genus is likewise rejected by Vuillemin (1913, p 521) as not satisfactorily characterized *Pacinia* may probably be regarded as a synonym of *Vibrio*

Pacinieae. The fifth subtribe of the tribe *Bacilleae* used by De Toni and Trevisan (1889, p 1015) with the following diagnosis "*Baculi nudi nunquam spiraliter torti Arthrosporae*" The following genera are included *Pacinia* and *Bacterium*

Palmella. A genus of algae named by Lyngbye Various bacteria have been incorrectly placed here Among them are *Palmella murifica* Rabenhorst = *Micrococcus murificus* Trevisan, *Palmella prodigiosa* Montagne = *Micrococcus prodigosus* Cohn

Panhistophyton. According to Zopf (1885, p 69) the name *Panhistophyton ovatum* was given by Lebert (1856) to the organism earlier grouped with the bacteria and now known to be a protozoon, *Nosema bombycis* Nageli Trevisan (1882, p 100) says

La prima opinione fu emessa dal Lebert, che li ritenne un alga unicellulare, cui diede il nome *Panhistophyton ovatum* (In Berlin Entomolog Zeitschr, 1858), ma il Cohn avendodimostrato che erano Batterii appartenenti al genere *Micrococcus*, rimaseso nelle tabelle sistematistiche coll 'apellativo *Micrococcus bombycis*

Panhistophyton is not a bacterial genus

Parachromatium. An alternative to *Azotobacter* proposed as a generic name by Beijerinck (1903, p 217) In a footnote he says

Peut être le nom de *Parachromatium* qui indique la parenté de notre microbe avec le genre *Chromatium* de Winogradsky, serait préférable Des considérations physiologiques m'avantent d'abord conduit à une tout autre opinion mais des études ultérieure me portent a croire que cette parenté generique est indubitable Zettnow, en examinant mes preparation, avait deja la meme opinion

The name does not seem to have been used, and may be regarded as invalid as a generic designation

Paracloster. A generic name used by Fischer (1895, p 141) to designate those non-motile rod-shaped bacteria in which the cells become spindle shaped when sporulating The description given is

Unbeweglich, ohne Geisseln, mit Endosporen in spindelförmig aufgeschwollenen Stäbchen, die Spore liegt oft genau in der Mitte, kann aber auch in

einem der Enden, also ausserhalb der grossten Schwellung liegen Bis jetzt ist kein Vertreter dieser Gattung bekannt

Paracloster butyracus, the *Granulobacter immobilis* of Grassberger and Schattenfroh was later given as the type (1903, p 60)

This generic name has apparently not been used by other authors It is rejected by Erwin F Smith (1905, p 162) It may be regarded as a synonym of *Clostridium*

Paracoccaceae. A subfamily of the *Coccaceae* proposed by Winslow and Rogers (1905, p 660) with the diagnosis.

Parasites (thriving only, or best, on, or in the animal body), thrive well under anaerobic conditions Many forms fail to grow on artificial media, none produce abundant surface growths Planes of fission generally parallel, producing pairs, or short or long chains

This diagnosis was emended in 1906, (p 545), by the addition of "never packets Generally stain by Gram Produce acid in dextrose and lactose broth Pigment, if any, white or orange" In their last contribution (1908, p 240) the Winslows include the following genera *Diplococcus*, *Ascococcus*, *Streptococcus*, *Albococcus* and *Aurococcus*

It would seem that the subfamily name *Paracoccaceae* is invalid for two reasons, it has the ending prescribed by the botanical code for families and not subfamilies, furthermore, it is not formed as prescribed in the code by the addition of an appropriate ending to the name of one of the genera included Inasmuch as no tribes are included in the Winslow's classification, the tribal name *Streptococceae* used by Trevisan might well be used in place of this subfamily

Parameningococcus. A casual name used by Dopter (1909, p 74) for a type of meningococcus It is differentiated primarily by its agglutination relationships

Paraplectrum. A generic name used by Fischer (1895, p 141) for non-motile rod-shaped bacteria which produce spores in the cell tips, with consequent enlargement and formation of "drumsticks" The description given is "Unbeweglich, ohne Geisseln, mit Endosporen in einem kopfig aufgeschwollenen Ende" The type given is *Paraplectrum Peroniella*, the *Bacillus Peroniella* of Klein (1889, p 65) The species mentioned later (1903, p 61) is *Paraplectrum foetidum* Weigmann

This genus has not been used by other authors. It is rejected by Erwin F Smith (1905, p 162) It may be regarded as a synonym of *Plectridium*.

Paraschizae. A subdivision of the group *Schizomycetes* named by Jennings (1899, p 315) to include the genera *Astrobacter* and *Pasteurella* characterized by longitudinal rather than transverse fission of the cell in multiplication. The coordinate subdivision is named *Diaschizae*.

Paraspirillum. A genus used by Dobell (1911, p 97) to include the species *Paraspirillum vejdoskii* found in a culture of *Oscillatoria*.

The living organisms are, for the most part, in the form of spiral or S-shaped rods, very like ordinary spirilla. Short individuals are about 8μ in length, the largest individuals attaining a length of about 25μ . The average length is about 15μ .

The organisms are variable in thickness, with a well marked thickening toward the middle of the body. They taper decidedly toward the ends. The cell is described as a "much elongated and spirally twisted spindle." "In the middle region, the organisms measure from 1.5μ to 2μ across—the thickness varying in different individuals."

In each living cell at thickest point is an oval or rounded structure regarded as a nucleus. Immediately surrounding the nucleus is an area of hyaline cytoplasm. Toward the ends of the cells are granular inclusions (volutin or metachromatic granules).

"Movement from place to place is effected by means of flagella, similar to those of common spirilla. In the living organisms, I have never succeeded in seeing any flagella." The flagella were demonstrated, however, in stained mounts, a single one, or one at each end.

The genus was recognized as the third of the family *Spirillaceae* by Buchanan (1918, p 179) with the description

Cells spiral or S-shaped, like spirilla, variable in thickness, with a well marked thickening toward middle of the body, and tapering toward the ends, a much elongated and spirally twisted spindle. Motile by means of flagella.

The type species is *Paraspirillum vejdoskii* Dobell.

Pasteurella. A genus created by Trevisan (1887, p 94) primarily to include the bacillus of fowl cholera. His description follows:

Pasteurella, Trevis (nuovo genere). Tre stadi di sviluppo vegetativo: 1 Baculi, 2 Filamenti, 3 Cocchi. Baculi (stadio primigeno tipico) brevemente cilindrici o ellissoidi, inarticolati incolori, protoplasma polare-diblastico. Filamenti (stadio anormale transitorio) visulanti da baculi rimasti in congiunzione per incompleta temnogenesi. Cocchi segregati (monococchi e cocchi ottoformi) o serati in catenelle. Spore provenienti nei baculi.

The species first named is *Pasteurella cholerae-gallinarum*. Other species named were *P. Davainei* and *P. suilla*. Later (1889, p 21)

twelve species were included. The genus as discussed in Saccardo's Sylloge contains eighteen species. The genus is characterized as follows: "Baculi plasmate polari-diblastice foeti Sporae (arthrospora-) isosomae, microsomae." In the analytical key the following additional characters are found, "Baculi recti vel incurvi, nunquam spiralter totti." "Baculi in coenobium reticulatum non connexi." "Baculorum divisio semper in unam eandemque directionem longitudinalem repetita."

The name is rejected by Erwin F. Smith (1905, p. 162).

The genus *Pasteurella* has been recognized by various authors particularly Lignieres (1900, p. 212) who worked with the hemorrhagic septicemia group of organisms, as the cause of the so-called "Pasteurelloses," but has not found its way into formal classification. Lignieres (1900, p. 331) gives the following diagnosis of *Pasteurella*:

Parmi les maladies qui forment le groupe des septicémies hémorragiques, on en trouve un certain nombre dont les microbes spécifiques répondent aux caractères invariables suivants: Cocco-bacilles, sans mouvements de translation, ne prenant pas le Gram, très polymorphes, donnant des formes d'involution, ne liquéfiant pas la gélatine, ne coagulant pas le lait dont la réaction reste normale, ne donnant pas de culture visible sur pomme de terre naturelle acide, ni d'indol dans le bouillon pancréatique, ne rougissant pas le gelose de Wurtz, surtout aérobies, mais aussi anaérobies, présentant une odeur sui generis dans les cultures. Pas de spores, pas de cils. Virulence très variable, en général grande. Par injection intra-veineuse, affinité spéciale pour les synoviales tendineuses et articulaires. Les caractères que je viens de mentionner sont fixes, absolus, l'absence de l'un ou de l'autre exclut le microbe du groupe des *Pasteurella*. Comme indication générale, on peut ajouter que ces microbes se colorent facilement aux deux extrémités en laissant le centre clair, qu'ils ne se colorent pas aussi aisément que d'autres microbes, le coli, par exemple, qu'ils poussent parfois discrètement sur les milieux de culture habituels, que dans certains cas enfin ils sont très difficiles à mettre en évidence au sein de l'organisme.

Vuillemin (1913, p. 521) accepts the view of Fischer that the bipolar staining characteristic of this group is an artefact, but believes that the organisms related to fowl cholera constitute a homogeneous group of pathogenic bacteria. However, he advocated abandoning the term *Pasteurella* to the bacteriologists who have vulgarized it into a purely nominal genus.

Kitt in Kolle and Wassermann (6. 37) improperly credits the name to Lignieres, and suggests its abandonment in the following statement:

Von Lignieres ist für die Septicaemia avicum und überhaupt die Septic haemorrhagica der Name *Pasteurellose* und für die Bakterien der Name *Pasteurella*.

einzuführen Einige französische Autoren haben daraufhin diese Nominierung gewählt, während Montfallet & Boschetti mit begründeter Darlegung Stellung dagegen nahmen Auch Referentem scheint die Neubenennung unnötig und willkürlich, und der Name, welcher den Charakter der Krankheit ausdrückt, passender

Winslow *et al.* (Committee Soc Am Bact, 1917, p 561) accepted the designation with the description

Short rods, single or rarely in chains, usually showing distinct polar staining Non-motile Gram-negative Without spores Aerobic and facultative Powers of carbohydrate fermentation slight, no gas produced Gelatin not liquefied Parasitic, frequently pathogenic, producing plague in man and hemorrhagic septicemia in the lower animals

The type species is *Pasteurella cholerae-gallinarum* (Flügge) Trevisan

Buchanan (1918, p 52) used the following description

Short rods, single or rarely in chains, usually showing distinct polar staining, non-motile. Gram-negative, without spores, aerobic and facultative, usually not producing gas, powers of fermentation slight, often pathogenic, not acid-fast, not liquefying gelatin

The type species is *Pasteurella cholerae-gallinarum* (Flügge) Trevisan

The genus is made the type of the tribe *Pasteurelleae* by Castellani and Chalmers (1919, p 943) Bergey (1923, p 264) also includes the genus in the tribe *Pasteurelleae*.

The name of the type species is somewhat puzzling Apparently the first species name for the fowl cholera organism was *Micrococcus gallicidus* Burrill (1883, p 320) His description is inadequate, but he claimed he was describing the cause of fowl cholera Zopf (1885, p 87) called the organism *Micrococcus cholerae-gallinarum* and Trevisan *Pasteurella cholerae-gallinarum* It would appear the correct designation would either be *Pasteurella cholerae-gallinarum* (Zopf) Trevisan or *Pasteurella gallicida* (Burrill) comb nov.

Pasteurelleae. A name proposed by Castellani and Chalmers (1919, p 932) for the eighth tribe of the family *Bacillaceae*. The description (1919, p 943) is.

Bacillaceae with good growth on ordinary media, without endospores, aerobic, without fluorescence or pigment formation, unable to liquefy gelatine, and Gram-negative, but with polar staining

Type genus *Pasteurella* De Toni and Trevisan, 1889

Winslow *et al.* (Committee Soc. Am Bact, 1920, p 211) independently proposed the same name with the following diagnosis "Gram-

negative rods, showing bipolar staining Parasitic forms of slight fermentative power "

The only genus included is *Pasteurella*

Pasteuria. A generic name proposed by Metschnikoff (1888, p 166) for *Pasteuria ramosa*, an organism described from the bodies of the water fleas *Daphnia pulex* and *D magna* The generic diagnosis given by De Toni and Trevisan (1889, p 941) follows

Baculi primitus in cumulos rotundatos, axi centrali ramoso instructos, arctissime acervati, confluentes, dein ob pluries repetitam divisionem longitudinalem segregati, demum soluti, ellipsodei, stipitulati Sporae (Endosporae) globosae, microsome, ad apicem baculorum liberorum obvenientes

Obs A caeteris Baculogenis ob formam valde peculiarem cumulorum primigeniorum et baculorum divisionem semper longitudinalem praedistincta

The organism apparently has not been observed by other investigators Lehmann and Neumann (1896, p 20) say that the organism can scarcely be regarded as belonging to the bacteria in a strict sense

Migula (1904, p 128) doubts the bacterial nature of this organism and excludes it from his classification The name is included by Erwin F Smith (1905, p 162) among the rejected genera

Vuillemin (1913, p 527) states that this organism may be differentiated from *Nocardia* (!) by its claviform ramifications, by its dichotomy and by the formation of spores inside the cells He proposes it as a *genus conservandum* of the *Microsiphoneae* with the type species *P ramosa* Metchnikoff

The nomenclatural status and position of this genus is uncertain The type species is *Pasteuria ramosa* Metchnikoff

Pasteuriaceae A family name proposed by Metschnikoff (1888, p 165) to include his genus *Pasteuria* and the bacteria of the root nodules of leguminous plants

Pasteurieae. A subtribe of the tribe *Bacilleae* named by De Toni and Trevisan (1889, p 941) with the following diagnosis "Baculorum divisio semper in unam eandemque directionem longitudinalem repetita Endosporae " The single genus *Pasteuria* is included

Pectinobacter. A generic name proposed by Makrinov (1916, p 440) for the species (type) *Pectinobacter amylophilum* Enlows (1920, p 69) summarized the description of the species as follows.

Rods 4 to 6 μ long by 0.5 to 1 μ wide Motile Spores are formed, prior to which a fusiform aspect is assumed Gram positive Grows better on starch media than on any other Active fermentative agent Isolated from soil

Pectobacillus. The second genus of the family *Butyribacteriaceae* proposed by Orla-Jensen (1909, p 342) It is defined to include those forms of his family *Butyribacteriaceae* which attack pectins They stain blue with iodine and ferment many carbohydrates An organic source of nitrogen is required, the species are anaerobic The *Granulobacter pectinovorum* of Beijerinck is the type

This name may be regarded as a synonym of *Clostridium* and *Granulobacter* unless the phenomenon of pectose fermentation should prove of generic significance

Pediococcus. According to Lindner (1888, p 97) a species of bacterium was named *Pediococcus cerevisiae* by Balcke (1884, p 183) This was an organism previously noted as occurring in diplococcus and tetrad forms in beer by Pasteur Hansen (1882) termed the organism a *Sarcina* According to Lindner the *Pediococcus cerevisiae* produces colorless colonies on gelatin, which later become yellowish or yellowish brown The gelatin is not liquefied On meat infusion gelatin it grows in stroke culture as a gray white, moist iridescent streak with relatively smooth edges

Lindner (1888) described two additional species of *Pediococcus*, *P. acidilactici* and *P. albus* Trevisan (1889, p 28) included this as the third genus of *Sarcinae* It is ascribed to Lindner (1887) The generic description is "Cocchi aggregati a 4 a 4 in famiglie unistrate nella ganga gelatinosa equalbilmente spianata a guisa di membranella parallelogrammica Temnogenesisi in due direzioni" Ten species were included

This generic name was accepted by De Toni and Trevisan (1889, p 1050) with the following description

Cocci globosi, v ovoides, quaternati, in exiguas familias regulares, unistrates muco amorpho tenuisculo, vix conspicuo, ceto dilabente obvolutas, laxe consociati Coccorum divisio in duas directiones

Three additional species are here listed These authors credit the generic name *Pediococcus* to Lindner and not to Balcke Eisenberg (1891) lists the three species recognized by Lindner under "*Pediococcus*" Lehmann and Neumann (1896, p 102) include this genus as a synonym of *Micrococcus*

The genus is recognized by Fischer (1897) with the following description "Two planes of fission, alternate and at right angles" "*Micrococcus tetragenus*" is given as an example, also the gonococcus This genus is again listed in the second edition (1903, p 60)

Sollhed (1903, p 491) described another species *P hennebergi* from beer

Klocker (1903, p 331) ascribes the genus to Francke and states "The cells are arranged in flat colonies, they divide along two planes"

Erwin F Smith (1905, p 162) placed this name among the rejected genera Vuillemin (1913, p 520) dismisses *Pediococcus* as an invalid synonym of *Merista* Van Tieghem (1883)

If the cocci which are arranged in tetrads are to receive generic recognition, the name *Pediococcus* would appear to be valid If not, this becomes a synonym, perhaps of *Micrococcus* The type species is *Pediococcus cerevisiae* Balcke

Pediokokkus. A variant of *Pediococcus* (*qv*) used by Eisenberg (1891) and other German writers

Pedioplana. A generic name proposed by Wolff (1907, p 9) for the species *P haeckeli* The genus corresponds to *Pediococcus* except that the cells are motile by means of flagella The name has apparently not been used by other investigators It may be regarded as a synonym of *Pediococcus*

Pelochromatium. A generic name proposed by Lauterborn (1913, p 99) for an organism morphologically resembling *Chlorochromatium* but containing bacteriopurpurin Included in his family *Rhodobacteriaceae* The only (type) species is *Pelochromatium roseum* Lauterborn

Pelodictyon. A generic name proposed by Lauterborn (1913, p. 98) for a genus of his family *Chlorobacteriaceae* containing organisms with elongate cells 2 to 3 μ long, yellowish-green in color, and united into a net in a fashion similar to *Thiodictyon* Winogradsky The single species (type) named was *Pelodictyon clathratiforme* (*Aphanothece clathratiformis*) (Szafer) Lauterborn The inclusion of this genus among the bacteria rather than the blue-green algae is questionable

Peloglea A generic name proposed by Lauterborn (1913, p 99) for a genus of his family *Chlorobacteriaceae* with the following description

Peloglea chlorina nov gen nov spec Zellen 0,003 bis 0,004 mm lang, kettenartig zu (fast *Nostocartigen*) locker gewundenen Faden zusammenschliessend und in eine gemeinsame Gallertmasse eingebettet Kolonien bis 1 mm im Durchmesser

The inclusion of this form among the bacteria is questionable

Peloploca. A name proposed by Lauterborn (1913, p 99) for a genus of bacteria belonging to the colorless forms The description is

Peloploca nov gen Fadenformige Zellreihen, in wechselnder Zahl parallel zu starren Bündeln oder Bandern vereint Zellen mit Pseudovakuolen Bewegungslos

P undulata nov spec Zellfaden alle sehr locker spiralg gedreht, zu einem scheinbar wellenformig gebogenen, parallel gestreiften Fadenbündel dicht zusammenschliessend

P taenrata nov spec Zellfaden zu ziemlich breiten, oft gedrehten Bandern vereint, die durch die Pseudovakuolen der Einzelzellen wie gegittert erscheinen Zellen 0,003-0,004 mm lang, Bänder bis über 0.7 mm lang

Peloploca ist im faulenden Schlamm von Characeen-Teichen nicht selten, scheint aber wegen einer gewissen Ähnlichkeit mit mazerierten pflanzlichen Gefässbündeln bisher übersehen worden zu sein

Pelosigma. A generic name proposed by Lauterborn (1913, p 100) to include *Pelosigma cohnii* This organism as *Spiromonas cohnii* the author states was erroneously placed in the flagellate genus *Spiromonas* by Warming

Pelosphaera. A generic name given by Lauterborn (1906, p 197) in the following description

Pelosphaera rotans nov gen nov spec Zellen keilförmig, vorn verbreitert und abgerundet, mit ziemlich derber Membran und meist körnigem Inhalt, in grosserer Zahl radial zu maubbeerförmigen kugeligen bis elliptischen Kolonien vereinigt, die sehr an gewisse Flagellaten-Kolonien erinnern Frei beweglich mit Hilfe von Geisseln

Jüngere Kolonien farblos durchsichtig, kugelig bis elliptisch, Zellen sehr dicht zusammenschliessend, im Centrum der Kugel oft einen Hohlraum freilassend, sehr lebhaft beweglich Aeltere Kolonien meist schwach gelbbraunlich gefärbt, oft unregelmässig im Umriss, Zellverband lockerer, die freien Zellenden mehr oder weniger halbkugelig über die Oberfläche vorspringend Einzelne Zellen bisweilen vergrössert, im Innern mit ansehnlichem, stark lichtbrechendem, kugeligem Gebilde (Spore?) Durchmesser der Kolonie 0,015-0,040 mm

Vermehrung Bisher nur Teilung der Kolonie beobachtet die Kolonie streckt sich und schnürt sich einseitig durch

Peptonococcus. A generic name for peptonizing lactic acid cocci proposed by Orla-Jensen (1909, p 340) in the following statement

Da die peptonisierenden Milchsäurekokken reichliche Mengen Pepton, aber sehr wenig weitere Eiweisszersetzungserzeugnisse bilden, während die von den Proteus-Bakterien abgeleiteten Kokken durch Indolproduktion charakterisiert sein müssen, so liesse sich vielleicht zwischen den Gattungen *Peptonococcus* und *Indolococcus* unterscheiden

Orla-Jensen himself does not include this genus in his formal classification. It is probably invalid as no species is designated.

Peritricha. The fourth group of the *Gymnobacteria* named by Messea (1891, p. 232) to contain bacteria with flagella on all sides.

Peritrichinales. Orla-Jensen (1921, p. 270) suggests splitting *Eubacteriales* (Buchanan) into *Pseudomonadales* and *Peritrichinales*. This latter is the equivalent of his previously used *Peritrichinae* (q.v.). He includes the following genera: 1 *Thermobacterium*, 2 *Streptobacterium*, 3 *Streptococcus*, 4 *Betabacterium*, 5 *Betacoccus*, 6 *Propionibacterium*, 7 *Microbacterium*, 8 *Tetracoccus*, 9 *Coccus*, 10 *Bacterium*, 11 *Bacillus*, 12 *Clostridium*.

Peritrichinae. An order named by Orla-Jensen (1909, p. 328) to include those bacteria with spherical or rod-shaped, never spiral, cells which, when motile, have peritrichous flagella. The group as a whole is not typically made up of water bacteria. The organisms of this group do not secure their growth energy solely by oxidative processes. The other of the two orders into which the *Schizomycetes* is divided is *Cephalotrichinae* (q.v.).

The order *Peritrichinae* includes the following families: *Acidobacteriaceae*, *Alkalibacteriaceae*, *Butyribacteriaceae* and *Putribacteriaceae*.

Perroncitoa. This genus of cocci was named by Trevisan (1889, p. 29), with the description "Cocchi compressi seriati lateralmente a due a due (diplococchi transversali) in filamenti semplici rivestiti di vagine cilindriche membranaceo-gelatinose. Spore macrosome provenienti nei filamenti. La forma compressa dei cocci, appaiata a guisa delle Neisserie, é affatto speciale. Quando si liberano dalla vagina divengono perfettamente globulosi." The single species listed is *Perroncitoa scarlatina* (*Micrococcus scarlatinus*, Trevisan). The diagnosis is given as follows by De Toni and Trevisan (1889, p. 1053):

Cocchi compressi, lateraliter duo per duos seriati (diplococchi transversales) in filamentis simplicibus, vaginis cylindricis membranaceo-gelatinosis obducta concatenati. Arthrospora macrosomae in filamentis obvenientes. Cocci e vagina liberati exakte globulosi fiunt.

One species is included, *Perroncitoa scarlatina* Trevisan, the *Streptococcus rubiginosus* Edington (1887, p. 1262).

This genus is undoubtedly founded upon a misconception of morphology. It has never been used by other authors. It is rejected by Erwin F. Smith (1905, p. 162). It may be regarded as invalid, or at least based upon an unidentified organism.

Petalobacillus. A name used by Maggi (1886, p 174) to designate a growth form of a *Bacillus*

Petalobacteria A name applied by Billroth (1874, p 16) to a growth form of his pleomorphic *Coccobacteria septica* in which the organism assumed a rod shape and formed a flat gelatinous layer of cells The term has apparently never been used as a generic name

Petalobacterium. A name used by Maggi (1886, p 84) to designate a growth form of *Bacterium aceti* Zopf

Petalococcus. A name applied by Billroth (1874, p 6) to a growth form of his pleomorphic *Coccobacteria septica* in which the cells are spherical and occur arranged in a flat gelatinous mass or plate

This name has not been used as a generic designation, although Smith (1905, p 162) includes "*Petalococcus*" in his list of invalid bacterial names

Petalococcus. A name included by Erwin F. Smith (1905, p 162) in his list of rejected genera See *Petalococcus* Maggi (1886, p 81) gives *Petalococcus aceti* var *Duclauxi* as a variety of his *Micrococcus aceti*

Petalo-Gliabacteria. A growth form of *Coccobacteria septica* as described by Billroth (1874, p 17) A casual name

Pfeifferella. A generic name¹ proposed by Buchanan (1916) and (1918, p 54) as the seventh genus of the subtribe *Bacterinae*, with the description

Non-motile rods, slender, Gram-negative, without spores, staining poorly, sometimes forming threads and showing a tendency toward branching Gelatin may be slowly liquefied Do not ferment carbohydrates Growth on potato characteristically honey-like

The type species is the glanders bacillus, *Pfeifferella mallei*

Winslow *et al* (Committee Society of American Bacteriologists, 1920, p 200) used the same description and designate the type species as *Pfeifferella mallei* (Loeffler) Buchanan

Bergey *et al* (1923, p 389) follow Winslow It is possible that *Malleomyces q v* has priority

¹ This generic name for the glanders group came into being through a clerical error The author had intended to propose the name *Pfeifferella* for the influenza group of bacteria, and *Loefflerella* for the glanders group It was suggested by the Committee of the Society of American Bacteriologists that *Hemophilus* would be an appropriate name for the influenza group, it was intended therefore in manuscript in preparation to eliminate *Pfeifferella* Through some inexplicable clerical mistake *Loefflerella* was dropped and *Pfeifferella* substituted in mimeo-

Photobacillus. A generic designation used by Miquel and Cambier (1902, p. 881) for certain photogenic bacteria described by Kat (1891, p 159). No species name was used. The name cannot be considered as adequately published without specific names. It may be considered a casual designation. The name is rejected by Erwin F. Smith (1905, p 162).

Photobacter. A name proposed by Beijerinck (1900) for the photogenic bacteria. He named *Photobacter luminosum* and *Photobacter indicum* with two varieties, *Ph. splendidum* and *Ph. splendor*, likewise *Ph. phosphorescens* and *Ph. Fischeri*. These were studied primarily with reference to their ability to split urea.

Later (1900, p 197) he states that this is to be regarded as a physiologic genus including three unrelated form groups, and giving the inference that the name should have no nomenclatural status.

Enlows (1920, p 71) suggests that apparently Beijerinck uses *Photobacter* interchangeably with *Photobacterium* (q v). It is probably to be regarded as a synonym of the latter.

Photobacterium. A generic name proposed by Beijerinck (1889, p 401) to include certain photogenic marine bacteria. He says

Comme il n'a pas été possible d'établir, entre les cinq (ou six) espèces que j'ai étudiées, de distinctions assez importantes pour autoriser quelque séparation générique, et comme, d'un autre côté, le besoin se fait sentir de pouvoir facilement, désigner par un nom ces organismes si intéressants, je me hasarde à les rapporter toutes au genre *Photobacterium*.

The species listed are *Ph. luminosum*, *Ph. phosphorescens*, *Ph. indicum* and *Ph. Fischeri*. The reasons for grouping these species, among them cocci, bacilli and spirilla, in a single genus are given as follows:

- 1 All the species grow best in media containing $3\frac{1}{2}$ per cent NaCl
- 2 All lose the power of photogenesis by the addition of 2 per cent or more of glucose
- 3 Peptone is the principal nutrient
- 4 A trace of acid stops photogenesis
- 5 All are motile and do not form spores
- 6 None produce any diastatic or inverting enzyme
- 7 All give a similar spectrum

graphed outlines distributed at the meeting of the Society of American Bacteriologists. This constituted valid publication, and to prevent further confusion no effort has been made to correct the error. While *Pfeifferella* is not so appropriate in this connection, it is not invalid in consequence.

Bernhard Fischer (1894, p 19, *et al*) notes the occurrence in the Atlantic Ocean of the West Indian Light Bacillus (*Photobacterium indicum*) He also described as new species *Ph delgadense*, *Ph digenerans*, *Pg tuberosum*, *Ph papillare*, *Ph glutinosum*, *Ph annulare*, *Ph. coronatum*, *Ph caraibicum* All are comma bacilli, showing considerable variation in form

Migula (1897, p 336) insists that this genus is to be regarded strictly as physiologic, and invalid in nomenclature All with the exception of Beijerinck's *Ph phosphorencens* he placed in *Microspira* He states that to bring all these forms together into a single genus is from a systematic point of view quite incorrect, for they belong not only in different genera, but in different families He states further that Beijerinck's *Ph phosphorescens* is a true *Micrococcus*, the *Ph delgadense* of Fischer a *Pseudomonas* and *Ph coronatum* a *Microspira* Apparently he is mistaken with reference to *Ph phosphorescens*, for although Beijerinck gives as a synonym *Micrococcus phosphorescens* Cohn, the pictures show the organism to be a rod Possibly Migula's suggestion that all these forms should be included in *Microspira* is appropriate The name is rejected as invalid

The type species is designated by Enlows (1920, p 71) as *Photobacterium phosphorescens* (Cohn) Beijerinck

If the phosphorescent bacteria of this group are regarded as deserving of generic recognition, *Photobacterium* may be regarded as a valid designation, otherwise it may be regarded as a synonym (probably of *Vibrio*)

Photobakterium. A spelling of *Photobacterium* used by Flugge (1896, p 333) and other German writers

Photomonas. A name suggested by Orla-Jensen (1921, p 271) in the following statement "As the phosphorescent bacteria form, biologically, a connected whole, I deem it correct to unite them so as to form one genus, for which a better name than *Photomonas* can hardly be found "

No species is named Probably may be regarded as a synonym of *Photobacterium*, and as invalid

Photospirillum. A generic name proposed by Miquel and Cambier (1902, p 888) for the species *Photospirillum dunbari*, described by Dunbar (1893, p 799) Enlows (1920, p 71) states "Photogenic Finest light is produced at 22° on gelatin prepared from peptonized beef bouillon. Pathogenic for guinea-pigs " The type species is therefore *Photospirillum dunbari* Miquel and Cambier. The genus may probably be regarded as a synonym of *Photobacterium*

Phragmidiothrix. Engler (1882, p 19) named an organism found on the body of a crustacean (*Gammarus locusta*) *Beggiatoa multiseptata* Later (1883, p 187), his diagnosis was as follows

Phragmidiothrix, Engl Filis rectis vel leviter flexuosis, gelatinosis, cellulis brevibus egranulosis *Ph multiseptata* Engl Cellulis brevissimis, saepe diametro diversis, multis semel vel pluries septatis

Zopf (1883, p 104) (1885, p 103) gave the name *Phragmidiothrix multiseptata* ascribing the name to Engler His description is "Faden scheidenlos, successiv Theilungen sehr weit gehend Zellen schwefelfrei Wasserbewohner" Flugge (1886, p 398) defined the genus as one showing variable growth forms

Trevisan (1889, p 10) included this genus with its single species as the third of his tribe *Leptotrichae*

De Toni and Trevisan place this organism next to the genus *Beggiatoa* among the filamentous water bacteria with the following description

Filamenta evaginata, articulata, articuli diametro longitudinali brevissimo, simplicia, ubique aequalata, basi affixa Multiplicatio cellularum articularum divisione in duas directiones (septis transversis et longitudinalibus in coccis (an sporiferis?) abeuntium Baculogonia nulla

The species is described as follows

Filamentis 3-6 μ in diameter, articulis diametro longitudinali quadruplo-sextuplo breviori quam transversali Hab in aqua marina, ad Gammarum Locustam affixa, in sinu Kulensi Germaniae (Engler) Cocci quamquam adhuc filamentis matricialibus incluse, huc illuc germinant novaque filamenta procreant, quae ob rem filamenta primigenia lateraliter ramosa, videntur

Hueppe (1891, p 31) says "Die Faden sind in niedrige Cylinder-scheiben gegliedert, welche in Halbscheiben, Quadranten und schliesslich in Kugeln zerfallen"

Hansgirk (1891, p 313) concluded *Phragmidiothrix* should be included in the genus *Crenothrix*, for the filaments of *Phragmidiothrix multiseptata* are not as Engler stated, sheathless, but are enclosed in a thin sheath which in all respects corresponds with that of *Crenothrix* He states that a species *Crenothrix marina* previously named by him (1890, p 20) is probably *Crenothrix multiseptata* or even *C foetida* (Fiorini-Mazzanti), or *C mucor* (Oerstedt) Hansgirk concludes that the genus *Crenothrix* Cohn may be divided into two sections, *Phragmidiothrix* including the marine forms with *C foetida* as the type and *Eucrenothrix* with fresh water forms and *C Kuhniana* as the type.

Migula 1894, p 238

Phragmidiothrix Engler, Zellen zu anfangs unverzweigten Faden verbunden, sich nach 3 Richtungen des Raumes teilend und so einen Zellenstrang derstellend. Später kommen einzelne Zellen durch die sehr feine eng anliegende Scheide hindurchgewachsen und zu Verzweigung Veranlassung werden

Migula (1895, p 38) believed also that *Phragmidiothrix* might be united with *Crenothrix*, but set aside the genus, "So lange aber nicht erwiesen ist, dass die von Engler einmal beobachteten seitlichen Ausstrahlungen keine Verzweigungen, sondern Epiphyten sind, ist sie als Gattung beizubehalten." His description of the genus is as follows

Fadenbacterium mit sehr zarter, eng anliegender nur alten Faden sichtbarer Scheide, bis 100 μ lange und 3-12 μ breite Faden bildend. Die Faden bestehen anfangs aus flachen Scheibenzellen, welche sich jedoch später nach 3 Richtungen des Raumes teilen und so *Sarcina*-artige Packete bilden. Die einzelnen Zellen runden sich ab und werden aus der wie es scheint gleichzeitung zerfließenden Scheide frei

The genus was again recognized by Engler in 1897 (p 47) and 1900 (p 1034), Chester (1901, p 371) gives the same description. Kendall (1902, p 484) says "Cell division in three planes, very delicate sheath." Others who have used the genus are Smith (1902, p 272), Migula (1904, p 145), E. F. Smith (1905, p 162), Ellis (1909, p 9), Frost (1911, p 59)

The type species is *Phragmidiothrix multiseptata* Engler. The genus probably should be merged with *Crenothrix*.

Phycobacteriaceae A family name used by Engler (1912) for the filamentous bacteria surrounded by a more or less definite sheath, and which divide but rarely in three planes. It is a synonym of *Chlamydobacteriaceae*.

Phytobacter. A generic name proposed by Groenewege (1912, p 217) for the species *Phytobacter lycopersicum*. Enlows gives the description of the species as follows

Rods of varying length, 1.5 to 2.5 μ by 0.5 to 0.7 μ . No spores. Very slightly resistant to heat. Young cultures motile. Zoogloea in old cultures, which appear as a complex of rods bound together by a viscous slime. Found in decaying tomato fruits.

Phytomonas. A name proposed by Bergey *et al* (1923, p 174) for a genus of bacterial plant pathogens. The genus is included as the second in the tribe "*Erwiniae*." The description is as follows:

"Rods, yellow or white, motile or non-motile, the motile species possessing either mono- or lophotrichous flagella. May or may not have yellow pigment"

The type species is *Phytomonas campestris* (Pammel) Bergey

Phytomyxa. A generic name used by Schroter (1886, p 135) for the organisms causing nodules upon the roots of leguminous plants. This organism had previously been named *Schinzia leguminosarum* by Frank (1879, p 376). Since the generic name *Schinzia* (qv) was preoccupied, Schroter (1886) gave the name *Phytomyxa leguminosarum*, including this organism among the *Myxomycetes* in the order *Phytomyxini*. He also named a similar species from the lupine, *Phytomyxa Lupini*. He based his inclusion of this genus with *Plasmodiophora* among the slime molds on the work of Prillieux (1879, p 98). The plasma-like filaments in the nodules he interpreted as the vegetative growth, and the bacterioids as spores. Beijerinck (1888, p 758) named this organism *Bacillus radicolus*. It was again included among the slime-molds a year later by Schroter (1889, p 7). Frank (1889, p 338) concluded that the assignment of the organism to the slime molds was incorrect, and that a new generic name should be chosen. He used *Rhizobium leguminosarum*. More recently this organism has been known by several names, as *Pseudomonas radicolus*, *Bacteridium radicolus* (Fischer 1903, Lotsy 1907), *Bacterium leguminosarum* (Smith 1905). Maire and Tison (1909, p 241) have emphasized the fact that even though the genus *Phytomyxa* was erroneously included by Schroter among the *Myxomycetes* this did not invalidate the genus name.

Phytomyxa is probably technically a valid generic name for organisms of the type which cause nodule formation on the roots of legumes if this type is worthy of generic recognition, with the type *Phytomyxa leguminosarum* (Frank) Schroter.

However, Winslow *et al* (Committee Society of American Bacteriologists 1917) recommended that *Rhizobium* be recognized as a *genus conservandum* for this group. In this they have been followed by Buchanan (1918), Bergey (1923) and others.

Planococcus. A genus of cocci proposed by Migula (1894, p 236) who says *Planococcus* nov. gen. Die Zellen teilen sich nach 2 Richtungen des Raumes wie bei *Micrococcus* besitzen aber geisselformige

Bewegungsorgane (Z B *Planococcus citreus* (Menge) Mig'') The generic diagnosis given later is

Zellen einzeln oder zu 2 oder 4 genahert, oft in grosserer Zahl unregelmässige Haufen bildend Die einzelnen freien Zellen Kugelförmig, Teilung abwechselnd nach 2 Richtungen des Raumes Die Zellen sind frei beweglich Die Bewegung wird vermittelt durch 1-4, meist 1 Geissel, die in der Regel vielmal länger ist als die Zelle

This generic name has been used quite generally by writers on systematic bacteriology, among others by Migula (1897, p 46, 1900, p 269, 1904, p 144), Fischer (1897, p 32, 1903, p 59), Chester (1897, p 63, 1901, p 114), Kendall (1902), A J Smith (1902, p 270), E F Smith (1905, p 160), Ellis (1909, p 4), Frost (1911, p 57)

Meyer (1912, p 5) uses *Planococcus* as the designation of a section of *Micrococcus* with the description "Die begreisselten Formen"

Vuillemin (1913, p 525) proposes that this be made a "*genus conservandum*" with the type species *Planococcus europaeus* (*Nitrosomonas europaea* Winogradsky)

The genus *Planococcus* is valid if this character of motility is sufficient to differentiate from *Micrococcus*, otherwise is a synonym of the latter name The type species is *Planococcus citreus* (Menge) Migula

Planomerista A "*genus conservandum*" proposed by Vuillemin (1913, p 525) to include those cocci which divide regularly in two planes and which remain united in tetrads and are motile It is a motile *Merista* (q v) The type species cited is *Planomerista ventriculi* (*Micrococcus tetragenus mobilis ventriculi* Mendoza 1889)

The genus name is valid if the organisms of this type are worthy of generic recognition, otherwise it is to be regarded as a synonym of *Micrococcus*, or *Pediococcus*

Planosarcina A genus of cocci, created by Migula (1894, p 236) who says "*Planosarcina* nov gen Die Zellen teilen sich wie bei *Sarcina* nach 3 Richtungen des Raumes, besitzen aber geisselförmige Bewegungsorgane (Z B *Planosarcina agilis* (Cohen) Migula, *Pl mobilis* (Mauria) Mig'") The cells when isolated are quite spherical usually arranged in pairs and fours, and flattened somewhat on the contiguous sides Division occurs in three directions of space, the cells may remain united into packets or break apart The packets are usually developed only in sugar media Cells are actively motile by means of flagella The two species (1895, p 19) first mentioned are *Planosarcina mobilis* and *Pl agilis*

This genus has been frequently recognized in the literature of systematic bacteriology. The following authors among others have used it. Migula (1897, p 46, 1900, p 275, 1904, p 144), Chester (1897, p 63, 1901, p. 115), Fischer (1897, p 32), Kendall (1902), A J Smith (1902, p 270), Fischer (1903, p 60), E F Smith (1905, p 160), Ellis (1909, p 4), Frost (1911, p 57)

Vuillemin (1913, p 525) has proposed that *Planosarcina* be recognized as a "*genus conservandum*"

Meyer (1912, p 5) uses *Planosarcina* as a designation for a section of the genus *Sarcina* with the description "Die begeisselte Spezies"

If the character of motility is sufficient to differentiate this group, the name is valid, but it is probably to be regarded as a synonym of *Sarcina*. The type species is *Planosarcina agilis* (Cohen) Migula

Planostreptococcus A name used by Meyer (1912, p 4) to designate a section of the genus *Streptococcus*. The description reads, "Begeisselte Spezies". No species are named

Plectridieae. A subfamily of bacteria created by Fischer (1897, p 33) to include those rod shaped organisms which become drumstick shaped during sporulation. The only genus included is *Plectridium*

Plectridium. A genus of rod-shaped bacteria named by Fischer (1895, p 147) with the following definition "Beweglich mit diffusen Geisseln und Endosporen in einem kopfig geschwollenen Ende der Stabchen (Kopfchenbakterien, Kloppel-, Trommelschlagen- oder Kaulquappenform), das andere Ende nicht geschwollen". Three species *Plectridium paludosum*, *P. tetani* and *P. des Rauschbrandes*. It is recognized that the organisms are anaerobic

Hueppe (1895, p 38) notes the genus. It is again described by Fischer in (1897, p 32) and (1903, p 61). In the latter he notes as species the butyric acid bacilli, the tetanus bacillus, and a decay form, *Plectridium putrificum*

Buchanan (1918, p 38) emended the genus as follows

Cells rod-shaped, straight or at least never spiral, motile by diffuse flagella or non-motile. Endospores produced under favorable conditions, causing an enlargement of one tip of the cell, giving rise to a drumstick appearance. Usually Gram-positive. Anaerobic or microaerophilic

The type species is *Plectridium tetani* (Nicolai) Fischer

The same type species was later (1922) used by Heller for the genus *Nicolaiellus*, which becomes a synonym

It is possible that *Plectridium* is not sufficiently distinct to require recognition separate generically from *Clostridium*. It is made a synonym of *Clostridium* by Winslow *et al* (Committee Soc Am Bact., 1917, and 1920) and by Bergey *et al* (1923).

Plectrillum. A generic name proposed by Fischer (1895, p 144) for rods motile by means of a tuft of polar flagella, and with spores terminal and causing enlargement of the rod (Drumstick shapes). No species were named.

Plectrinum. A genus of rod shaped bacteria named by Fischer (1895, p 142) with the following definition

Plectrinum nov gen. Beweglich mit polarer Einzelgeissel, Endosporen in einem aufgeschwollenen Ende. Auch für diese Gattung fehlt es vorläufig noch an einem sicheren Beispiel. Zopf bildet zwar eine Köpfchenbakterie mit einer Geissel am nicht geschwollenen Ende ab, giebt aber weder den Fundort noch sonst etwas an.

No species was named. The genus is apparently invalid.

Plennobacterium. A spelling of *Plennobakterium* (qv) used by Kossowicz and others.

Plennobakterium. Gonnermann (1907, p 887) named an organism belonging to the hay bacillus group *Plennobakterium*. It is non-motile and gram positive, forms spores and liquefies gelatin. The rods are 0.4 to 0.6 wide, 2.5 to 5.0 μ long. Sometimes in filaments. Characteristic spiral bands were observed within the cell. It was isolated from sugar beet sap.

No species is named. It may be regarded as a casual designation and not entitled to generic standing.

The name is spelled *Plennobacterium* by certain authors as Kossowicz.

Pleurococcus. A genus of algae named by Meneghini (1842) to which certain bacteria have been assigned as *Pleurococcus Beigelii* Rabenhorst 1867 which became *Hyalococcus beigelii* Schroter and *Micrococcus beigelii* Migula, and *Pleurococcus roseo-persicinus* Rabenhorst which became *Lamprocystis roseo-persicinus* Migula.

Pleurospora. A subgenus of the bacterial genus *Cornelia* used by De Toni and Trevisan (1889, p 1002) with the description "*Sporae macrosomae e latere protuberantes*" to contain two species, *Cornelia tremula* and *C. sphinx*. The name has apparently not been used by other authors.

Plocamobacterium. Lowi (1920, p 730) proposed the name *camobacterium crassum* for *Bacillus crassus* Lipschutz. The organism is a gram positive facultative anaerobe, with characteristic growth on Drigalski agar, maltose nutrient agar and dextrose agar. The cells in young cultures are small rods, while in older cultures they are filamentous. Non-motile. No spores. A common inhabitant of the normal vagina, but occasionally pathogenic causing acute ulcer of the vulva.

Pneumobacillus. This name has been used commonly as a quasi-scientific name for the organism described by Friedländer as the causal organism of pneumonia. Arloing (1889, p 428) proposed the name *Pneumobacillus liquefaciens bovis* for the organism which he isolated as the cause of contagious peripneumonia of cattle. The use of the name in a trinomial would appear to invalidate it as a generic designation. It has never come into common use. This organism was later termed *Bacterium bovis* (Arloing). Migula (1900, p 442).

The name is placed by Erwin F. Smith (1905, p 162) among the rejected genera.

Pneumococcus. A name usually used as a casual designation of the organism of pneumonia. It was apparently first used in a generic sense by Arloing (1889, p 430) who named three species, *Pneumococcus gutta-cerei*, *P. lichenoides* and *P. flavescens*, all present together with *Pneumobacillus liquefaciens bovis* in bovine pleuropneumonia. The species are inadequately described.

It has occasionally apparently been used in a generic sense by other writers as witness Schmidlechner (1905, p 291) who uses the designation *Pneumococcus lanceolatus* Fraenkel in a list of species.

The name *Pneumococcus mucosus* is used also by Dochez and Avery (1915, p 114) and by Lyall (1915, p 146). It is placed by Erwin F. Smith (1905, p 162) in his list of rejected genera.

Pneumococcus is probably best regarded as a synonym of *Diplococcus*.

Pollendera. According to De Toni and Trevisan (1889, p 943) this generic name was used by Trevisan (1884) for the anthrax bacillus which he named *Pollendera anthracis*.

The original reference to this use by Trevisan has not been discovered by the writer. The name has not been used by other authors, and is rejected later by Trevisan as a synonym of *Bacillus*.

It would seem that this name would become valid if non-motile rod-shaped bacteria are to be grouped in a genus. It certainly would have priority over *Bacterium* in the sense of Migula and over *Aplanobacter*. E. F. Smith.

It may be noted that Smith (1905, p 174) rejects the name *Pollendera* (which he spells *Pollendra*) at the same time he proposes *Aplanobacter*

Pollendra. A spelling of *Pollendera* (*qv*) used by Erwin F Smith (1905, p 174)

Polyangium. This genus was first created by Link (1795, p 40) according to Zukal (1897, p 550) for the species *P vitellinum*. He regarded it as a species of fungus, and as such it appeared in several floras and works on fungi. What is evidently the same species was described later by Thaxter (1892, p 394) as *Myxobacter aureus*. He first recognized the true affinities of the organism as a genus of the *Myxobacteria*, and that it is not a true fungus, nor, as had been urged by Bonordon, an insect egg. That the genus *Myxobacter* of Thaxter is a synonym of *Polyangium* Link was pointed out by Zukal (1 c). *Myxobacter aureus* Thaxter therefore became *Polyangium vitellinum* Link. This explanation was accepted by Saccardo (1899, p 844) who states

Polyangium Link Syll VII p 47 (inter dubia) Huc ducendum est genus *Myxobacter* Thaxt Syll XI p 461 cuius species *M aureus* Thaxt collidit cum *Polyangio vitellino* Link jam ab anno 1795 descripto, sed huiusque inter dubia, relato

This was accepted later by Thaxter (1904, p 37)

Cystobacter Schroter (*qv*) is given by Thaxter as a synonym, in part. Seven species are recognized, as follows: *Polyangium vitellinum* Link, *P simplex* Thaxter, *P fuscum* (Schroeter) Thaxter, *P septatum* Thaxter, *P compositum* Thaxter, *P solediatum* Thaxter, and *P primigenium* Quehl

The genus is apparently valid. It is so recognized by Smith (1905, p 165) and by Quehl (1906, p 16)

The diagnosis of the genus given by Thaxter is "Rods forming large, rounded cysts, one or more free within a gelatinous matrix raised above the substratum." Together with the genera *Myxococcus* and *Chronodromyces*, it constitutes the family *Myxobacteriaceae*

Buchanan (1918, p 542) included this as the second genus of the family *Myxobacteriaceae* with the following description

Rods forming large rounded cysts, one or more, free within a gelatinous matrix raised above the substratum

The type species is *Polyangium vitellinum* Link

Bergey *et al* (1923, p 416) followed Buchanan

Polybacteria. A genus (?) of bacteria named by Van Tieghem (1880, p 149) to include bacteria growing in naked masses made up of cylindrical cells *Polybacteria catenata* was found in a decoction of horse manure A *Polybacteria sulphurea* was described from rotting beans

The descriptions are so inadequate that probably the organisms cannot be identified The generic name does not seem to have been used by other authors

Polycephalum. A fungus genus named by Kalchbrenner and Cooke (1880, p 22) with the species *Polycephalum aurantiacum* Thaxter (1892, p 399) states that this is a possible synonym of *Chondromyces* (*q v*)

Propionibacterium. A genus of bacteria proposed by Orla-Jensen (1909, p 337) The genus is included in the family *Acidobacteriaceae*, order *Peritrichinae* It includes those rod-shaped bacteria which ferment lactose and lactates with formation of considerable amounts of propionic acid

Orla-Jensen's statement is as follows

Da die Propionsäurebakterien wie die Coli-Bakterien sich dadurch auszeichnen, dass sie die gebildeten Spaltungsprodukte sowohl oxydieren als auch reduzieren können, ist es natürlich, auch sie von den letzteren abzuleiten In kultureller Beziehung nehmen sie eine Zwischenstellung zwischen den Aerogenes-Bakterien und den Milchsäurebakterien ein, indem sie in der Stichkultur sich auf der Oberfläche um so weniger ausbreiten, je grösser ihre Fähigkeit zur Bildung von Propionsäure ist Sie bilden unbewegliche Stäbchen oder Diplokokken, die völlig an *Streptococcus lacticus* erinnern Sie vergären sowohl Laktose als auch Laktate Sie greifen das Kasein nicht an, sondern spalten die von den peptonisierenden Bakterien gebildeten Produkte weiter Wegen ihrer reduzierenden Eigenschaften entwickeln sie Schwefelwasserstoff, wenn sie Pepton zur Verfügung haben Wir wollen die Gattung *Propionibacterium* nennen

Later (1921, p 272) he again emphasizes the genus

The status of the genus is doubtful, as no species is described or referred to

Propionicoccus A generic name suggested by Orla-Jensen (1921, p 272) in the following statement

Whereas the shape of cells was formerly used as a family character, I have adopted it only as a generic one, and if we do not want to further confine its significance and only consider it as a specific character, we shall doubtless have to set up the genus *Propionicoccus* besides the genus *Propionibacterium*

Proteae. The seventh tribe of the family *Bacillaceae* named by Castellani and Chalmers (1919, p 932) with the description: "*Bacillaceae* growing well on ordinary laboratory media, not forming endospores, aerobic, without fluorescence or pigment formation, but liquefying gelatine"

Two genera are included, *Proteus* and *Cloaca*.

Proteobacter. A name used by Beijerinck (1900, p 195) In a discussion of hydrogen sulphide production he says,

Wenn die Quantität der organischer Körper im Schlamm sehr gross ist, wie, z B in den Stadtgraben zu Delft, wo Gerberien und Brennerieien ihre Abwasser seit langer Zeit darin geleert haben, so sind die Lebensbedingungen für die gewöhnlichen Anaeroben der Eiweissfaulnis gegeben, während *Spirillum desulfuricans* infolge der Anhaufung organischer Körper, zurücktritt Von den bei der Eiweissfaulnis in Betracht kommenden Arten habe ich die drei wichtigsten (*Proteobacter septicum*, *P shatol*, *P pseudopulcher*) bei einer anderen Gelegenheit besprochen. Es hat sich gegeben, dass diese Mikroben nicht nur Schwefelwasserstoff erzeugen, sondern auch die entsetzlichen Sulfiden der Mercaptangruppe

Proteus. A generic name used by Hauser (1885, p 1) to designate three species of putrefactive bacteria, *Proteus vulgaris*, *P mirabilis* and *P Zenkeri*. No discussion or definition of the generic name is given apart from the species description. All the species agree in showing great variability in morphology, short and long rods, filaments, and spirals, and many involution forms may be observed. All are motile by means of peritrichous flagella, none produce spores. All the species produce "islands" on the surface of moist solid media, these seem to be endowed with some motility. All the species are closely associated with decay and putrefaction.

This name is regarded as a synonym of *Bacillus* by De Toni and Trevisan (1889) and of *Spirulina* by Hueppe (1891). The genus is recognized by Cornil and Babes (1890, p 151) subsequent writers have practically all until recently regarded the term as synonymous with *Bacillus* and have not considered its pleomorphism as distinctive.

Lehmann and Neumann (1896, p 103) make it a synonym of *Bacterium*.

Macé (1897, p 487) says that *Proteus* cannot be separated from *Bacillus*. Erwin F Smith (1905, p 174) includes *Proteus* in his list of genera to be rejected. Orla-Jensen (1909, p 338) makes *Proteus* a synonym of *Liquidobacterium*.

Buchanan (1918, p 54) emended the generic description as follows:

Short rods, showing great variation in morphology, filamentous and bent rods as involution forms frequent Motile by means of peritrichous flagella The species commonly produce motile "islands" on the surface of moist solid media No spores Gram-negative Usually producing acid and gas from certain carbohydrates In general the species are closely associated with decay and putrefaction, sometimes pathogenic

The type species is *Proteus vulgaris* Hauser

Winslow *et al* (Committee Soc Am Bact, 1920, p 210) have described the genus as follows

Highly pleomorphic rods, filaments and curved cells being common as involution forms Gram negative Actively motile Characteristic amoeboid colonies on moist media Liquefy gelatin rapidly and produce vigorous decomposition of proteins Ferment glucose and sucrose (but usually not lactose), with formation of acid and gas (the latter being CO₂ only)

Type species, *P vulgaris* Hauser

The genus is also recognized by Castellani and Chalmers (1919, p 943)

Bergey *et al* (1923, p 209) include *Proteus* as the third genus of the tribe *Bactereae* with the following description

Highly pleomorphic rods Filamentous and curved rods are common as involution forms Gram-negative Actively motile, possessing peritrichous flagella Produce characteristic amoeboid colonies on moist media and decompose proteins Ferment dextrose and sucrose but not lactose Do not produce acetyl-methyl carbinol

Protobacteriés. A name used by Maggi (1886, p 81) as the designation of his first order of bacteria, with one genus *Aphaneroglia*, but without description

Protococcus. A genus of green algae proposed by Agardh One species *Protococcus roseo-persicinus* Kutzing (1849) was later described by Schroter as *Lamprocystis roseo-persicina* and included among the bacteria *Protococcus* is a valid algal generic name and should not be used for the bacteria

Miller (1892, p 123) makes the following statement

Klencke (1850) beschreibt auch einen, von ihm entdeckten Parasiten, eine *Protococcus* art, *Protococcus dentalis*, welcher Schmelz und Zahnbein in ahlicher Weise verflüssigt, wie der Hausschwamm, *Merulius lacrymans*, das Holz der Hauser oder Möbel erweicht

Protomycetum. Lewis (1879, p 394) makes the following statement concerning *Spirillum obermeieri*

Since the period of its discovery in the blood by Obermeier it has been referred to under various names *Spirothrix*, *Protomycetum recurrentis* in Lebert's article on recurrent fever and in Ziemss' 'Handbuch' of Medicine, *Spirillum* by Erichsen, Litten, Birch-Hirschfeld etc , *Spirillum tenue* by Naunyn, and *Spirochoete Obermeieri* by Cohn (fig 15)

Pseudobacterium. Trevisan (1888, p 781) included this name in the following sentence

Senze dipartirsi da *Klebsiella*, basterebbe ricordare come tra le specie che attualmente se ne conoscono, altre s'ebbero le denominazioni generiche *Micrococcus*, *Diplococcus*, *Pseudodiplococcus*, ed altre quelle di *Bacterium*, *Pseudobacterium*, *Proteus*, *Bacillus*, *Ascobacillus*, etc

What organism is referred to is not evident

Pseudodiplococcus. A generic name used by Bonome (1888, p 321) to designate *Pseudodiplococcus pneumonicus*, an organism which he isolated from a fibrinous exudate The organism is stated to resemble closely morphologically and biologically the "*Diplococcus pneumoniae*"

The organism is termed *Klebsiella Leydeni* by Trevisan (1889, p. 1030) The genus has not been recognized by other authors It may be regarded as a synonym of *Diplococcus*

Pseudomeningococcus A casual name proposed by Elser and Huntoon (1909, p 384) for "a group of organisms which cannot be differentiated from the meningococcus excepting by serum reactions"

Pseudomonadaceae. The third family of the *Eubacteriales* named by Winslow *et al* (Committee Soc Am Bact, 1917, p 555) with the following description

Short rods, usually motile Flagella single, polar Gram-negative Not obligate aerobes Many species active ammonifiers Many species produce water-soluble pigments or green fluorescence, yellow pigment common Some species are photogenic Soil and water bacteria, with many plant parasites

The single genus *Pseudomonas* is included

Pseudomonadales A name proposed by Orla-Jensen (1921, p 270) as one of the two orders into which he would divide *Eubacteriales* Buchanan. He would include in this group some eleven genera as follows 1. *Methanomonas*, 2 *Carboxydomonas*, 3 *Hydrogenomonas*, 4. *Nitrosomonas*, 5 *Nitromonas*, 6 *Azotomonas*, 7 *Rhizomonas*, 8. *Acetimonas*, 9 *Fluormonas*, 10 *Photomonas*, 11 *Spiromonas*.

Pseudomonas. A genus name proposed by Migula (1894, p 237) with the following description "*Pseudomonas* nov gen Zellen mit polaren Bewegungsorganen, Endosporenbildung kommt bei einigen Arten vor, ist aber seltener (A B *Pseudomonas violacea*)"

It will be noted that no clue to what previously described *violacea* he is referring is given This cannot therefore be accepted as the type, and the genus was technically not adequately published until the next year (1895, p 29) when the following description is given

Kurzer oder langercylindrische Zellen, welche zuweilen kleine Faden bilden, lebhaft beweglich mit polarer Begeizung Die Zahl der an einem Pol stehenden Geizeln schwankt bei den verschiedenen Arten zwischen 1-10 und ist am häufigsten 1 oder 3-6 Endosporenbildung kommt vor, aber nur bei wenigen Arten

Eine Trennung der hierhergehorigen Arten in zwei Gattungen, je nach dem am Pol nur 1 Geizel oder ein Buschel von Geizeln steht, wie dies von Fischer vorgeschlagen wurde, ist unthunlich, da alle Übergänge zwischen streng eingezelligen und vielgeizeligen Arten vorhanden sind

The first species named and described is *Pseudomonas pyocyanea* which in its corrected form, *Pseudomonas aeruginosa* (Schroter) may be regarded as the type

Migula (1900, p 876) notes the close relationship of this genus to *Microspira* in the following statement

Die Grenze gegenüber *Microspira* ist vielleicht eine kunstliche, es ist nicht immer möglich, zu beurteilen, ob es sich bei eingezelligen Formen um eine *Microspira* oder eine *Pseudomonas* handelt Leichte Krümmungen, namentlich in gefärbten Präparaten, beobachtet man bei vielen *Pseudomonas*arten Gewöhnlich lässt sich die Entscheidung, zu welcher Gattung eine Art gehört, durch Beobachtung von mehreren zusammenhängenden Zellen ermöglichen Bei *Pseudomonas* kommen dann niemals regelmässige Schraubenwindungen vor

The genus has been accepted by many subsequent writers, among them Migula (1897, p 47, 1900, p 29, 1904, p 145), Chester (1901, p 306), Schmidt and Weis (1902, p 91), A J Smith (1902, p 270), Ellis (1909, p 6), Frost (1911, p 57), Schneider (1912, p 23) and Meyer (1912, p 3)

Conn (1909, p 12) defines the genus as made up of rod-shaped bacteria with a single flagellum

Erwin F Smith (1905, p 160) and Vuillemin (1913, p 522) independently reached the conclusion that the generic name *Bacterium* (*qv*) has priority for this group of organisms

Winslow *et al.* (Committee Soc Am Bact, 1917, p 556) used the following generic description

Rod-shaped, short, usually motile by means of polar flagella or rarely non-motile Aerobic and facultative Frequently gelatin liquefiers and active ammonifiers No endospores Gram stain variable, though usually negative Fermentation of carbohydrates as a rule not active Frequently producing a water-soluble pigment which diffuses through the medium as green, blue, purple, brown, etc In some cases a non-diffusible yellow pigment is formed Many yellow species are plant parasites

Buchanan (1918, p 48) discussed at length the nomenclatural status of the genus, and gave the following description

Rod-shaped bacteria, never spiral, usually motile by means of polar flagella or rarely non-motile Aerobic and facultative Frequently liquefying gelatin Without spores Gram stain variable Usually producing a water-soluble pigment which diffuses through the medium as a green, blue, purple, or brown or in some cases a yellow pigment Fermentation of carbohydrates usually not active

The type species is *Pseudomonas aeruginosa* (Schroter) Frost?

Later Winslow *et al* (1902, p 204) designated the type species as *Ps aeruginosa* (Schroter) Frost? Enlows (1920, p 74) states, "*Type species* (monotype)—*Pseudomonas violacea* (Schroter 1886) Migula" This designation can be made only by reference to Migula's publications after the first in which *Pseudomonas* was proposed There had been previously described

Pediococcus violaceus Trevisan 1889

Lampropedia violacea Trevisan 1889

Thiocystis violacea Winogradsky 1889

Spirillum violaceum Warming

Bacteridium (*Bacillus*) *violaceum* Schroter

Chromobacterium violaceum Bergonzini

Chromococcus violaceus Bergonzini

Micrococcus violaceus Cohn

Streptococcus violaceus (Cohn) Trevisan

Migula (1895) particularly emphasizes the fluorescent pigment production as characteristic The first species described is one which is typical and should be designated as the type as was done by Winslow *et al*

Bergey *et al* (1923, p 122) include *Pseudomonas* as the fourth genus of the tribe *Chromobacterae* with the following description: "Principally water and soil bacteria producing a water-soluble pigment which

diffuses through the medium as green, blue or yellowish-green Motile or non-motile Gram-negative "

Pseudorhizobium. A generic name used by Hartleb (1900, p 887) for an organism isolated from the nodule of a legume. He named the form *Pseudorhizobium ramosum*. The organism resembled the normal legume bacteria but appeared to be non-infectious. The name has apparently not been used by other writers.

Pseudosarcina. According to Lohms (1913, p 449) the genus *Pseudosarcina* was named by Macé (1903, p 887). An examination of the statement of the latter author fails to show such generic name. The term used is pseudosarcine, employed apparently simply in a casual sense. He says

Ce microbe se présente insolé, et alors il est sphérique, ou sous forme d'agregats plus ou moins volumineux, d'un aspect mûriforme rappelant une grosse sarcine, en raison de cette analogie je le désignerai provisoirement par le terme de pseudosarcine car il ne semble pas que ses bipartitions se fassent suivant deux plans perpendiculaires.

Pseudo-sarcine. See *Pseudosarcina*.

Pseudospira. A subgenus of the genus *Pacina* used by Trevisan (1889, p 1018) with the following diagnosis

Baculi curvi, non raro semicirculares, saepissime in filamenta undulato-flexuosa vel irregulariter pseudospirilia, nunquam vere spiraliter ut in *Spirille*s torta, consociati. Arthrosporaë notabiliter quam in *Eu-Pacini*s minores, nunquam duplo diametri transversalis baculorum latiore.

Eight species are included in the subgenus. The type species may be designated as *Pacina cholerae asiaticae*. The organisms of this group are those ordinarily assigned to *Vibrio* or to *Microspira*.

Punctula. A generic name used by Van Tieghem (1880, p 150). The following summary is from Journal Roy. Mic. Soc. (1880, p 1001).

Punctula. The spherical cells are ordinarily extremely minute, they appear like innumerable dots united by a gelatinous cement. A close examination is required to distinguish the colonies composed of them from simple naked cells consisting of a finely granular protoplasm.

In *Punctula rosea* the colonies are of a bright rose colour, they are spherical, and with a sharply defined outline, the dots, which are so many elementary cells, are arranged in them with perfect regularity in radial rows and concentric circles. After each division, the two halves of the colony become rounded off, and separate completely. When one of these spheres is crushed, it is resolved into its elementary cells, and the formation can then be followed of so many new colonies by the repeated increase and division of each of the cells.

Two other species, *P. cubica* and *P. glomerata* are named

The species are so inadequately described as probably to represent practically *nomina nuda*.

Punctum. A generic name apparently proposed by Muhlhauser (1884, p. 97) Enlows gives the following summary

Very small (0.0005 mm) of varied form, but usual oval, at first not motile, later very lively, moving around in a circular fashion. It can traverse 0.1 mm in one second. In young cultures (stagnant water) the longer forms are found. No chains or filaments

Type species *Punctum saltans* Syn (Trevisan, Saccardo's Sylloge Fungorum, v. 8, 1889, p. 1008) *Spirillum obermeieri* Cohn, 1875 (Beit. z. Biol. der Pflanzen, v. 1, p. 196)

Putribacillus. A generic name proposed by Orla-Jensen (1909, p. 342) as one of the genera of his family *Putribacteriaceae*. It includes the anaerobic putrefactive bacteria which do not produce an exotoxin. Jensen makes the following statement:

Einen anderen Zweig bilden die anaeroben Faulnisbakterien, die Gattung *Putribacillus*. Diese sind dadurch gekennzeichnet, dass sie die Eiweiszstoffe unter Verbreitung eines furchtbaren Gestankes zersetzen. Die Entwicklung geht hier wie bei den aeroben Faulnisbakterien von saurebildenden zu saureempfindlichen Arten. So entsprechen z. B. *Bacillus bifermens sporogenes* und *Bacillus oedematis maligni* (welche Traubenzucker, aber nicht Milchzucker vergären), sowie *Bacillus perfringens* und *Paraplectrum foetidum* (die sowohl Traubenzucker als auch Milchzucker vergären, ja das letztere zerlegt sogar ebenso wie *Butyribacillus Chauvoei* Milchsäure) den Proteusbakterien, während *Bacillus putrificus*, den wir *Putribacillus vulgaris* nennen wollen, den aeroben Bacillen entspricht. Letztere Bakterie ist gar nicht zuckerspaltend, was sonst für fast alle fakultativ oder obligat anaeroben Bakterien charakteristisch ist, und sie muss sich daher die ganze nötige Energiemenge lediglich durch Zersetzung der Proteinstoffe verschaffen können. Innerhalb der anaeroben Bakterien hat sie deshalb die höchste Entwicklung als Eiweiszvergärer, ebenso wie *Cellulobacillus methanogenes* als Kohlenhydratvergärer, erreicht.

This genus perhaps may be regarded as a synonym of *Clostridium* or possibly of *Plectridium*.

It is difficult to determine what is regarded as the type species. The only one of the species mentioned which he specifically places in this genus is his *Putribacillus vulgaris* or properly *Putribacillus putrificus*. This may therefore be termed the type. With this understanding, this generic name has priority over *Putrificus* Heller (q.v.).

Putribacteriaceae. A family of bacteria proposed by Orla-Jensen (1909, p. 343) to include obligate anaerobic or microaerophilic bacteria

which act largely on proteins and bring about putrefactive changes. The family is included in the order *Peritrichinae*. It contains the following genera *Putribacillus* and *Botulobacillus*.

Putriclostridium. Orla-Jensen (1921, p. 273) states

The use of the term *Clostridium* as a generic name presents the inconvenience that under the same we must group together not only the clostridia but also the plectridia. The division into true butyric acid bacteria (*Butyriclostridium*), the requirements of which in regard to nitrogenous nutriment are very moderate (they are able to assimilate even the nitrogen of the air) and anaerobic, putrefying bacteria (*Putriclostridium*) seems natural to me.

Putrificoideae. A name given by Heller (1921, p. 550) to the second subfamily of the *Clostridiaceae* with the following description

Clostridiaceae which on meat medium produce after twenty days' incubation at 37° under vaseline a reaction of pH 7.1 or a more alkaline reaction, the reaction being read after the culture had been boiled.

Eleven genera are included *Segumillus*, *Regillus*, *Robertsonillus*, *Nicollaserillus*, *Martellillus*, *Recordillus*, *Tissierillus*, *Putrificus*, *Ermen-genillus*, *Metchnikovillus* and *Weinbergillus*.

Putrificus. A generic name proposed by Heller (1922, p. 27) with the following description

Putrificoideae that do not attack sugars. Slender Gram-negative or Gram-positive rods with oval terminal spores. Colonies in deep agar have radiate periphery and opaque center. Putrefactive organisms found in soil and wounds.

Type species *P. Bienstocki* (*Bacillus putrificus* Bienstock) as defined by Bienstock (1906). References, Bienstock (1884, 1901, 1906), Klein, Rodella (1905).

Pyobacillus. A name used by Kóppányi (1907, p. 429) to designate his *Pyobacillus capsulatus cuniculi* or *Bacillus pyemiae cuniculi*. The organism was isolated from rabbits. It is a non-motile capsulated rod, grows on weakly alkaline nutrient media only if at same time exudate is carried over on to the nutrient medium, or serum, or serum agar. Protein-rich substances necessary for growth. No growth on gelatin, room temperature not suitable. It probably is to be regarded as non-valid, as binomial nomenclature is not used.

Pyobacterium. A generic name used by Kuttner (1895, p. 760), for the species *Pyobacterium Fischeri*. The organism was found in an epigastric abscess. It is closely related to *Bacterium coli commune*, but is more slender, forms colonies pure white in color and is actively motile. Milk is slowly coagulated with flocculent curd. It ferments

sugars less vigorously and does not make meat infusion alkaline. Produces indol and nitrites. Pathogenic for mouse, guinea-pig, rabbits and pigeons. Animal infected intravenously quickly succumb to septicemia. In subcutaneous inoculation there is pus production (abscess formation).

Pyococcus. A term used by Ludwig (1892, p. 27) as a synonym of *Staphylococcus*. It evidently has not been used as a generic designation.

Radiobacter. A name occasionally used in a pseudo-generic sense in reference to *Bacillus radiobacter*, as by Stoklasa (1908, p. 484).

Radiococcus. A name used by Maggi (1886, p. 81) to designate a growth form of his *Micrococcus aceti*.

Rasmussenia. A genus of bacteria created by De Toni and Trevisan (1889, p. 930) to include the slender filamentous bacteria of the mouth usually known as *Leptothrix*. The generic diagnosis is as follows.

Filamenta cylindrica, vagina tenui gelatinosa facile evanescente obducta, simplicia, basi subiculo zoogloeico affixa, articulata, articulis aetate proventa in baculos et coccus solutis. Multiplicatio baculis primitus vivacites mobilibus, cito immotis. Arthrospora transformatione cocci singuli ortae.

Five species are described: *R. gigantea*, *R. buccalis*, *R. maxima*, *R. anceps* and *R. variabilis*. The type first described may probably be regarded as *Rasmussenia gigantea* (Miller) Trevisan, from the mouth.

Clements (1894, p. 28) is apparently the only other author who has used this generic designation. His generic description is "Cells elongate-cylindrical, flexuose, simple, with a thin, gelatinous sheath, often separating into bacilli or cocci, from which the arthrospores arise."

One species is described, *Rasmussenia buccalis* (Rob. and Leb.) Trev.

Unless the type, *R. gigantea* (Miller) Trevisan is so different from *R. buccalis* (Rob. and Leb.) Trev. as to warrant genera for the two species, the genus *Rasmussenia* is a synonym of *Leptotrichia* *q. v.* of which *L. buccalis* is the type.

Recordillus. A name proposed by Heller (1922, p. 26) for a genus of the tribe *Putrificoraceae* with the following description.

Putrificoraceae that, though they sporulate, are exceedingly delicate and soon die in meat medium and other media. Their growth in meat medium resembles that of the organisms of genus *Martellillus*, they do not produce much gas, they color the meat particles a greyish color and form a blackish pigment. Gram-negative rods with central or subterminal cocoon-shaped spores. Parasitic forms which infect cattle in California and Nevada. I venture to place such organisms in a separate genus on account of their parasitic habit and on account

of their delicacy It may be that they are descended from organisms of the genus *Martellillus*

Type species *R. fragilis* nov sp Characters of genus

Reducibacteriaceae. A family of bacteria proposed by Orla-Jensen (1909, p 333) to include those genera of his order *Cephalotrichinae* which are spiral or bent in shape, which are active reducers, which do not contain sulphur or bacteriopurpurin, which are in filaments and which do not show a tendency to branching The characteristic reaction may be regarded as the reduction of nitrates to nitrites. The genera included are *Solidovibrio* and *Spirillum*

Regillus. A generic name proposed by Heller (1922, p 23) for a genus of the *Putrificoideae* with the following description

Putrificoideae that show to the eye little proteolytic action on meat They may or may not blacken it slightly, but the meat particles do not diminish in size Sugars are attacked Milk is digested Gram-positive rods which readily form oval subterminal or median spores Colonies minute, opaque, complex, yellow Typically toxic tissue invaders which produce in the guinea-pig a clear white oedema that does not rapidly lose its oedematous nature on section

Type species *R. progrediens* nov sp To be described in a future paper Two species, one found in a case of human gas gangrene (see Barney and Heller), the other in a sheep

Rhabdochromatium A genus of sulfur bacteria proposed by Winogradsky (1888, p 100) as differentiated from *Chromatium* by the elongated rod and spindle shape and through its variable growth forms He regarded it as possibly of subgeneric rank under *Chromatium* The type species, *Rhabdochromatium roseum* Win he regards as probably identical with *Rhabdomonas rosea* Cohn Two other species, *Rh minus* and *Rh fusiforme* were also described

De Toni and Trevisan (1889, p 942) include it as a subgenus of *Mantegazzaea* with the diagnosis "*Species chromogenae, baculis granula sulphuris foventibus*" Three species are included

Rhabdochromatium was recognized by Migula (1900, p 1048) with the description "Zellen frei, stab- und spindelförmig, zeitlebens schwarmfähig, mit Geisseln an den Polen" This description has also been used by Smith (1905, p 164)

Orla-Jensen (1909, p 334) makes this a synonym of *Rhabdomonas* This is probably the correct assignment of the genus

Rhabdomonas A generic name proposed by Cohn (1875, p 167) for the species *Rhabdomonas rosea*, with the following description

Ausser der *Monas Okenii* enthielt das rothe Wasser aus Kahla noch vereinzelte spindelformige, blass rosenrothe Korperchen, welche nach beiden Enden verjungt, in ausgewachsenem Zustande etwa 8 mal langer als breit sind ich bestimmte die Breite zu 3.8-5 μ (0.0038-0.005 mm) die Lange je nach dem Zustande der Theilung 20-30 μ . Die Vermehrung durch Quertheilung ist haufig zu beobachten die Theilhalften erreichen fast ihre normale Grosse, ehe sie sich trennen Charakteristisch sind auch fur diese Korperchen die dunkelen, stark Licht brechenden, in die rosafarbene Substanz eingelagerten Kornchen, die sich bald in grosserer, bald in geringerer Zahl vorfinden, auch wasserhelle vacuolen in der Mitte und an den Enden wurden beobachtet Die Bewegung ist langsam zitternd, abwechselnd vor und ruckwärts unter bestandiger Drehung um die Längsachse, ein wirbel am hinteren Ende deutet auf eine nachschleppende Flimmergeissel, wie bei *Monas Okenii*, die ich jedoch nur einmal wirklich unterscheiden konnte So viel ich weiss, ist diese rothe Spindelmonade noch nicht beschrieben sie gehort unter die Ehrenbergsche Section *Rhabdomonas* als eine neue Art, die ich als *Rhabdomonas rosea* bezeichnen will

Trevisan (1879, p 137) introduced the generic name *Mantegazzaea* (*q v*) to replace *Rhabdomonas* on the plea that the latter was "nomen ineptum" Winogradsky (1888, p 100) proposed still a third generic name *Rhabdochromatium* (*q v*) with two additional species

Rhabdomonas is included by most subsequent writers as a synonym of *Rhabdochromatium* It is so included by Migula (1900, p 1048). It is rejected as invalid by E. F. Smith (1905) Orla-Jensen (1909, p 334) suggests that it should be used to replace *Rhabdochromatium*

Buchanan (1918, p 471) included this genus as the first of the tribe *Chromatiaceae* with the following description

Differentiated from *Chromatium* by the elongated rod-shaped or spindle-shaped cells Cells red, with sulphur granules, polar flagella

The type species is *Rhabdomonas rosea* Cohn

Bergey (1923, p 402) followed Buchanan

Rhizobacterium A generic name used by Kirchner (1895, p 213) for the species of root nodule organism found in the nodules of *Lupinus*, *Ornithopus* and *Soja*, under the name *Rh japonicum*. He proposed the name because he regarded *Rhizobium* as invalid on account of the previous use in entomology of the generic name *Rhizobius* for a genus of aphids This reason is not valid, for both botanical and zoological codes recognize the validity of identical names in the two fields of science, though they are not recommended Apparently other writers

have not used this name It may be regarded as a synonym of *Rhizobium*

Rhizobium. A generic name proposed by Frank (1889, p 338) to include the organisms causing the formation of nodules on the roots of leguminous plants He expresses the belief that the organism probably belongs to the fission fungi or bacteria and states

Um in der vielleicht noch sich aufwerfenden Discussion, ober zu den Myxomyceten oder zu den Spaltpilzen, sei, in der Namengebung nichts zu präjudicieren, ist vielleicht die Bezeichnung *Rhizobium leguminosarum* für ihn empfehlenswerth

This generic name has been used by several authors, among them Ludwig (1892, p 99) Schneider (1892, p 203) and Hiltner and Stormer (1903, p 151)

The name is rejected by Erwin F Smith (1905, p 174) Orla-Jensen (1909, p 334) proposed the substitution of the name *Rhizomonas*

Winslow *et al* (Committee Soc Am Bact, 1917, p 553) make the following statement

Phytomyza Schroter has priority over *Rhizobium*, but because of the confusion which would arise from the substitution of the older correct name for the better known term *Rhizobium*, the committee recommends the adoption of the latter

Minute rods, motile when young by means of polar flagella Involution forms abundant and characteristic when grown under suitable conditions Obligate aerobes, capable of fixing atmospheric nitrogen when grown in the presence of carbohydrates in the absence of compounds of nitrogen Produce nodules upon the roots of leguminous plants

The type species is *Rhizobium leguminosarum* Frank

Buchanan (1918, p 46) gave the following description

Rod-shaped cells, motile when young by means of polar flagella, obligate aerobes, fixing atmospheric nitrogen when grown in a nitrogen-free medium containing suitable carbohydrates, involution forms abundant and characteristic, usually growing in the nodules of the roots of leguminous plants

The type species is *Rhizobium leguminosarum* Frank

The name of this genus has been a source of confusion The organism of leguminous nodules was placed in the mold genus *Schizzia* by Frank as *Schizzia leguminosarum* Schroeter (1886, p 134) concluded the organism to be one of the slime molds and created the genus *Phytomyza*, including it in the order *Phytomyzini* among the *Myxomycetes* He based his conclusions as to the position of this organism among the slime molds upon the work of Prilleaux (1879, p 98) Beijerinck (1888, p 758), named the organism *Bacillus radiculicola* Frank (1889, p 338) renamed the organism *Rhizobium leguminosarum* The fact that Schroeter included this genus incorrectly among the slime molds does not invalidate the

name The Committee on Classification of Bacteria, of the Society of American Bacteriologists, however, has recommended the use of the generic name *Rhizobium* as better known and probably resulting in less confusion than the use of *Phytomyra*

The genus is included by Bergey *et al* (1923, p 40) as the second in the tribe *Azotobacterae* with the following description

Minute rods, motile when young, branching forms abundant and characteristic when grown under suitable conditions Obligate aerobes, capable of fixing atmospheric nitrogen when grown in the presence of carbohydrates and in the absence of organic nitrogen compounds Produce nodules upon the roots of leguminous plants

Rhizobunae. The name of a subtribe proposed by Buchanan (1918 p 43) for the third subtribe of the tribe *Bacteriae* with the description "Rod-shaped organisms, securing their growth energy by the oxidation of carbonaceous compounds, as carbohydrates, alcohol, etc Do not require serum, etc Not parasitic in animals "

Three genera are included, *Mycoderma*, *Rhizobium* and *Azotobacter*

Rhizomonas A generic name proposed by Orla-Jensen (1909, p 334) as a substitute for *Rhizobium* (q v) to designate the organisms which are responsible for the development of nodules on the roots of leguminous plants It may be regarded as a synonym of *Rhizobium* and *Phytomyxa* (q v)

Rhodobacillus. A genus of bacteria named by Molisch (1907 p. 14) to include bacteria containing bacterio-purpurin and bacterio-chlorin but no free sulfur granules The organisms are short rods, rounded at the ends, usually solitary and motile One species is described, *Rhodobacillus palustris* Molisch

Meyer (1912, p 2) used *Rhodobacillus* as a designation for the second section of the genus *Bacillus* with the description "Sie sol die durch Bakteriopurpurin gefarbtten Spezies der Gattung *Bacillus* enthalten, sowohl die schwefelfreien wie die schwefelhaltigen Spezies "

Buchanan (1918, p 473) included this as the fifth genus of the tribe *Rhodobacterioidae*, as "Rod shaped cells, solitary, usually motile" with the type species designated as *Rhodobacillus palustris* Molisch

Bergey *et al* (1923, p 406) followed Buchanan

Rhodobacteria. An order of bacteria named by Molisch (1907, p 27) with the following description "Bakterien, deren Zellinhalt durch Bakteriopurpurin und (sowert untersucht) durch Bakteriochlorin rosa, rot, violett oder karminrot gefarbt sind " Two families are included, *Thiorhodaceae* and *Athiorhodaceae*

Rhodobacteriaceae. A family of bacteria created by Migula (1900, p 1042) to include all bacteria containing bacteriopurpurin, and in consequence colored rose, red or violet They also contain sulfur granules. This family is recognized by Smith (1905, p. 163), Orla-Jensen (1909, p 313), Benecke (1912), Frost (1911, p 60) and others. An ordinal name *Rhodobacteria* (*q v*) is used by Molisch (1907, p. 27)

Buchanan (1918, p 464) included this as the third family of the order *Thiobacteriales* with two subfamilies *Chromatioideae* and *Rhodobacterioideae* His description is "Cells of various types, not filamentous, containing bacterio-purpurin with or without sulphur granules also "

Bergey *et al* (1923, p 394) included this as the first family of the order *Thiobacteriales*

Rhodobacterioideae. A name given by Buchanan (1918, p 471) to the second subfamily of the *Rhodobacteriaceae* The name may be considered a synonym of *Athiorhodaceae* Molisch The diagnosis is. "Cells not filamentous, containing bacteriopurpurin but not granules of sulphur" The following genera were included *Rhodocystis*, *Rhodonostoc*, *Rhodosphaera*, *Rhodobacterium*, *Rhodobacillus*, *Rhodovibrio* and *Rhodospirillum*

Rhodobacterioideae. A variant spelling of *Rhodobacterioideae q v* used by Bergey *et al* (1923, p 404)

Rhodobacterium. A genus of bacteria first described by Molisch (1907, p 16) to include those bacteria containing bacteriopurpurin, devoid of sulphur granules, rod shaped, and non-motile A single species, (type) *Rhodobacterium capsulatum* Molisch is included

Buchanan (1918, p 472) included this as the fourth genus of the tribe *Rhodobacterioideae* Bergey *et al* (1923, p 406) followed Buchanan

Rhodocapsa. A genus of sulfur bacteria described by Molisch (1906, p 223) The cells contain granules of free sulfur, and are free that is, not united into families, not capable of swarming In mass the organisms are rose or cherry red, due to their content of bacteriopurpurin One species was described, *Rhodocapsa suspensa* (the type)

This name was also used by Orla-Jensen (1909, p 334) as the tenth genus of his family *Rhodobacteriaceae* It appears that he proposes to use this name to replace an older designation *Thiocapsa*. In this sense the name would be invalid

Buchanan (1918, p 471) includes this as the fourth genus of the tribe *Chromatieae* with the description "Cells spherical, free (not united

into families), not capable of swarming (non-motile) In mass the organisms are cherry red Contain sulphur granules "

Bergey *et al* (1923, p 404) follow Buchanan

Rhodococcus. This generic name has been introduced three times into bacteriological nomenclature It was first used by Zopf (1891, p 28) to apply to two species of red bacteria whose physiological characters had previously been studied at length by Overbeck (1891, p 399) These organisms had been known as *Micrococcus erythromyxa* and *M rhodochrous* He characterized the genus (or subgenus as he terms it) as follows

Colonien auf gewöhnlicher Nahrgeelatine gebirgsrückenartig, roth gefarbte Zellen weder ausgesprochen fädige noch flächenförmige oder körperliche Verbände bildend, sondern unregelmässig zusammengelagert, ohne Gallerthülle, einen rothen Fettfarbstoff enthaltend, der nach der Ausscheidung in rothen, auffälligen Aggregaten krystallisirt, und durch ein einziges breites Absorptionsband bei F ausgezeichnet ist

The species *Rhodococcus erythromyxa* was isolated by Zopf from tap water Overbeck describes it as producing round sharply circumscribed red colonies in plate cultures In stab cultures in agar and gelatin growth occurs only in the upper portion of the stab Grows well on potato, egg, etc, forming an abundant red pigment, cultures generally being slimy Cells are about 1.1μ in diameter, spherical In general they are grouped irregularly, but packets are sometimes formed in suitable media Gelatin is not liquefied The pigment was determined to be a lipochrome

These species are disposed of in various ways of subsequent authors Migula (1900, p 487) uses the designation *Bacterium erythromyxa* (Zopf) Mig Matzschita (1902, p 389) changes it to *Bacillus erythromyxa*, because of its shape He (p 441) however uses the name *Micrococcus rhodochrous* He describes it as

Grosse Zellen Kleine, mattglänzende, kreisrunde gewölbte, dunkelkarminrosa gefarbte Kolonien Auf Agar erst karminrosa, dann tief zinnrothe Auflagerung In Bouillon bildet er sich eine dicke rosa Haut, glatt, feucht und ein roter flockig-brockeliger Bodensatz

This shows that at least one of the species is a true coccus It would seem evident therefore if the red pigmented cocci are worthy of generic designation, that *Rhodococcus* Zopf is valid

Winslow and Rogers (1906, p 546) evidently without knowledge of the previous use of the name by Zopf, proposed *Rhodococcus* as a generic

designation for the red cocci. In their later publication (1908, p. 261) they give the following diagnosis:

Saprophytes. Cells in groups or regular packets. Generally decolorize by Gram. Growth on agar abundant, with formation of a red pigment. Dextrose broth slightly acid, lactose broth neutral. Gelatin rarely liquefied. Nitrates generally reduced to nitrites, but not to ammonia.

They recognize two species only, *Rhodococcus roseus* (Flügge) Winslow and *Rh. fulvus* (Cohn) Winslow. It is apparent that this description may well be regarded simply as an emendation of that of Zopf.

Rhodococcus was independently introduced as a bacterial genus by Molisch (1907, p. 20). The only species described is named *Rhodococcus capulatus* Molisch. This organism is described as belonging with the *Athiorhodaceae*, that group of the sulphur bacteria which contain bacterio-purpurin, but not free sulphur granules. This generic name is a homonym of *Rhodococcus* Zopf. This genus is recognized by Orla-Jensen (1909, p. 334) as the seventh in the family *Rhodobacteriaceae* as a synonym of *Thiopedia*. In this sense the name is invalid.

Meyer (1912, p. 4) included *Rhodococcus* as a section of the genus *Streptococcus* with the description "Alle durch Bacteriopurpurin gefärbten Streptococcusarten." He also includes a section by the same name under *Micrococcus* with the description "Mit Schwefelkörnchen."

Buchanan (1915, p. 12) discussed the synonymy and relationships of the genus.

Buchanan (1917, p. 614) included this genus as the second of the tribe *Micrococceae* with the description:

Saprophytes. Cells in groups or regular packets. Usually Gram-negative. Abundant growth with red pigment on surface of culture media. Slight acid from dextrose, none from lactose. Gelatin rarely liquefied. Nitrate usually reduced to nitrites, but not to ammonia.

The type species is probably *Rhodococcus roseus* (Flügge) Winslow.

This genus was also recognized by Winslow *et al.* (Committee Soc. Am. Bact., 1917, p. 560 and 1920, p. 208) with the designation of the type species as *Rhodococcus rhodochrous* Zopf. Castellani and Chalmers (1919, p. 931) state *Rhodococcus roseus* (Flügge) emendavit Dyar (1895) to be the type. Bergey *et al.* (1923, p. 75) follow the Committee.

It would appear that *Rhodococcus* Zopf, is a valid name to apply to the red cocci as a generic designation if they are to be grouped separately, and that the type is *Rhodococcus rhodochrous* Zopf.

Rhodocystis. A name used by Molisch (1907, p 22) to designate a genus of bacteria characterized by the presence of bacteriopurpurin, the absence of sulfur granules, the union of cells into families, the division of cells in only one direction of space, the cells rod-shaped, embedded in a common slimy capsule. One species *Rhodocystis gelatinosa* is described.

Orla-Jensen (1909, p. 334) proposed this name to replace *Thiocystis*. In this sense it is probably invalid.

Meyer (1912, p 2) included this genus in his section *Rhodobacillus* of the genus *Bacillus*.

The genus was recognized by Buchanan (1918, p 472) and the type species designated as *Rhodocystis gelatinosa* Molisch. The description reads "Cells rod-shaped, dividing in only one plane embedded in a common slimy capsule."

Rhododictyon. A generic name proposed by Orla-Jensen (1909, p 334) to replace the older name *Thiodictyon* Winogradsky, as the third genus of the family *Rhodobacteriaceae*. It is probably to be regarded as a synonym.

Rhodomonas. A generic name proposed by Orla-Jensen (1909, p 33) to replace the older name *Chromatium*. It is included as the first genus of the family *Rhodobacteriaceae*. It is probably an invalid synonym.

Rhodonostoc. A genus described by Molisch (1907, p 23) to include bacteria containing bacteriopurpurin without sulfur granules, with cells united into families, cells spherical or short rods, in chains like a rosary and embedded in a common gelatinous capsule. One species, *Rhodonostoc capsulatum* is included.

It is recognized by Buchanan (1918, p 472) with the designation of *Rhodonostoc capsulatum* Molisch as type, and the description "Cells spherical or short rods, in rosary like chains, and embedded in a common gelatinous capsule."

Bergey *et al* (1923, p 405) followed Buchanan.

Rhodopolycoccus. A generic name proposed by Orla-Jensen (1909, p 334) to replace the older name *Thiopolycoccus*. It is included as the sixth genus of his family *Rhodobacteriaceae*. It may be regarded as a synonym of *Thiopolycoccus*.

Rhodosarcina. A generic name used by Orla-Jensen (1909, p 334) to replace the older name *Thiosarcina*. It may be regarded as an invalid synonym of this latter term.

Meyer (1912, p 5) also used *Rhodosarcina* as a designation for a section of the genus *Sarcina* to include those species containing bacteriopurpurin. He includes here the genera *Thiocystis*, *Lamprocystis* and *Thiocapsa*.

Rhodosphaera. A name proposed by Buchanan (1918, p 472) to replace the homonym *Rhodococcus* Molisch (*qv*). The genus is included as the third in the tribe *Rhodobacterioideae*, with the description

Cells spherical, non-motile, free not united into families

The type species is *Rhodosphaera capsulatus* (Molisch) Buchanan

Bergey *et al* (1923, p 405) followed Buchanan.

The name is invalid as a generic designation for the bacteria, as the name *Rhodosphaera* was given by Engler in 1881 to a genus in the family *Anacardiaceae* among the flowering plants.

Rhodospira. A name proposed by Meyer (1912, p 4) for the first section of the genus *Microspira* with the description "Bakteriopurpurin enthaltende Spezies."

Rhodospirillum. A generic name used by Molisch (1907, p 24) to include bacteria containing bacteriopurpurin, without sulfur granules, with cells free, bent in a spiral, actively motile by means of a polar flagellum or cluster of flagella. Two species are described as new *Rhodospirillum giganteum* Molisch and *Rh. photometricum* Molisch, and Esmarch's species *Spirillum rubrum* is included as *Rh. rubrum* (Esmarch) Molisch.

This name was also used by Orla-Jensen (1909, p 334) to replace the older name *Thiospirillum* as the twelfth genus of his family *Rhodobacteriaceae*. This use of the name is invalid.

Meyer (1912, p 3) makes *Rhodospirillum* the second section of the genus *Spirillum* with the diagnosis "Spirillen welche Bakteriopurpurin in dem Protoplasten enthalten." He believes Molisch's genus *Rhodospirillum* should be thus disposed of.

Buchanan (1918, p 473) included the genus as the seventh in the subfamily *Rhodobacterioideae* with the description "Cells spiral, actively motile by means of polar flagella." The type species is designated as *Rhodospirillum rubrum* (Esmarch) Molisch.

Bergey *et al* (1923, p 407) followed Buchanan.

Rhodothece. A genus of bacteria named by Molisch (1906, p 223) to include organisms containing bacteriopurpurin and sulfur granules, cells not united into families, not capable of swarming, each cell surrounded by a spherical or ellipsoidal capsule, cell usually spherical.

and occurring in pairs One species, *Rhodothece pendens* Molisch is described

This name was also introduced by Orla-Jensen (1909, p 334) to replace the older name *Thiothece*, as the fifth genus of the family *Rhodobacteriaceae*. In this sense the name is probably invalid

Buchanan (1918, p 471) included this as the fifth genus of the tribe *Chromatiaceae* with the description

Cells usually spherical and in pairs, each surrounded by a spherical or an ellipsoidal capsule Non-motile Cells not united into families Cells contain bacteriopurpurin and sulphur granules

The type species is *Rhodothece pendens* Molisch

Bergey *et al* (1923, p 404) followed Buchanan

Rhodovibrio. A generic name used by Molisch (1907, p 21) to include those bacteria containing bacterio-purpurin, without sulfur granules, with cells short, comma shaped, actively motile by means of a single terminal flagellum One species *Rhodovibrio parvus* Molisch is described

Buchanan (1907, p 21) included this as the sixth genus of the sub-family *Rhodobacterioidae* with the description

Cells short, comma shaped, free, actively motile by means of a single terminal flagellum

The type species is *Rhodovibrio parvus* Molisch

Bergey *et al* (1923, p 407) followed Buchanan

Rickettsia. A generic name proposed by Da Rocha Lima (1916, p 567) for the organism causing typhus fever The description was elaborated by Arkwright, Bacot and Duncan (1919, p 76) Enlows (1920, p 79) gives the following summary:

Very small, 0.3 to 0.5 μ by 1.5 to 2 μ Morphologically like a coccus, diplococcus or a short bacillus Gram-negative Not acid fast Stains well by Giemsa, appearing as small dots, paired cocci or bipolar staining bacilli with an unstained central part Not motile Occurs sparsely in blood films Not successfully grown on artificial media

Prowazek regards this organism as a protozoon largely because it is insect borne, and Da Rocha Lima seems rather inclined to this view on account of its peculiar staining reactions Arkwright, Bacot, and Duncan, however, regarded its Giemsa staining reactions as quite like those of other bacteria They conclude "Nevertheless this class of microorganism and its associated diseases appear to have sufficiently distinct characteristics to justify the retention of the name *Rickettsia* for the present" They state that Da Rocha Lima found these bodies in the midgut of lice (*Pediculus humanus*) fed on trench fever patients,

and that he considered the species causing typhus and trench fevers distinct. In typhus he claimed that the organism (*Rickettsia prowazekii*) invaded the epithelial cells of the gut wall, while only rarely is this the case with the trench fever organism (*R. quintana*), and with *R. pediculi* which is found in normal lice. He also claims that morphological differences are easily discernible if serial sections are cut. He infected a few guinea pigs, but was not able to pass the disease on from pig to pig, nor to infect mice.

The type species is *Rickettsia prowazekii* Lima.

Rivoltellus. A name proposed by Heller (1922, p. 20) for the twelfth genus of *Clostridiaceae* with the following description.

Clostridiaceae possessing moderately strong saccharolytic powers. Liquefy gelatin but do not produce H_2S demonstrable by a lead-acetate-paper test in blood broth. Produce in meat medium gas and a pink coloration which does not rapidly fade. Clot milk. Do not liquefy serum or egg or disintegrate meat particles. Gram-positive rods, usually short, with median, sub-terminal, or terminal spores, which usually bulge the sides of the bacillus. Sporangia not often much larger than vegetative rods. In tissue the sporangia may be uneven in their staining reactions, "granulose" being present, oocysts are long, frequently with parallel sides. Usually form chains on the liver of animals. Colonies in deep agar, though they may start as lenticular structures, consist later of a dense center and a wide loose woolly periphery. They vary in size, etc., according to species. Typically highly pathogenic tissue invaders that produce haemolysis in the animal body. Pathogenic for a wide range of species.

Type species *R. vibron* (the vibron septique of Pasteur), as defined in a future paper. Robertson (1920) has divided the group into four sub-groups on the basis of the agglutination reaction.

Robertsonillus. A name proposed by Heller (1922, p. 24) for a genus of the *Putrificaceae* with the following description.

Putrificaceae that produce on meat medium a little gas and a terra cotta coloration, multiplying slowly for a long period. A black pigment is absent. Sugars not fermented. Weakly Gram-positive rods with oval spores, usually subterminal, that somewhat distend the bacillus. Two spores are produced. One species, found twice in moulds. Non-pathogenic for guinea-pigs in pure culture.

Type species *R. primus* (Bacillus I, Hempl) as described by Hempl. Descriptions of bacilli similar to this have not been noted. Proteolytic organisms producing a terra cotta coloration in meat medium are not uncommon, however.

Saccharobacillus. A species of organism named *Saccharobacillus pastorianus* by Van Laer (1889) was found by this author to be the cause of the disease of beer called "turning," in which the beer loses its brilliancy, becomes disagreeable in smell and taste and forms a sediment. According to Henneberg (1909, p. 480) the colonies on wort

agar are smooth, gray-white, with flocculent sediment in the center. In the stab the growth is better than on the surface. Mash becomes clouded in two to three days with gas formation, and becomes clear after five to fourteen days. In wort flocculent masses form on the walls, when shaken it is clouded. In liquids the organism consists of slender, long rods which usually are united in 2, 3 or more straight, bent or wavy lines. Maximum growth temperature is about 37°, optimum 29 to 33° and minimum 11°C. Lactic acid to amount of 1.5 per cent is produced.

The following sugars are acidified: arabinose, trehalose, dextrose, levulose, galactose, maltose, methyl glucosid, raffinose, mannite, dextrin, cane sugar. Starch and lactose are slightly fermented. Xylose, rhamnose, inulin, erythrit, quercit and dulcitol are not fermented.

This generic name is rejected by Erwin F. Smith (1905, p. 174).

The genus *Saccharobacillus* has not come into general use, and is probably to be regarded as a synonym of *Lactobacillus*.

Saccharobacter. A generic name used by Beijerinck (1900, p. 200) for organisms closely related to his genus *Aerobacter* (q.v.). The types mentioned are those generally termed *Bacillus megatherium* and *B. hortulensis*. The name is listed as invalid by E. F. Smith (1905, p. 174).

Salmonella. A generic name used by Lignieres (1900, p. 389) and others for the organisms of the hog cholera bacillus or intermediate group of the intestinal bacteria. Lignieres states:

Le microbe de la Schweineseptikemie rentre dans le groupe des *Pasteurella*, celui du Hog-cholera ou schweinepest est très différent, il pourrait servir de prototype pour la création d'un autre groupe, celui des *Salmonella*.

Again (1901, p. 734) he states: "J'ai, de plus, jeté les bases d'un autre groupe, celui des salmonelloses, ayant pour type le microbe du hog-choléra de Salmon."

Buchanan (1918, p. 53) used this name as a designation of his second subgenus of the genus *Bacterium* with the definition: "Fermenting glucose but not lactose with formation of acid and gas. The type species is *Bacterium (Salmonella) cholerae suis*?"

Castellani and Chalmers (1919, p. 934) recognize this as the sixth genus of the tribe *Ebertheae*. The definition is:

Ebertheae which completely ferment glucose, but do not ferment lactose, and partially or completely ferment mannitol, in addition to other carbohydrates. Milk not clotted.

Type species: *Salmonella paratyphi* (Schottmüller, 1902).

Apparently Castellani and Chalmers are in error in fixing the type species Lignières included only the organism associated with hog cholera in the group or genus It remains to determine just what specific name should be applied

Bergey *et al* (1923, p 212) state

Motile forms occurring in the intestinal canal of animals, in various types of acute, inflammatory conditions Attack numerous carbohydrates with the formation of both acid and gas In general do not form acetyl-methyl-carbinol

The type species is *Salmonella schotmulleri*

Saprospira. A genus of free living spirochetes named by Gross (1911, p 190) with the following diagnosis. "Spirohemacea mit spiralig gebogenem Körper Vermehrung durch Zerfallstheilung, Freilebend The species, *Saprospira grandis* (100 μ in length) and *S. nana* (36 μ in length) were described Later (1912, p 84) he says "Die Saprospiren (Fig 2) gleichen den Cristispiiren in hohem Masse, nur fehlt ihnen die Crista Es erscheint daher geboten, für sie ein eigenes Genus zu schaffen "

Winslow *et al* (Committee Soc of American Bacteriologists, 1917, p 563) state.

Non-parasitic forms similar to *Cristispira*, but without the flattened ridge or "crista" which is, if present, here replaced by a straight columella or thickening of the periplast

The type species is *Saprospira grandis* Gross

Buchanan (1918, p 544) included this as the second genus of the family *Spirochaetaceae*

Slender spiral cells living free in salt water, actively motile, flexuous Cross section circular

The type species is *Saprospira grandis* Gross.

The genus is also recognized by Castellani and Chalmers (1919, p 439) and by Bergey *et al* (1923, p 422)

Sarcina A genus created by Goodsir (1842, p 432) to include the species *Sarcina ventriculi* which he discovered in microscopical examination of vomit His publication created a good deal of interest and many papers were published in the next quarter century upon this "Sarcinosis" The organism was apparently found repeatedly under similar conditions This form is of interest because it is one of the first organisms which we now include among the bacteria to be

described definitely as a plant Enlows (1920, p 80) has summarized the characteristics of the species as follows

Frustules 16, color light brown, transparent matrix very perceptible between the frustules, less so around the edges, size "0 800 to 0 1000 of an inch" "The individual organisms were transparent and slightly yellow or brown When carefully examined under favorable circumstances the cell walls appeared rigid, and could be perceived passing from one flat surface to the other as dissepiments These dissepiments, as well as the transparent spaces, were from compression of contiguity rectilinear, and all the angles right angles, but the bounding cells bulged somewhat irregularly on the edges of the organism by reason of the freedom of pressure These circumstances gave the whole organism the appearance of a wool pack, or of a soft bundle bound with cord, crossing it four times at right angles, and at equal distances" Found in the human stomach

Robin (1847, p 47) characterized the genus as follows

Plante coriace, transparente, consistant en masses cubiques ou allongées, composees de 16 à 64 cellules cubiques, partagées en 4 frustules, arrangées l'une à côté de l'autre Espèce *S. ventriculi* (Goodsir) Frustules 16, de couleur brune claire Masse transparente, laissant voir son intérieur entre les frustules Les bords de celles sont emousses Diametre 1/120 a 1/100 de ligne Habite l'estomac humain

The first definite inclusion of the genus in a scheme of classification was the assignment of this form to a position in the tribe *Glaeogenae* by Cohn (1875) The genus was also discussed by Luerissen (1879, p 21) who mentions the species *S. ventriculi* Goodsir He states that the spherical cells divide in three directions of space and form cubical families Magnin (1880, p 96) states that "*Sarcina* is very nearly allied to *Merismopedia*, from which it only differs by the absence of chlorophyll, its siliceous skeleton allies it with the diatoms" He mentions no species

Trevisan (1879, p 145) accepted the genus among the bacteria under the subtribe *Sarcinae*, with the following definition

Cellulae globosae, divisionis tempore ovoideae, quaternatim in colonias tabulatas consociatae, tegumentis propriis gelatinosis crassiusculis confluentibus obvolutae Coloniae tegumento communi destitutae

Two species are listed, the *Sarcina ventriculi* Goodsir and *Sarcina urnae* Welcher

Before 1885 the genus is discussed by at least three other writers, Van Tieghem (1884, p 1114), Grove (1884, p 20) and Winter (1880, p 48) Grove gives the definition of Winter, stating

Cells roundish, dividing in two or three dimensions of space Daughter cells connected for some time, forming small solid families or plates which are often again in their turn united to form larger colonies Families usually consisting of four or a multiple of four cells

He is thus enabled to bring into the genus some four other organisms which had previously been termed *Merismopedia* Zopf (1885, p. 51) limited the genus to those forms in which cell division occurs regularly in three planes with formation of packets Falkenheim (1885, p. 229) made use of the newly developed gelatin plate methods to secure cultures of *Sarcina ventriculi* Goodsir He found that this species would grow readily upon artificial media, forming light yellow, round colonies in from thirty-six to forty-eight hours Packets characteristic of *Sarcina* were missing in all media tried, except hay infusion In this medium the packets were well developed A study of this paper shows how extensive a literature had grown up concerning *Sarcina* He notes that twenty years earlier (1865) fifty titles were already known It would seem that the organism described by Falkenheim might well be taken as the type of the genus

The genus as defined by Zopf was recognized by many writers up to 1895, among them Schroter (1886, p. 153, *S. ventriculi* and 3 new species), Flugge (1886), De Toni and Trevisan (1889, p. 1044, 15 species), Cornil and Babes (1890, p. 148), Billet (1890, p. 24), Hueppe (1885, 1891, p. 30), Ludwig (1892) and Sternberg (1892, p. 17)

Migula (1894, p. 236) says

Sarcina Goodsir Die Zellen teilen sich nach 3 Richtungen des Raumes, wodurch, wenn sie nach der Teilung verbunden bleiben, warenballenartig eingeschnurte Packets entstehen können Bewegungsorgane fehlen (Z. B. *Sarcina ventriculi* Goodsir, *S. rosea* Schroter)

Migula (1895, p. 18) defined the genus somewhat more accurately insisting that the outstanding generic character is division in three planes and not necessarily the formation of the packets He also separated the motile forms into the new genus *Planosarcina* He notes that about 45 species are known Gruber (1895, p. 239) monographed the genus He gives the following generic description

Zellen rund, einzeln oder zu warenball artig eingeschnurten packetformiger Verbänden vereinigt, oder zu 2 (Diplococcenform) oder 4 einander genähert Teilung nach drei Richtungen des Raumes Bleiben die Tochterzellen nach der Teilung vereinigt, so sind sie an den Berührungsflächen mehr oder weniger abgeplattet, trennen sie sich nach der Teilung so sind sie völlig kugelförmig

He recognizes in all 39 species With Migula's diagnosis the genus has been used by Lehmann and Neumann (1896, p 135), Chester (1897, p 62), Migula (1897, p 27, 1900, p 18 and 1904, p 144), Macé (1897, p 335), Fischer (1897, p 32), Hewlett (1898), Baumgarten (1900, p 127), Chester (1901, p 109), Lehmann and Neumann (1901), Schmidt and Weis (1902, p 288), Kendall (1902), Matzuschita (1902), A J Smith (1902, p 270), Fischer (1903, p 59), Klocke (1903, p 331), E F Smith (1905, p 160), Fluegge (1908), Ellis (1909, p 4), Orla-Jensen (1909, p 340), Heim (1911, p 249), Frost (1911, p 57), Schneider (1912, p 22), Lohnis (1913, p 45), and Vuillemin (1913, p 519)

The Winslows (1905 p 669) have proposed a somewhat radical emendation of the genus In its latest form (1908, p 260) it reads

Facultative parasites or saprophytes Division occurs under favorable conditions in three planes, producing regular packets Generally decolorize by Gram Growth on agar abundant, with formation of yellow pigment Dextrose broth slightly acid, lactose broth generally neutral Gelatin frequently liquefied Nitrates may or may not be reduced

They recognize three species *Sarcina lutea*, *S citrea* and *S flava* To these Kligler (1913, p 432) has added *Sarcina aurantiaca*, an orange organism, emending therefore the generic description by the insertion of "and orange" after "yellow" The Winslows (1 c, p 236) in discussion of *Sarcina ventriculi* state

Another interesting *Sarcina*, possibly related to *S lutea*, is the form originally isolated by Goodsir in 1842 and named by him *S ventriculi* It was described as a non-liquefying sarcina, isolated from the stomach in cases of hyperacidity of the gastric juice More recent investigations suggest that there is nothing specific in the relation of this organism to the pathological condition in question (Flügge, 1896) *S ventriculi* was distinguished from the type of *S lutea* by the production of an orange, instead of a yellow, pigment It corresponds therefore to the non-liquefying *S aurantiaca* Whether Goodsir's form was a packet-forming *Aurococcus* or an orange *Sarcina* can only be decided from a study of similar form which the future may bring to notice

The writer has been unable to find the authority for terming *S ventriculi* orange, Falkensheim (1885), Flügge (1886), Eisenberg (1891) and Gruber (1897, p 259) all state that this organism is yellow on culture media, except that the latter states that on potato "Später orangegelb werdende Colonien" It would appear therefore that this name, *Sarcina ventriculi* should have been given to one of their species,

and constitute the type of the genus Beijerinck (1906, p 200) identified *Sarcina ventriculi* Vuillemin (1913, p 525) proposes that this genus should be a "*Genus conservandum*."

Buchanan (1915, p 10) discussed in detail the validity of the generic name and concluded the type should be *Sarcina ventriculi* Goodsir. Later (1917, p 614) he included it as the third genus of the tribe *Micrococceae* with the following description

Saprophytes or facultative parasites Division occurs under favorable conditions in three planes, producing regular packets Usually Gram-negative. Growth on agar abundant with formation of yellow pigment Dextrose broth slightly acid, lactose broth generally neutral Gelatin frequently liquefied. Nitrates may or may not be reduced

The same disposition and practically the same diagnoses have been used by Winslow *et al* (Committee Soc Am Bact, 1917, p 559, 1920, p 208), Castellani and Chalmers (1919, p 925) and Bergey, *et al* (1923, p 70)

Sarcinacoccus According to Enlows (1920, p 81) this casual name was used by Billroth (1874, p 8) for one of the growth forms of his *Coccobacteria septica*

Sarcinaglobulus A generic name used by Poulsen, (1879, p 232). Enlows (1920, p 81) gives the following as the description used

Schizophytarum corpora perpava hyalina e cellulis incoloribus minimis composita formans Cellulae plantae vivae vix conspicuae reagentibus chemiciis additis apparent Divisionis modus ut in Sarcina, quacum maximam similitudinem habet Ab hoc genere eo differt, quod non fasciculos hexaedricos sed globulos vel flebas subisodiametricas vel irregulares rotunditas format Nucleum cellularum non observati Species adhuc una cognita, scil

Type species (monotypy) *S punctum* Char gen Magn 2 to 200 μ In limo putido litoris freti cresund prope Haunias legi

The position and status of this genus is uncertain

Sarcinastrum. A generic name proposed by Lagerheim (1900, p. 280) for an organism *Sarcinastrum Urosporae* found producing galls upon the seaweed *Urospora mirabilis* The following description of the organism is given

Die zu Stäbchen ausgewachsenen Coccen erzeugen wahrscheinlich durch Langspaltung Kolonien, bestehend aus parallelen Stäben, die zu einer runden, gebogenen Scheibe angeordnet sind Diese Scheibe wird später durch fortgesetzte Langsteilung der Stäbe halbkugelig, zuletzt fast hohlkugelig Nachdem die Kolonie eine gewisse Grosse erreicht hat, tritt Quertheilung der Stäbe zu Coccen ein, die Kolonie besteht schliesslich aus einer sehr grossen Anzahl

sehr kleiner, reihenweise angeordneter, kugelliger Zellen. Die gallertartige Substanz, die die Zellen zusammenhält und die ganze Kolonie umgibt, löst sich darauf allmählich auf, so dass die winzigen Coccen von einander frei werden und durch Wasserströmungen etc. fortgetrieben werden können. Eine Eigenbewegung der Coccen wurde nicht beobachtet. Durch die Schleimschicht können dieselben der *Urospora-Membran* angeklebt werden und sich weiter entwickeln.

Der Parasit ist von dem in mehreren Hinsichten ähnlichen *Sarcinoglobulus punctum* Pouls stofflich verschieden. Jodjodkalium und Anilinfarben bewirken bei jenem eine schwächere Färbung, als bei diesem.

The genus is probably valid, although the location of this organism among the bacteria might be questioned.

The type species is *Sarcinastrum urospora* Lagerheim.

Sarcineae. A subtribal name under *Bacteriae* used by Trevisan (1879, p. 136) with the following description: "Cellulae substantia intercellulari in colonias mucosas determinatas consociatae." The genera included were *Sarcina*, *Chlamydatomus* and *Ascococcus*. De Toni and Trevisan (1889, p. 1044) also used the name as a tribal designation with the following description: "Cocci in familias pluristratas vel unistratas, muco matricali plus minus conspicuo involutas, consociati. Cystides nullae. Endosporae microsomae in coccis normalibus obvenientes."

The following genera are included: *Thiopolycoccus*, *Sarcina*, *Lampropedia*, *Thiocapsa*, *Pediococcus*.

Sarcinee. A tribal designation used by Trevisan (1889, p. 27) for the second tribe of the suborder *Coccogene*. Four genera are included: *Thiopolycoccus*, *Sarcina*, *Pediococcus* and *Thiocapsa*. The description is: "Cocchi più o meno densamente aggregati senz'ordine, o ad 8 ad 8, o a 4 a 4, in famiglie pluristrate od unistrate, più o meno avvolte in ganga gelatinosa, non mai racchiuse in cisti speciali." It is a variant of *Sarcineae*.

Sarcinoglobulus. See *Sarcinaglobulus*. The original spelling has not been determined, as the article is not available. Enlows (1920, p. 81) gives *Sarcinaglobulus* while Lagerheim (1900, p. 280) and others (see Journ. Roy. Mic. Soc., 1880, p. 1001) write *Sarcinoglobulus*.

Schinzia. According to Schroter (1886, p. 134) this generic name was first used by Dennstatt. He says: "Eine Gattung *Schinzia* ist schon früher 1818 im Dennstatt aufgestellt worden." The same name in another sense was used by Naegeli (1842, p. 279) as the generic designation of a fungus found in the rhizome of the Iris. Frank (1879, p. 376) concluded the organism responsible for the root nodules of leguminous plants belonged to the genus defined by Naegeli. To it

he gave the name *Schinzia leguminosarum*. When later it was found that these organisms are bacteria, and not fungi, they were removed from this genus. *Schinzia* is not tenable as a genus name among bacteria.

Schizomycetacea. The class name of the bacteria as written by Castellani and Chalmers (1919, p. 924) with the description

Thallophyta without chlorophyll and as a rule without chromatophores, with the vegetative body consisting of a single cell, in which the nucleus is not present in the form typical for other thallophytes. Reproduction by fission or spore formation.

Three orders are recognized, *Eubacterales*, *Thiobacterales* and *Myxobacterales*.

Schizomycetaceae. De Toni and Trevisan (1889, p. 923) were apparently the first to use this name as a family designation for the entire group of bacteria. They give the following description.

Fungi mycelio proprio carentes, typice unicellulares, minutissimi, simplicissimi (globosi, elliptici, lineares, filiformes, recti v. incurvi v. spirales), vegetatione apicali genuina destituti, rarissime pseudoramosi. Multiplicatio vegetativa fit cellularum divisione in unam v. duas vel tres directiones repetita, individua in statu vegetativo immobilia vel vivide oscillantia, saepe etiam in colonias vel alio modo congregata. Sporarum formatio duplex: tum plasmatis homogenei pars a liqua vel cellulae plasma totum contrahitur, condensatur et membrana peculiari induta in sporam transmutatur (ENDOSPORAE), tum cellula tota, sine plasmatis contractione, in sporam transformatur (ARTHROSPORAE).

OBS. Fungilli minimi sed in oeconomia universali maximi penderis, sunt enim pathogeni, zymogeni vel saprogeni.

These authors subdivide the family into the following tribes, *Trichogengae*, *Baculogengae* and *Coccogengae*.

Schizomycetes. A name first used by Naegeli (1857, p. 760) as a general designation of the forms we now recognize as bacteria. He included the organism causing silk worm disease, *Nosema bombycis*, also *Umbina acetii* (mother of vinegar), *Bacterium*, *Vibrio*, *Spirillum*, *Hygrococcus* and *Sarcina*.

This designation for the group of the bacteria has been quite generally accepted by bacteriologists.

Cohn (1872a, p. 236) used term "Klasse der Schizomyceten."

Cohn (1872a, p. 237) uses term "Familie der Schizomyceten."

Thielsen-Dyer (1875, p. 325) used *Schizomycetes*, as did Lankester (1876, p. 37), Grove, (1884, p. 1), Balbiani (1886, p. 543).

The name has frequently been used in the form *Schizomyceten*, *Schizomycetacea* (g v) and *Schizomycetaceae*

Hansgirg (1888, p 229) gives this as a synonym of Bacteria

Clements (1909, p 8) recognizes this as a class name with the two orders *Bacteriales* and *Myxobactrales*, and the description

Typically one-celled fungi, dividing by fission in 1, 2 or 3 planes, sometimes forming true filaments, but then motile or sheathed, and without true branches, resting cells often developed, sexual reproduction lacking

Vuillemin (1913, p 512) gave an excellent characterization of the group He states that they are simple organisms, formed of a single element without septa and unbranched The element is circumscribed by a rigid vegetable like membrane, elastic, but not contractile, sometimes also with a capsule It may undergo plasmolysis or plasmoptysis The protoplasm is less differentiated than that of most cells, the chromatin particles do not form an individual nucleus of a permanent type Division is amitotic Some forms are motile with flagella which traverse the membrane at points characteristic of the species (polar or diffuse) They are not broader than 5μ when not in bunches or unimpregnated with metal or colloidal coloring material The resting stage may be either an arthrospore resulting from a simple modification of the membrane or an endospore In some cases the spore bearing element retains its form or is modified passively by the enlargement of the spore, in other cases it is a specialized element for spore production Only the latter type of sporulation is satisfactory for generic characters The amitotic division usually occurs by a pinching transversely with rapid separation of two individuals These may remain united into families, in chains, layers or packets as determined by the successive planes of division He emphasizes that all forms showing contractility should be placed in the protozoa, that the myxobacteria constitute a distinct group, and that forms like the tubercle bacillus which show branching should be placed in the mold group *Microsiphones*

Buchanan (1917, p 155) used the following characterization

Typically unicellular plants, cells usually small and relatively primitive in organization The cells are of many shapes, spherical, cylindrical, spiral or filamentous, cells often united into groups, families or filaments, occasionally in the latter showing some differentiation among the cells, simulating the organization seen in some of the blue-green, filamentous algae No sexual reproduction known Multiplication typically by cell fission Endospores are formed by some species of the *Eubacteriales* (see below), gonidia (conidia, arthrospores) by some of the filamentous forms Chlorophyll is produced by none of the bacteria (with

the possible exception of a single genus) Many forms produce pigments of other types The cells may be motile by means of flagella, some of the forms intergrading with the protozoa are flexous, a few filamentous forms (as *Beggiatoa*) show an oscillating movement similar to that of certain of the blue green algae (as *Oscillatoria*)

He recognized six orders, the *Eubacteriales*, *Chlamydobacteriales*, *Thiobacteriales*, *Actinomycetales*, *Myxobacteriales*, and *Spirochaetales*

Winslow *et al* (Committee Soc Am Bact, 1917, p 549) used the following description

Minute, one-celled, chlorophyll-free, colorless, rarely violet-red or green-colored plants, which typically multiply by dividing in one, two or three directions of space, the cells thus formed sometimes remaining united into filamentous, flat, or cubical aggregates Capsule or sheath composed in the main of protein matter The cell plasma generally homogeneous without a nucleus Sexual reproduction absent In many species resting bodies are produced, either endospores or gonidia Cells may be motile by means of flagella

The same definition is used by Bergey *et al.* (1923, p 29)

Schmidlea. A generic name proposed by Lauterborn (1913 p 98) for one of the family *Chlorobacteriaceae* His diagnosis is

Schmidlea nov gen mit *Sch luteola* (*Aphanothece luteola* Schmidle) Zellen elliptisch, 0,0015–0,002 mm lang, zu wolkenartigen, rundlichen bis ovalen, oft gelappten gallertigen Kolonien vereint, die bisweilen vakuolenartige Raume umschliessen Die Kolonien umfassen oft Tausende von Zellen und erreichen nicht selten mehr als 0,2 bis 0,3 mm Durchmesser

The organism is found in decomposing plant remains in lake bottoms, particularly in decomposing *Chara*

The position of this genus among the bacteria is questionable

Schuetzia. A genus proposed by Trevisan (1889, p 29) as the second genus of his tribe *Streptococceae* The description is 'Cocchi globulosi o per incipiente temnogenesi brevemente ellissoidi (monococchi), seriatim in filamentis moniliformi, simplici, racchiusi in tenui capsula membranaceo-gelatinosa Spore macrosome provenienti nei filamenti' Three species are included, *Sch Poelsii*, the organism of equine distemper, *Sch Laughlini* and *Sch Dissel* The diagnosis given by De Toni and Trevisan (1889 p 1052) follows

Cocci globosi vel divisim tempore ovoidei, in filamenta moniliformiter concatenati, capsulis membranaceo-gelatinosis, arctic, tenuiusculus, homogeneis, non lamellosis obducta Arthrospora macrosomae, in filamentis obvenientes

Five species are included, one *Schuetzia equi* (Sand et Jensen) Trevisan, is the organism commonly known as *Streptococcus equi*, the cause of equine "strangles"

The first species listed by De Toni and Trevisan (1889, p 1052) is *Schuetzia Lagerheimii* (Ludwig) Trevisan. Probably *Sch equi* may be considered the type Its exact nomenclatural status is uncertain

Sclerothrix This name was apparently first used by Kuetzing (Ag Aq dulec Dec 11) for one of the blue green algae under the designation *Sclerothrix Callitrichae* Later Kuetzing (1849, p 319) gave the following generic description "Trichomata vaginata subramosa radiatim congesta, aequilonga, in cespitem globosum coalita (Parasiticae palustres)" This author had previously described this alga (1843, p 229) as *Hyphaethrix Callitrichae*

Metschnikoff (1888, p 70) proposed this as a generic name for organisms resembling the tubercle bacillus He states that this organism in its perfect state may grow into short threads, and also differs from other forms except the leprosy bacillus, in the possession of a dense envelop The name *Sclerothrix Kochii* was proposed for the organism causing tuberculosis

This genus was regarded as a synonym of *Bacillus* by De Toni and Trevisan (1889) The name is rejected by Lehmann and Neumann in the following statement

Since we proposed this name (*Mycobacterium*) in the first edition, we have seen that Metschnikoff (Virchow's Archiv, 113, p 70, 1888), who first recognized the peculiar position of the tubercle bacillus as opposed to the other then known bacteria, in a work "Regarding the Phagocytic Rôle of the Tubercular Giant Cell," has said "If one considers that in the perfected stage the tubercle bacteria have grown into (although short) threads, and also differ from other analogous forms (except the lepra bacteria) in the possession of a very dense envelope, then perhaps it will be easier to accept the designation *Sclerothrix* for the genus, and *Sclerothrix Kochii* for the species of the tubercle bacterium" We should have immediately accepted these names if we had known of them, but believe that according to the rules of botanical nomenclature our names should now stand, since Metschnikoff only made a conditional proposal, did not accurately define his new genus, and never made any use of the new name himself, while we have ourselves already established a name

This is also rejected by Erwin F Smith (1905, p 174)

Sclerothrix has been proposed by Vuillemin (1913, p 527) as a "Genus conservandum of the Microsiphones with *Sclerothrix tuberculosis* Metschnikoff as the type"

If the organisms of the type of the tubercle bacillus are to be given generic recognition, *Sclerothrix* can be used as a generic name only if it can be shown that the previous use by Kuetzing does not invalidate it, and that *Coccothrix* (Lutz (*qv*)) is invalid, for the latter term has priority.

Seguinillus. A name proposed by Heller (1922, p. 23) for a genus of *Putrificoraceae* with the following description

Putrificoraceae that do not exert a marked action on protein. Soften and may blacken meat. Milk clotted, may later be digested. Attack a few sugars. Very reluctant to form spores. Minute Gram-positive or Gram-negative rods, uniform in thickness but not in length. Spores oval, sub-terminal. Deep colonies lenticular or modified lenticular. Occasional tissue invaders which produce a proteolytic gangrene or phlegmon.

Type species *S. aerofoetidus* (*Bacillus aerofoetidus* Weinberg and Séguin) as described by Weinberg and Séguin (p. 161). The organism forms oval subterminal spores.

Semiclostridium. A genus of bacteria proposed by Maassen (1905, p. 5) with *Semiclostridium commune* as the type. This organism is aerobic bacillus which form spores, the sporangia being of a type which would place it in Fischer's *Clostridiaceae*. It has an unusual growth temperature range, from 18° to 55°. Its optimum is about 45°, it is a facultative thermophile. The organism is motile, with peritrichic flagella. It peptonizes milk. In sugar solutions it forms larger gelatinous zoogloal masses. Other species described are *S. citreum*, *S. flavum* and *S. rubrum*.

This is probably to be regarded as a synonym of *Bacillus*.

Serratia. The original article in which the name was proposed by Bizio (1823, p. 288) for fungus genus is apparently not available in America. However, the original report is reviewed by Bizio in a communication to the Paris Academy of Sciences (1844, p. 951). He states that in July 1819 in the province of Padua there was observed a phenomenon of reddening of the surface of "polenta," a porridge made from corn (maize) meal. In the course of eight to ten hours this was covered by a substance having the color of blood. It also appeared on bread and on rice kept under the same conditions. The phenomenon excited so much surprise and astonishment among the natives that it came to the attention of the public authorities and a commission of professors from the University of Padua was appointed to investigate. Bizio thought the phenomenon might be due to some fungus. He placed some polenta in a saturated atmosphere at 21° Reaumur. In

eight hours purplish spots were observable, and within twenty-four hours the entire surface was covered. The results of this experiment were published in *Gazetta privilegiata*, Venice, Aug. 24, 1819. He observed further that a bit of the reddened polenta placed in contact with freshly prepared polenta caused the latter to redden more quickly. He concluded it to be due to the "seeds" of some microscopic fungus present in the air.

He says

J'examinaï aussi les taches purpurines de la polenta de maïs avec le microscope, et, j'aperçus un agrégat de très-petites vésicules hémisphériques, couvertes d'une pellicule mince, un peu luisante, et parsemée de petits points d'une couleur pourpre si foncée, qu'elle paraissait presque noire. J'ai jugé que des points étaient les sporules ou étaient placées les semences qui reproduisent l'espèce. Après avoir fait l'examen de ce cryptogame, il me sembla ne pouvoir le classer avec justesse dans aucun des genres alors connus, c'est pourquoi j'en établis un nouveau et je l'appelai *Serratia*, du nom d'un illustre Italien, en y joignant les phrases génériques et spécifiques qui suivent: *Funguli acaules, hemisphericî, capsulis contortis S. marescens Vesicula tenuissima, latice primo roseo, dehinc rubro repleta*

(Breed in a personal communication states that "contortis" is a misprint for "confertis," and "marescens" for "marcescens.")

Bizio found the color of dried reddened polenta was insoluble in water, and very soluble in alcohol. By use of mordants, it will dye silk and linen, but the color fades promptly in direct sunlight. A summary of his work was published in 1827 in the first volume of his *Opuscoli chimici-fisici*.

In 1844 Bizio called attention to the report of a French commission to the Minister of War relating to discoloration of war bread.

What is commonly regarded as the same organism was named *Zoogalactima imetropa* Sette 1824, *Monas prodigiosa* Ehrenberg 1848, *Palmella prodigiosa* Montagne, *Bacteridium prodigiosum* Schroter 1872, *Micrococcus prodigosus* Cohn 1872, *Bacillus imetrophus* Trevisan 1887, and *Bacillus marcescens* De Toni and Trevisan 1889.

Pfeiffer (1887) ascribes the genus to Bergamaschi (1822).

Vuillemin (1913, p. 521) concludes that the genus name *Serratia* should be revived for the rod-shaped organisms that have diffuse flagella. He states that the characters as described by Bizio have no generic value. However, inasmuch as the genus *Bacillus* had been "vulgarized" by bacteriologists he proposed that *Serratia* be made a genus *conservandum* with the type species *Serratia subtilis*.

Buchanan (1918, p. 51) has used this as a designation for the red rod-shaped bacteria in the following description

Cells rod-shaped, without spores Motile by means of peritrichous flagella or non-motile Gram stain variable Aerobic, producing a red or pink pigment, a lipochrome Possibly closely related yellow and orange lipochrome-forming bacteria should be included here as well

The type species is *Serratia marcescens* Bizio, the organism usually termed *Bacillus prodigiosus*

Erythrobacillus Fortineau (*qv*) was proposed by Winslow *et al* (1917, 1920, p 209) as a substitute

Bergey *et al* (1923, p 85) include *Serratia* as the first genus of the tribe *Chromobactereae* with the description

Small, aerobic rods, producing a red or pink pigment, usually a lipochrome Gram-negative Motile or non-motile

The type species is *Serratia marcescens* Bizio

Shigella A generic name proposed by Castellani and Chalmers (1918, p 934) for the third genus of the tribe *Ebertheae*, with the following description

Ebertheae non-motile, partially fermenting glucose with the production of acid, but no gas, lactose not fermented Milk not clotted

Types *Shigella dysenteriae* (Kruse, 1899)

It is also used by the same authors (1919, p 937) as the name of a subgenus of the genus *Shigella* to include those types which do not ferment mannitol

Siderocapsa. A generic name proposed by Molisch (1909, p 29) for an organism from the surfaces of immersed leaves of aquatic plants showing iron stains The organisms are epiphytic on parts of plants in fresh water, such as the older stems and leaves of *Elodea*, the under side of leaves of *Nymphaea*, etc, which show a brownish crust Under a magnification of 300 to 500 diameters the surface is seen to be made up of irregularly circular areas, with a lighter center The bacteria are invisible in direct light, and can be differentiated by staining only with difficulty He found colorless "Schiff" solution best When the areas are immersed for a time in this reagent the iron oxid area colors up red-violet and the bacteria of the central area are recognizable In the species *Siderocapsa treubii* there occur one, two or occasionally as many as six cells The diameter of the "rusted" area was 5 to 18 μ , of the clear central area 1.8 to 3.6 μ , and of the coccus 0.4 to 0.6 μ .

In *Siderocapsa major* the cells are coccus-like short rods united into colonies of 2 to 100 cells 5 to 28 μ in diameter. The cells are 0.7 to 1.8 μ long.

The organism does not appear to have been grown in culture media.

Siderocapsa is probably a valid generic designation, with *Siderocapsa treubii* Molisch as the type.

Siderocapseeae. A name given by Buchanan (1917, p. 615) to the third tribe of the *Coccaceae* with the description

Cells spherical or ovoid, non-motile, epiphytic upon the leaves and other parts of water plants. Have not been cultivated. Thick capsules enclosing the cells become encrusted with iron oxide.

One genus only has been described, *Siderocapsa*.

Siphonomyxa. A generic name used by Billroth (1874, p. 27) for his *Siphonomyxa Nosocomii Viennensis* an organism, apparently one of the iron bacteria found in a tap water. He regarded it as a pleomorphic organism resembling his *Coccobacteria septica*, but distinct from it. The name would appear to be invalid, as it is used in a trinomial, and the description does not make it possible to recognize the original.

Solidococcus. A generic name proposed by Orla-Jensen (1909, p. 332) to include cocci which do not as a rule require organic nitrogen, and which belong to the family *Luminibacteriaceae* of the order *Cephalotrichinae*, and do not liquefy gelatin.

The generic name is probably a synonym of *Micrococcus*.

Solidovibrio. A generic name proposed by Orla-Jensen (1909, p. 333) to include short spiral organisms which do not as a rule require organic nitrogen, which often reduce sulfates to hydrogen sulfid, and which do not liquefy gelatin. The description is as follows

Als Schraubenformen, die sich aus der Gattung *Dentritomonas* entwickelt haben, rechne ich die erwähnten sulfatreduzierende Vibrionen, denn das Reduktionsvermögen von Nitrat und von Sulfat müssen ja verwandte Eigenschaften sein. Die weitere Entwicklung dieser bereits spirillenähnlichen Vibrionen, *Solidovibrio*, führt zu den echten (schwefelfreien) Spirillen. Ein wesentlicher Unterschied zwischen den monotrichen Vibrionen und den lophotrichen Spirillen existiert nicht, denn es ist selbstverständlich, dass, während dem die kürzeren Bakterien mit unvollständigen Schraubengang sich mit einer einzigen Geißel als Propell begnügen können, so müssen die längeren, starker gewundenen Formen einen ganzen Buschel haben. Die Gattungen *Solidovibrio* und *Spirillum* wollen wir zur Familie *Reducibacteriaceae* sammeln.

The generic name is probably to be regarded as a synonym of *Vibrio* (q v).

Sphaerobacteria. A group name used by Cohn (1872, p 146) to include all the spherical bacteria. It is equivalent to the modern name *Coccaceae*.

This name was also used by Hansgirk (1888, p 266) for the third order of Bacteria, including one family only, *Mycococcaceae*.

Sphaerobacteries. A name used by Maggi (1886, p 81) for his second order of bacteria, to include the genera *Micrococcus* and *Leuconostoc*, but without description.

Sphaerococcus. Agardh (1821, p 227) used this as a generic algal name with the following description "Char essent Functus uniformis capsulae glomerulum seminum minutissimum sphaericum includentes". On the following pages over 100 species are listed. J S Agardh (1842, p 154) says "*Sphaerococcus* Ag Grev Alg Butt, p 137."

This genus of algae was recognized by Kuetzing (1843, p 408) with the type species *Sphaerococcus confervoides*. In (1849, p 772) he included this in his family *Sphaerococceae*. This name was used by Marpmann (1885, p 121) as the generic designation of a coccus capable of souring milk. The name *Sphaerococcus lactis acidæ* was given to this organism. This genus name has apparently not been used by other writers. It was rejected by Erwin F Smith (1905, p 174). It is invalid for bacteria.

Sphaerokokkus. An aberrant spelling of *Sphaerococcus*, Marpmann used by Eisenberg (1891, p 50).

Sphaerotilus. This genus was described by Kutzin (1833, p 385), with the species *Sph natans*. Later the same author included this genus in his *Phycologia generalis* (1843, p 150) naming three species *Sph natans*, *Sph thermalis* and *Sph lacteus*. The generic description is "Stratum floccosum ex globulis minutissimis aggregatis compositum". The description of *Sphaerotilus natans* is *S natans*, fuscens, floccis fugacissimis."

Bildet sich an Frühlings—und Sommertagen auf stillen Gewässern fast überall. Diese Alge stellt anfangs ein feines braunliches Hautchen dar, welches durch den kleinen Wellenschlag des Wassers zusammengeschoben und an einzelnen Stellen wieder in Gestalt einzelner Flocken denselben und ganzen sich häufig an hervorragende Gegenstände, ohne hangen sich häufig an hervorragende Gegenstände, ohne ihren Zusammenhang zu verlieren. Sie bilden sich besonders bei ruhigem, warmen Wetter und oft so schnell, dass sie in wenigen Tagen einen ganzen Teich bedecken. Diese Decke besitzt eine schmutzige Farne, und wird auch wol häufig für blossen Schmutz gehalten, aber man erkennt unter dem Mikroskope sehr feine organische Kugeln, die einen geringen Zusammenhang untereinander haben.

Indessen kommt es allerdings auch vor, dass fremdartige kleine Körper, besonders Diatomeen sich ansetzen und von den Flocken eingeschlossen werden. So waren unter den Exemplaren, die ich in der *Linnaea* abgebildet habe, eine Menge kleiner Kalktheilchen, daher die mikroskopische Zeichnung nicht genau und hiernach zu berichtigt ist.

In 1849 (p. 147) two other species, *Sph. ochraceus* and *Sph. glaucus* were named.

Apparently the same organism was described under the name *Cladotrix dichotoma* by Cohn (1875, p. 185). Eidam (1876, p. 133) used the generic name *Sphaerotilus*. Winter (1880, p. 62) gives the following diagnosis:

Zellen rundlich-eckig oder langlich, an den Ecken abgerundet, in grosser Zahl reihenweise in einer farblosen Schleimscheide vereinigt zu langen Faden, die dicht zopfartig verflochten und verwirrt schwimmende Flocken bilden. Vermehrung durch sich isolirende, vegetative Zellen, die durch fortgesetzte Theilung neue Faden erzeugen. Fortpflanzung durch Sporen, die endogen in den vegetativen Zellen sich bilden.

Grove (1884, p. 53) uses the same description. It was regarded by Flugge (1886) as a doubtful genus. The generic description was revised by Schroter (1886, p. 173) and included as a valid genus in "Die Pilze Schlesiens."

De Toni and Trevisan (1889, p. 926) recognize both *Cladotrix* and *Sphaerotilus*. The description given of the latter follows:

Filamenta primitus affixa, basi ab apice superiori distincta, mitio simplicia, dein Cladotricis more pseudoramosa a basi ad apicem subaequilata articulata, vagina gelatinosa obducta, in fasciculos crassos floccosus varie divisos consociata. Multiplicatio fragmentis filamentorum secedentibus, quae filamenta et fasciculos, novos efficiunt. Arthrospora numerosissimae, articulorum divisiones in tres directiones ortae.

The genus was also recognized by Ludwig (1892). Migula (1909, p. 1035) redescribed the genus. This description as given by Smith (1905, p. 162) is as follows:

Cells cylindric, enveloped in sheaths, forming dichotomously branched threads with no differentiation into base and apex. Multiplication by means of conidia which swarm out of the sheath, attach themselves anywhere, and immediately grow out into new threads. The conidia possess a tuft of flagella inserted sidewise under one pole.

It should be noted that this same author (Smith 1905) includes *Sphaerotilus* in a list of generic names to be abandoned. Orla-Jensen

(1909, p 331) regards *Sphaerotilus* as a synonym of *Cladothrix* Molisch (1910, p 18) retains *Cladothrix* as the correct generic designation. He says

Migula stellt *Cladothrix* zu *Sphaerotilus* Ich habe vorläufig davon Abstand genommen, weil Bau und Entwicklung der letzteren Gattung noch mangelhaft bekannt sind und weil sich nach Gewinnung von Reinkulturen der *Sphaerotilus natans* und nach einem genaueren Studium dieser vielleicht doch herausstellen dürfte, dass die Aufstellung der Gattung *Cladothrix* neben *Sphaerotilus* berechtigt ist Migula sagt, die Gattung *Sphaerotilus* erzeuge dichotom verzweigte Faden ohne Gegensatz von Spitze und Basis Aber die Faden der *Cladothrix dichotoma* die er zu *Sphaerotilus* stellt, lassen doch deutlich Spitze und Basis unterscheiden

The generic name *Cladothrix* (*qv*), has been more generally used by authors than has *Sphaerotilus*, although the latter term would seem to have priority

Buchanan (1918, p 305) gave the following description

Attached colorless threads showing false branching, making a pseudodichotomy Filaments consist of rod or oval cells, surrounded by a thin, firm sheath Multiplication occurs both by non-motile and motile gonidia, the latter with a clump of flagella near one end

Sphaerotilus natans Kuetzing is the type

Bergey *et al* (1923, p 392) have used the same description and arrangement

Spherobacteria. A variant spelling of *Sphaerobacteria* (*qv*) used by Magnin (1880, p 71)

Spirella. A name of a genus proposed by Dubosco and Lebailly (1912, p 835) to include the new species (type) *Spirella canis* The description of the species follows

Spirella canis ressemble à une petite spiruline qui serait pourvue d'un cil à chaque pôle Certains individus, issus sans doute d'une division transversale recente, paraissent manquer de cil à l'une des extrémités ou n'avoir là tout au plus qu'un court prolongement Le corps est constitué par un filament épais contourné en hélice, à pas serres et à tours plus ou moins nombreux

Les formes les plus communes (formes courtes) ont de 3 à 7 tours de spire Ce sont les seules qui aient été vues par Bizzozero Elles ont l'aspect de petites vis effilées seulement aux extrémités et toujours rectilignes dans leur ensemble Il existe, en outre, dans l'estomac du chien, des formes longues ayant jusqu'à 20 tours de spire, et Regaud en figure de plus longues chez le spirille du chat Ces longues *Spirella*, qui sont souvent intracellulaires (dans les canaux intracellulaires de Golgi des cellules bordantes), sont courbées en crochet, en U, en S ou de façon plus compliquée Leur longueur, comme celle des formes précédentes, peut s'exprimer en μ par le nombre de leurs tours Il faut compter, en plus, les cils terminaux qui mesurent de 3μ à 4μ , quelle que soit la longueur du corps

Etant très réfringentes, les *Spirella* sont très visibles à l'éclairage ordinaire, qui permet même de voir leurs cils terminaux. Par le Giemsa, le corps se colore en bleu et montre de nombreux grains rouges distribués en série assez régulière, souvent par couples. Le bleu de toluidine donne une coloration analogue, le corps étant bleu avec des grains rouges. Mais ceux-ci sont beaucoup moins nombreux, toujours périphériques et faisant même saillie en déformant la spire. Nul doute que ce ne soient des grains métachromatiques (volutine) tandis que certains grains rouges du Giemsa représentent peut-être la vraie chromatine.

A chaque pôle, à la base du cil, est un corpuscule chromatique très bien mis en évidence par l'argent se colorant en rouge par le Giemsa et le Mann. C'est le granule basal découvert par Swellengrebel (1907) chez *Spirillum giganteum*, et comparé par Fuhrmann (1910) à un blépharoplaste.

Dans certaines *Spirella*, les grains rouges sont remplacés par des corpuscules plus gros, de couleur pourpre, qui paraissent être mis en liberté par dissociation du filament. Des corpuscules pareils de *Spirillum volutans* sont interprétés comme spores par Amato (1909).

Edkins (1923, p. 288) has discussed a similar organism from the cat, which he terms *Spirella regaudi*.

Spirilina. The first genus of Bory de St Vincent's (1826, p. 533) family *Vibrionides*. The organisms are without appendages, linear or worm like, opaque, coiled in a discoid spiral.

Spirillaceae. A family name probably first used by Migula (1894, p. 237). He says "*Spirillaceae* Mig. Zellen schraubig gewunden oder Teile eines Schraubenumgangs darstellend. Teilung nur nach einer Richtung des Raumes nach vorausgegangener Langsstreckung." He included the genera *Spirillum*, *Spirosoma*, *Microspira* and *Spirochaeta*. The name was also used by Fischer (1897, p. 32) to include the genera *Vibrio*, *Spirillum* and *Spirochaete*. This family has been quite commonly recognized with the Migula conception by bacteriologists, among them Chester (1899, p. 63), Lehmann and Neumann (1896, p. 107, 1901, p. 125), A. J. Smith (1902, p. 270), Schmidt and Weis (1902, p. 91), Kendall (1902), Fischer (1903, p. 61), Erwin F. Smith (1905, p. 161), Swellengrebel (1907, p. 581), Flugge (1908), Ellis (1909, p. 3), Frost (1911, p. 57), Benecke (1912, p. 188), Schneider (1912, p. 23).

Blanchard (1906, p. 1) makes *Spirillaceae* a synonym of the older name *Spirobacteria*, used by Cohn (1875).

Winslow *et al.* (Committee Soc. Am. Bact., 1917, p. 556) use the following description, including this as the fourth family of the *Eubacteriales*:

Cells elongate, more or less spirally curved. Cell division always transverse, never longitudinal. Cells non-fertuous. Usually without endospores. As a

rule motile by means of polar flagella, sometimes non-motile Typically water forms, though some species are intestinal parasites

Buchanan (1918, p 175) and Bergey *et al* (1923, p 97) used this designation for the third family of the *Eubacteriales*, including the genera *Vibrio* *Spirillum* and *Paraspirillum* Castellani and Chalmers (1919, p 924) include this as the third family of the *Eubacteriales* with the genera *Spirosoma*, *Vibrio* and *Spirillum*.

Spirilleae. The name given by De Toni and Trevisan (1889, p. 1006) to a subtribe of bacteria belonging to the tribe *Bacilleae* The diagnosis given is "*Baculi spiraliter torti Endosporae*" The following genera are included *Spirillum* and *Spiromonas*

Spirillina. A group created by Perty (1852, p 179) to include the genera *Spirillum* and *Spirochaete* The name has never come into common use

Spirillineae. A subfamily name used by Rabenhorst (1865 p 70) to include the following genera *Vibrio*, *Spirillum* and *Spirochaeta* The diagnosis given is

Spirillineae (Cohn in ed) Trichomata abbreviata vel elongata flexuosa vel spiraliter torta flexibilia et mobilis, plus minus distincte articulata, nuda vel mucosulo hyalino achromatico involuta, articuli saepe leviter constricti

Spirillobacteriaceae. A family name proposed by Orla-Jensen (1921, p 264) in the following statement

As to the family names of the bacteria, it will be convenient to let all of them end in *-Bacteriaceae*, by which it will be seen directly what is in question If there are to be formed families of the cocci and spirilla, they must consequently be termed *Coccobacteriaceae* and *Spirillobacteriaceae* (or by the older name of Zopf, *Spirobacteriaceae*)

Spirilloflagellata. According to Dobell (1911, p 536), this name was given to the group of spirochetes by Krzyształowicz and Siedlicki (1908) They include this group among the protozoan group *Mastigophora*

Spirillum. A generic name first used by Ehrenberg (1830, p 38) to include the spiral bacteria The species named is *Spirillum volutans*, the *Vibrio spirillum* of Muller (1786)

Dujardin (1841, p 223) gave the generic description "*Corps filiforme contourne en helice, non extensible quoique contractile*" He described three species, *S undula*, *S volutans* and *S plicatile* Perty (1852, p 179) included species descriptions of *Spirillum volutans*

Sp. undula and *Sp. rufum* Davaine (1859) interpreted the genus as containing any spiral rod Rabenhorst (1865, p 72) gives the following diagnosis: "Trichomata filiformia, divisione spontanea imperfecta articulata, spiralter torta, cochleam rigdam vel cylindriforream simulantia" Cohn (1872, p 181) also used this generic name The organisms described all were found in stagnant water and in infusions Cohn later (1875) included the genus in his tribe *Nematogenes*, and defined it to include short spiral, rigid filaments without phycochrome, free and unbranched The generic designation was used in this sense by Luerssen (1879, p 25), and Magnin (1880, p 94) Trevisan (1879, p 148) gives the generic description "Somatia cylindrica, articulata, valida, abbreviata, rigida, spiralter, torta, anfractibus paucis latis" Three species are listed, *Spirillum tenue*, *S. undula* and *S. volutans* Winter (1880, p 62) included *Vibrio* Mueller (Cohn) and *Ophidomonas* Ehrenberg as synonyms His diagnosis reads

Zellen cylindrisch oder schwach zusammen gedruckt, einfach bogenformig gekrümmt oder spiralig gewunden, starr, an jedem Ende mit einer Geissel (ob bei allen Arten?) Vermehrung durch Querteilung, die Tochterzellen trennen sich meist bald von einander Mitunter kommt auch Zoogloeabildung vor Sporenbildung anhnlich wie bei *Bacillus*.

The same description was used by Grove (1884, p 46)

Zopf (1885, p 61) in accordance with his ideas of pleomorphism redefined the genus to include organisms which are spiral, and which consist of rods only or of rods and cocci De Bary (1884) included in the genus the endospore-producing spiral rods This conception was also held by Hueppe (1885) Zopf (1885) on the other hand stated that spores were lacking or unknown Fluegge (1886) used *Spirillum* in a general or broad sense to include *Spirochaeta*, and in a more limited sense He differentiated in the latter from the genus *Vibrio* because of the more pronounced and extended spirals Schroter (1886, p 167) followed De Bary in regarding the presence of endogenous spores as characteristic Hueppe (1886, p 143) agrees with Schroter

Hansgurg (1888, p 264) recognized this as the sole genus of his subfamily *Spirobacteria* He subdivided the genus into two sections, *Vibrio* and *Ophidomonas* De Toni and Trevisan (1889, p 1006) give the following diagnosis "Baculi cylindrici, spiralter torti, plasmate uniformiter diffuso instructi Sporae (endosporae) microsomae in baculis normalibus immutatis ortae" They include as synonyms *Ophidomonas*, *Spirochaeta* and *Thiospirillum* and list 36 species

Billet (1890, p 25) included all spiral forms Ludwig (1892) used the term to designate all stiff or rigid spiral rods Sternberg (1892) included all curved rods and spiral filaments, whether rigid or flexible.

Migula (1894, p 237) divided the old genus *Spirillum* into *Spirosoma*, *Spirillum* and *Microspira* He says "*Spirillum* Ehren Zellen starr mit polaren Buscheln von 5-20 halbkreisförmig gebogenen Geisseln " The definition of Migula has been quite commonly used by bacteriologists, among others by Fischer (1897, p 32), Mace (1897, p 484), Chester (1899, p 63), Migula (1897, p 6, 1900, p 1017), Lehmann and Neumann (1896, p 344), 1901, p 126), Chester (1901, p 342), Matzuschita (1902), Kendall (1902), A J Smith (1902, p 270), Fischer (1903, p 61), Migula (1904, p 145), E F Smith (1905, p 161), Fluegge (1908), Ellis (1909, p 6), Frost (1911, p 59), Schneider (1912, p 23), Lohnis (1913)

A few others have adhered to the definition that the genus *Spirillum* includes all spiral organisms Among such are Hewlett (1898) and Baumgarten (1900)

Blanchard (1906, p 1) defines the genus as follows

Corps spirale, à section cylindrique, non effilé aux extrémités Pas de membrane ondulante Un ou plusieurs flagelles inflexés en combe régulière, soit aux deux extrémités, soit seulement à l'une d'elles Formation de spores endogènes constatée chez nombre d'espèces Organismes de taille relativement considérable, se cultivant assez aisément sur divers milieux usités en bactériologie

He lists seventeen species that had been described to 1906

Orla-Jensen (1909, p 333) includes this genus of elongated spiral forms as the second in his family *Reducibacteriaceae*, order *Cephalotrichinae* He states that as a rule combined organic nitrogen is not necessary, that in general free oxygen is not required and that the organisms are active reducers

Vuillemin (1913, p 519) has proposed that *Spirillum* Ehrenberg be made a *genus conservandum* with the type species *Spirillum undula* (Müller) Ehrenberg The genus is defined as made up of bacteria with spiral body, polar flagella and an undulating motion

Winslow *et al* (Committee Soc Am Bact, 1917, p 557) give the following generic description

Cells rigid rods of various thicknesses, length, and pitch of the spiral, forming either long screws or portions of a turn Cells motile by means of a tuft of polar flagella (5 to 20) which are mostly half circular, rarely wavy-bent These flagella occur on one or both poles, their number varies greatly and is difficult to determine, since in stained preparations several are often united into a common strand

Endospore formation has been reported in some species Habitat, water or putrid infusions

Buchanan (1918, p 179) includes the genus as the second in the family *Spirillaceae* with the description

Cells definitely spiral, not enlarged at center, motile by means of 5 to 20 polar flagella, or non-motile Not readily cultivated in ordinary culture media

The type species is *Spirillum undula* (Muller) Ehrenberg

This designation of type was also accepted by Winslow *et al* (Com Soc. Am Bact, 1920, p 204) and Bergey *et al* (1923, p. 82)

It should be noted that from the standpoint of zoological nomenclature, *Spirillum* was invalid as a generic designation for a protozoan Stiles (1905, p 34) says:

Judged from the zoological point of view, *Vibrio spirillum* is the type species by absolute tautonymy, but the zoological name *Spirillum*, 1830 was a stillborn homonym, having been used by Oken, 1815, for a genus of polychaete worms It is not the function of the zoological code to determine the names in bacteriology, but it is undoubtedly not the most wise or farseeing policy for either zoologists or botanists (including bacteriologists) to accept unnecessarily a generic name for organisms so near the border line when that name is a homonym in either zoology or botany, should further investigations again bring bacteriology into zoology, the generic names would come under the zoological code, and in this case *Spirillum*, 1830, would be rejected If *Spirillum* is retained in bacteriology, consistency calls for the rejection of the name *Spirillum volutans* Ehrenberg in favor of *Spirillum spirillum* (Muller)

Stiles is in error in holding that the type species should be called *Spirillum spirillum* for Article 55 of the Brussel's Code reads "Specific names must also be rejected when they merely repeat the generic name " The opposite is true in zoology

The question whether the fact that *Spirillum* was a homonym when first given by Ehrenberg should invalidate its use for Ehrenberg's types in bacteriology in one which is not adequately answered by the botanical code Its general acceptance by bacteriologists would argue against its being discarded

The generic name *Spirillum* may probably be regarded as valid

Probably the designation of *Spirillum volutans* Ehr as the type by Stiles (1905) is valid, and has priority over the designation of *Spirillum undula* Ehr by Buchanan, Vuillemin and the Committee

Spirobacillus. A generic name proposed by Metschnikoff (1889, p 61) for an organism *Spirobacillus cienkowskii* This organism was found in *Daphnia* producing a disease which colored this entomostracan

red In the younger stages the organism appeared as a long oval, like certain yeasts, but not more than 3 to 5 μ . Later the cells become longer, narrower, rod-shaped, and finally spiral, similar to *Spirillum volutans*. These spirals break up again into short bent segments, somewhat more slender. Small ovoid bodies often develop. On the death of the infected animal very numerous, small, strongly bent, motile rods, which become *Spiromonas*-like, escape. At the ends of these cells there develop strongly refractive swellings which were regarded as spores.

De Toni and Trevisan (1889, p. 1019) have included this organism in their genus *Pacina*. Their description of the organism reads

Baculis prima aetate magnis ellipsoideis, mox rectis, dein incurvis cylindricis, apice rotundatis, demum in filamenta pseudospiralia consociatis, filamentis postremo in baculos commae vel litterae S administar incurvis, minutissimos, post Daphniae mortem uno apice arthrosporiferos solutis. In ullo medio nutritivo huiusque adhibito non crescit.

Certes (1889) described a giant spirillum as *Spirobacillus gigas* (according to Zettnow, 1903, p. 383) and later studied it in greater detail (1900, p. 714). Zettnow also gives a description of this species.

Blanchard (1906, p. 1) emended the diagnosis somewhat. He states

Bactériacées de forme spiralée atteignant de grandes dimensions. Spores endogènes. Pas de flagelles polaires, mais flagelles disposées, en petit nombre, sur les côtés de chaque segment, comme chez les *Bacillus*. Deux espèces acutellement connues. *Sp. cienkovskiyi* Metschnikoff 1889. *Sp. gigas* Certes 1889.

The genus is rejected as invalid by Erwin F. Smith (1905, p. 174) and by Vuillemin (1913, p. 521), but is recognized by Meyer (1912, p. 3).

It is probably that this is a valid generic designation, but its exact position and nomenclatural status is uncertain.

Spirobacteria. The fourth tribe of bacteria recognized by Cohn (1872, p. 180) as including the spiral or bent forms. Hansgirk (1888, p. 264) used this for his first subfamily of his family *Bacteraceae*. Two genera were included, *Sprochaete* and *Spirillum*. The same designation was also used by Blanchard (1906, p. 1) to include the four genera *Spirosoma*, *Vibrio*, *Spirobacillus* and *Spirillum*. The diagnosis given is

Bactéries plus ou moins incurvées en vis, l'incurvation ne formant dans d'autres cas des tours de spire plus ou moins nombreux. Organisms peu ou pas flexibles, se multipliant par division transversale. Formation de spores endogènes constatée dans nombre d'espèces.

of *Elephantopus* Baker (1902, p 53) insists that the single species described is worthy of generic recognition, and uses *Spirochaeta*. These uses in botany are of course invalidated by the earlier use of *Spirochaeta* by Ehrenberg, if this organism is to be included among plants. The species *Spirochaeta plicatilis* was recognized by Perty (1852, p 179).

Rabenhorst (1865, p. 73) states. "Genus a Spirillo vix discernendum" The single species is described as "Sp longissima, distincte articulata anfractibus numerosissimis"

Cohn (1872, p 180) changed the spelling to *Spirochaete*, (*qv*) a form commonly used at present among bacteriologists. He says "*Spirochaete* mit flexibler und langer enggewundener Schraube" He discusses his observations of *Sp plicatilis*. Later (1875, p. 196) he named the slender spiral organism found in the blood in cases of relapsing fever, *Spirochaete obermeieri*. He included the genus in the tribe *Nematogenes*, and described it as having long filaments, spiral, flexible, without phycochrome, unbranched and free. Trevisan (1879, p 148) gives the following description "Somatia cylindrica distincte articulata, valida, elongata flexilia, spiralter torta, anfractibus numerosis densis" This genus is included in the subtribe *Eurythronaeae*. Three species are included, *Spirochaete plicatilis*, *S obermeieri* of relapsing fever, and *S cohnii* from the tartar of teeth. Luerssen (1879, p 25), Winter (1880, p 60), Grove (1884, p 43), Van Tieghem (1884) and Schroter (1886, p 168) followed closely the diagnosis of Cohn. Cohn (1883, p 198) described a new species from gelatinous growths from wine cellars *Spirochaete Schroeteri*.

Hueppe (1885) differentiated *Spirochaeta* from *Spirillum* in that the former had no endospores. Fluegge (1886) combined the genus with *Spirillum*. Hansgirtl (1888, p 264) included the genus in his subfamily *Spirobacteria*, with the species *Spirochaete ferruginea* (Ehrb.) Hansgirtl.

De Toni and Trevisan (1889, p 1006) regard *Spirochaete* as a section of the genus *Spirillum*. They say "Spirilli regulariter spirales flexiles, valde elongati, spiris primis et secundariis instructi" Eleven species are included in this section.

Billet (1890, p 24) followed Cohn in his definition, as did Sternberg (1892, p 18), Ludwig (1892) and Fischer (1897, p 32). Migula (1894, p 237) says "*Spirochaete* Ehrenb. Zellen schlangenartig biegsam. Bewegungsorgane unbekannt, vielleicht eine undulierende Membran

(Z. B. *Spirochaete plicatilis* Ehrenb.) " Migula (1897, p 87) and later (1900, p. 1029) revised the diagnosis as follows

Zellen schraubig gewunden, dünn, aber meist verhältnissmässig lang, beweglich und fleil, schlangenartig sich windend, aber auch schraubig sich fortbewegend. Bewegungsorgane bisher unbekannt. Endosporenbildung nicht beobachtet. Der Algengattung *Spirulina* nahe verwandt, aber farblos und nicht in einzelne Zellen geghedert.

This description has been adopted by many writers, among them Chester (1897, p 63), Chester (1901, p 347), Lehmann and Neumann (1896, p 348, 1901, p 126), Kendall (1902), Matzschita (1902), A J Smith (1902, p 270), Fischer (1903, p 61), Migula and (1904, p 145).

Stiles (1905, p 94) called attention to the fact that this generic name is usually misspelled, and that the original and correct spelling is *Spirochaeta*. This same point was later brought out by Blanchard (1906, p 2) who showed the spelling *Spirochaete* to be incorrect. He recognizes sixteen species.

Among the later writers who have followed Migula in his treatment of this genus may be mentioned E F Smith (1905, p 161), Ellis (1909, p. 7), Orla-Jensen (1909 p 334), Frost (1911, p 59), Bosanquet (1911, p 69) and Lohnis (1913).

Much interest was centered in the group of flexuous spiral organisms after the discovery of the form causing syphilis by Schraudinn. Difference of opinion was at once expressed as to the propriety of including them with the bacteria. Vuillemin (1905) proposed the generic name *Spironema* (q v) which was found to be invalid. Later in the same year Schaudinn (1905) proposed *Treponema* (q v) and Stiles and Pfender (1905) *Microspironema*.

Blanchard (1906, p 2) places the genus *Spirochaeta* with the protozoa. He gives the following description:

Corps excessivement grêle, spiralé, aplati, l'ectoplasme s'étalant en une étroite membrane ondulante qui entoure en spirale tout le corps. Pas de flagelles, pas de spores endogènes. Un noyau très allongé, filiforme, occupant l'axe du corps, avec grains de chromatine distribués à sa surface. La multiplication se fait selon toute apparence, par division longitudinale. La culture de ces organismes ne réussit sur aucun des milieux usités en bactériologie.

He regards this genus as related to the other flagellates through the genera *Treponema* and *Trypanosoma*. He lists sixteen species as belonging to this genus.

Thesing (1906, p. 351) takes exception to placing the spirochetes among the protozoa, holding them to be typical bacteria, and *Spirochaeta* to include all

Swellengrebel (1907, p. 582) gave the following diagnosis

Cellules sans cils, avec appendice periplastique bien développée et souvent de structure alvéolaire. Quelquefois, il y a une bande de myonemes dans l'appendice. Type *Spirochaeta plicatilis* Ehrenberg

It will be noted that he excludes from the genus the forms parasitic in higher animals. The genus *Spirochaeta* together with *Treponema* and *Borrelia* are included together under the subfamily *Spirochaetaceae* of the family *Spirillaceae*. He regarded the organisms as belonging definitely with the bacteria.

Dobell (1910, p. 536) restricts the generic designation *Spirochaeta* to include only the groups closely resembling the original *Spirochaeta plicatilis* Ehrenberg. The genus therefore includes only the free living water or marine forms, all parasitic types being excluded.

Zuelzer (1911, p. 51) after a careful study of *Spirochaeta plicatilis* came to the same conclusion as Dobell—*Spirochaeta* includes the fresh water and free living spirochetes, *Cristispira*, the mussel spirochetes, and *Treponema*, the group of pathogenic forms. She defines the genus *Spirochaeta* to include the strict anaerobic mesosaprophilic organisms, living usually in the presence of H_2S and growing well in the presence of *Beggiatoa*. The organism is actively flexible, the protoplasm being spirally wound around a straight elastic axis filament. In the plasma spiral there regularly occur volutin granules. The plasma is naked, that is, there is no differentiation of membrane or periplast. The cell is circular in cross section. The organism multiplies by transverse fission. Three new species of *Spirochaeta* were described by this author.

Bosanquet (1911, p. 69) includes all spirochaetes in the genus *Spirochaeta*. This author is inclined to include these under the heading of bacteria. Heim (1911, p. 252) lists the pathogenic forms under the generic name *Spirochaete* with the species *S. obermeieri*.

Doflein (1911, third edition, p. 351) used the generic name *Spirochaeta* in two senses, in a limited sense, to include all slender spiral organisms, and in a somewhat narrower sense, as a term to include fresh water spirals and their near relatives, differentiating the genera *Cristispira* and *Treponema*.

Gross (1911, p. 109) has accepted the definition of *Spirochaeta* used by Zuelzer and insisted upon the necessity for another generic name

for the organism specific for syphilis. The same conclusion was reached by Gleitmann (1913, p 31), but he stated that there is as yet no uniformity in the nomenclature of these forms.

Gonder (1914, p 674) gives the following diagnosis, agreeing in essential points with Zuelzer:

Charakterisiert ist demnach die Gattung *Spirochaeta* wie folgt (stark aktiv flexibel, ein in echten Spiralen stark gewundener, nackter, einliedigen Achsenstab mit Volutinkörnern in den Plasmaspiralen, Fortpflanzung durch echte Querteilung.)

It would seem that there is sufficient evidence to show the advisability of including only those forms closely related to *Spirochaeta phcatilis* in the genus *Spirochaeta*.

Winslow *et al* (Com. Soc. Am. Bact., 1917, p 563) give the generic diagnosis:

Non-parasitic, with flexible undulating body and with or without flagelliform tapering ends. Common in sewage and foul waters.

The type species is *Spirochaeta phcatilis* Ehrenberg.

Buchanan (1918, p 543) states:

Slender, spiral cells, living free, usually in water containing hydrogen sulphide, actively motile, flexuous. Flagella unknown. Anaerobic. Protoplasm is spirally wound around a flexible or elastic axis filament. Volutin granules regularly present in the plasma. No differentiation of exterior. Cell circular in cross section.

The type species is *Spirochaeta phcatilis* Ehrenberg.

The genus is also recognized in this limited sense by Castellani and Chalmers (1919, p 436) and by Bergey *et al* (1923, p 419).

Spirochaetaceae. An order of flagellates named by Fantham (1908) according to Castellani and Chalmers (1919, p 435) who give the following description:

Plasmodiomata, generally parasitic, in form narrow, wavy, and thread-like, with or without an undulating membrane. The cytoplasm is divided into endoplasm and ectoplasm, and is bounded by a flexible, chitinous periplast. The nucleus consists of a spiral achromatic filament, on which are arranged transverse bars or rodlets of deeply staining chromatin.

Two families are included, *Spirochaetidae* and *Treponemidae*.

Spirochaetaceae. A subfamily name used by Swellengrebel (1907, p 581) under the family *Spirillaceae*. This author includes in the

subfamily those spiral bacteria in which the cells are flexible. Three genera are recognized *Spirochaeta*, *Treponema*, and *Borrelia*. As a family name this is used by Schneider (1912, p. 23), coordinate with six other families of bacteria. It is defined to include long, unicellular, flexible, spirally twisted threads without flagella. One genus only, *Spirochaeta*, is recognized.

Winslow *et al.* (Com. Soc. Am. Bact., 1917, p. 562) regard this as a group intermediate between the bacteria and the protozoa with the following description:

Free living or parasitic spirilliform organisms with or without flagella, with undulating or rigid spiral twists. Reproduction by transverse division and by "coccoid bodies," the equivalent of spores.

Spirochaetales. The name proposed by Buchanan (1918, p. 542) for the sixth order of the *Schizomycetes* with the description:

Protozoan-like in many characters. Cells usually relatively slender flexuous spirals. Multiplication of cells apparently by longitudinal division in some types, by transverse division in others, or both.

One family is recognized, *Spirochaetaceae*.

Spirochaete. A variant of the spelling of *Spirochaeta*. Ehrenberg first used by Cohn (1872, p. 180). It was later used by Luerksen (1879, p. 25), Trevisan (1879, p. 148), Van Tieghem (1884), Schroeter (1886, p. 168), De Toni and Trevisan (1889, p. 1006), Billet (1890, p. 24), Sternberg (1892, p. 18), Ludwig (1892), Fischer (1897, p. 32), Migula (1897, p. 47), Lehmann and Neumann (1901, p. 126), Kendall (1902, p. 484), Matzschita (1902), Fischer (1903, p. 61), Migula (1904, p. 145), Thiesing (1906, p. 351), Ellis (1909, p. 7), Heim (1911, p. 252), and by many writers in the modern literature of medical bacteriology. Attention has been drawn to the fact that this form is incorrect by Stiles (1905, p. 94) and by Blanchard (1906).

Spirochaetidae. A family (zoological) name ascribed to Ehrenberg 1833 by Castellani and Chalmers (1919, p. 436). They state: "*Spirochaetacea* with an undulating membrane. Body may be ribbon shaped on transverse section." The genera included are *Spirochaeta*, *Cristispira*, *Saprospira*, *Pseudospira*, and *Spiroschaudinna*.

Spirochaetoidea. A name used for a group of spiral organisms by Dobell (1911, p. 536). Three genera are recognized *Spirochaeta*, *Treponema* and *Cristispira*. He says:

The organisms commonly called Spirochaetes may be conveniently collected into a single group, for which I propose the name *Spirochaetoidea*. This group

comprises three different sets of forms, which may be correspondingly classified in three different genera *Spirochaeta*, *Treponema*, *Cristispira*. These three groups of organisms, whilst showing certain resemblances to one another, possess no definite relations with Protozoa, Bacteria, or Cyanophyceae. The *Spirochaetordea* should therefore be regarded—for the present—as a group of Protista which stands apart.

Spirochetaceae. A variant spelling of *Spirochaetaceae* (*q v*) used by Schneider.

Spirochoeta. A spelling of *Spirochaeta* used by Dujardin (1841) according to Stiles (1905) and occasionally used by medical writers.

Spirodiscus. A generic name used by Ehrenberg (1828, p. 34) as the fifth genus of the Vibrionen. One species, *S. fulvus*, was described from a mountain stream, growing among *Conferva*. The organism is described as forming a flat spiral. Its relationship to the bacteria is very doubtful. It apparently has not been observed since the first description. Ehrenberg states, "Animale e familia Vibrioniorum, divisione spontanea imperfecta (et obliqua?) in catenam filiformem *S. cochleam rigidam disciformem accrescens*."

Dujardin (1841, p. 226) regarded this as a doubtful genus.

Perty (1852, p. 179) questioned the location or identification of this genus in the following statement:

Hier wurde sich noch die Sippe *Spirodiscus* E. anschliessen, da die einzige bekannte Species (*Sp. fulvus* E. p. 86, t. 5, f. 1) nicht etwa eine Pilzspore ist, bei welcher öfters Bewegung wahrgenommen wird.

Migula (1897, I, p. 6) states that perhaps this form belongs with the sulphur spirilla, but that it is in a high degree doubtful whether it should be regarded as belonging to the bacteria. The genus may be regarded as of questionable value.

Spiromonas. A genus described by Perty (1852, p. 171). The organism was described as related to the spirilla, but with the cells flat and band shaped. It multiplies through transverse fission. One species is included, *Spiromonas volubilis*.

The genus was recognized by Trevisan (1879, p. 138) with the description "Somatia compressa ligammiformia, valida, abbreviata, spirallita torta." It was included in his subtribe *Euvibrionae*. Winter (1884, p. 62) gave the following generic description "Zellen "blattartig" (flach) zusammengedrückt, um eine ideale Achse der Länge nach gewunden. Vermehrung durch Quertheilung."

The description given by Winter of *S. volubilis*, is as follows.

Farblos, durchsichtig, glatt, ohne irgend auffallende Differenzirung, Bewegung ziemlich schnell unter rascher Drehung um die Achse, um welche der blattförmige Körper gewunden ist Körper manchmal sehr wenig gewunden, nie mehr als einen Umgang bildend

The second species *S. cohnii* was described by Warming (1876, p 370) Both species are recognized by De Toni and Trevisan (1889, p 1015)

Migula (1897, p 62) states that the morphology of this organism is so poorly understood that it is possible that it should not be included among the bacteria, but rather among the flagellates Erwin F Smith (1905, p 162) includes *Spiromonas* in his list of doubtful genera

Orla-Jensen (1921, p 269) independently proposed this as generic designation in the following statement

Certainly the lophotrichic spirilla, both in regard to their morphological and to their biological properties, form a natural group This thoroughly justifies the setting up of a genus, *Spirillum*, or better *Spiromonas*, a new designation, which would also make it possible to incorporate nearly related monotrichic species in this group

In this sense the genus is invalid

The nomenclatural status of the genus is uncertain The type is *Spiromonas volubilis* Peity

Spironema This name was proposed by Vuillemin (1905) for the generic designation of the organism causing syphilis, which had been named *Spirochaeta pallida* by Schaudinn (1905) This organism was regarded by Schaudinn as a protozoan Schaudinn (October 26, 1905) noted that *Spironema* was preoccupied in zoological nomenclature (by *Spironema* Meek (1864) a genus of molluscs and by *Spiromema* Klebs (1893) a genus of protozoa) He accordingly proposed the generic name *Treponema* Blanchard (1906, p 4) agrees that *Spiromema* is invalid, as does Dobell (1911, p 535) also

Stiles and Pfender (December 2, 1905, p 936), evidently unaware of the previous publication by Schaudinn of the name *Treponema* proposed *Microspironema*

Gross (1912, p 89) gives the following diagnosis "*Spironema* Vuillemin Spironemacea mit wellenförmig gebogenem Körper und Endfaden Theilung mit Ausziehung einer Plasmabrücke"

Gross (1911, p 109) contends that the organism causing syphilis belongs with the bacteria and not the protozoa, and that as *Spironema*

has not been previously used in botany, Vuillemin's designation is tenable. In this he is apparently in error, for *Spironema* was used by Hochstetter (1842, p. 226) for a genus of *Verbenaceae*. This is given as a synonym of *Clerodendron* L. Lindley (1840) used *Spironema* as the name of a genus of the family *Commelinaceae*. Rafinesque (1836) used this name for a genus of the family *Lauraceae*, the name being now listed as a synonym of *Cassytha* L.

It is apparent therefore from the preceding statements that *Spironema* is valid as a generic name for the syphilis organism neither in botany or zoology.

Gross (1912, p. 84) again emphasized his belief in the validity of *Spironema* as a bacterial genus. Gleitsmann (1913, p. 31) regarded as invalid. The name is also used by Gonder (1914, p. 683).

Bergey *et al.* (1923, p. 424) have recognized *Spironema* Vuillemin as a valid generic name with the description "A spiral, flexible body with terminal filaments but no membrane."

Spironemaceae. A group (family?) name used by Gross (1910, p. 88) to include the genera *Spirochaeta*, *Saprospira*, *Cristospira* and *Spironema*. His description is "Spironemaceae nov. fam. Langgestreckte, spiralig oder wellenformig gebogene Bacterien mit flexiblen, aus einer Wabenreihe bestehendem drehrunder Körper. Vermehrung durch Quertheilung."

A later (1911, p. 202) description reads "Vielzellige, lang gestreckte spiralig oder wellenformig gebogene Bacterien mit flexiblen, aus einer einzigen Zellenreihe bestehendem Körper. Vermehrung durch Quertheilung."

Spirophyllum. A genus of bacteria described by Ellis (1907, p. 516) with one species, *S. ferrugineum*. The following description (1907, p. 516) is given:

Body of the cell elongated, flattened and spirally twisted. The number of spiral turns may vary from a quarter to fifteen turns and more. The width varies from 1 to 6 μ . There is no definite membrane, but the edge is thickened so as to form a kind of rampart all around the cell. The ends are usually irregular, angular and unsymmetrical. The spirals may be very close or very wide apart, but the majority of the individuals have spiral lengths three or four times greater than the width. The youngest stages are coated with a thick deposit of ferric hydroxide.

The name is accepted by Orla-Jensen (1909, p. 331) as one of the genera of his family *Trichobacteriaceae*.

Molisch (1910) states that *Spirophyllum* is a synonym of *Gallionella* and that *S. ferrugineum* is a peculiar growth form of *Gallionella ferruginea*.

It is probable that *Spirophyllum* is to be regarded as a synonym of *Didymohelix*.

It may be noted that this name was used by Schindler (1905, p. 82) as the name of a subsection of the section *Pentapteris* of the subgenus *Eumyriophyllum* of the genus *Myriophyllum* of the family *Halo-rhagidaceae*.

Spiroschaudinna. Sambon (August, 1907, p. 833) used this generic name for the organism which is commonly known as *Spirochaeta recurrentis*. He says

Spirochaeta Ehrenberg 1843, pro parte *Spirochaete* Cohn 1875 pro parte. The name *Spiroschaudinna* is here proposed to designate an important group of blood parasites hitherto referred to the genus *Spirochaeta*.

It apparently is an invalid synonym, as *Borrelia g v* has priority.

Gleitsmann (1913, p. 31) called attention to the priority of *Borrelia* for this type.

The genus is recognized by Castellani and Chalmers (1919, p. 439) with the following description.

Spirochaetidae parasitic in the blood and tissues of vertebrates and in some blood-sucking invertebrates.

Remarks. This genus, as we believed would happen, is now recognized by many authorities, and therefore we adopt it, though we had hesitated to do so in the previous edition.

Type species *Spiroschaudinna recurrentis* Lebert, 1874.

Morphology. This has been most carefully studied by Fantham in *S. recurrentis*, *S. duttoni*, and *S. marchouxii*, and he finds that they have long, narrow bodies, bent into many spirals or coils, enclosed in a firm periplast, with a very tenuous membrane, which is often invisible. The nucleus consists of granules of chromatin distributed along the body.

Spirosoma. A generic name proposed by Migula (1894, p. 237). He says "*Spirosoma* nov. gen. Zellen ohne Bewegungsorgane, starr (Z. B. *Spirosoma lingualis* (Weibel) Mig.)". He included non-motile spiral bacteria. Later (1905) five poorly described species are included, among them the *Myconostoc gregarium* of Cohn. Two subgenera were included, *Euspirosoma* and *Myconostoc*. Chester (1899, p. 63), Migula (1900, II, p. 955), A. J. Smith (1902, p. 270), Kendall (1902), Migula (1904, p. 145), E. F. Smith (1905, p. 161), Blanchard (1906, p. 1), Ellis (1909, p. 6), Frost (1911, p. 59), Meyer (1912, p. 3) and others have used this name.

It is rejected by Lehmann and Neumann (1896, p 107) Vuillemin (1913, p 523) proposes that *Spirosoma* shall be retained as a form genus

The genus is probably to be regarded as a synonym in part of *Myconostoc* and in part of *Spirillum*

Spirothrix. Guttman (1880, p 1) in an article on recurrent fever states-

Der erste und eigentlich der einzige Autor, welcher diese Körperchen in charakteristischer Weise beschreibt, ist Bliesener, ohne aber der Neuheit dieses Befundes Erwähnung zu thun Bliesener theilt in seiner Inauguraldisserlation (Berlin), die im Juli 1873 (also wenige Monate nach der Veröffentlichung von Obermeyer über die Spirillen im Blute im Recurrensanfall) erschien, 3 Fälle von Febris recurrens mit, von denen 2 bei wiederholter Untersuchung des Blutes diese Körperche zeigten, nemlich "feine, stark lichtbrechende Kornchen, einzeln oder gepaart, letztere durch einen feinen kurzen Faden verbunden, so dass sie in handtelnähnlicher Form erscheinen, die Kornchen zeigen eine zitternde Bewegung mit langsam fortschreitender Locomotion" Sie fanden sich im Anfall und in der Remission Ob zwischen ihnen und der "Spirothrix" ein engeres Verhältniss bestehe, lässt Bliesener unentschieden

Lewis (1879, p 394) says

Since the period of its discovery in the blood by Obermeier it has been referred to under various names *Spirothrix*, *Protomycetum recurrens*, in Lebert's article on recurrent fever and in Ziemss' "Handbuch" of Medicine, *Spirillum* by Erichsen, Litten, Birch-Hirschfeld, etc, *Spirillum tenue* by Naunyn, and *Spirochaete Obermeieri* by Cohn (fig 15)

The nomenclatural status is uncertain Probably a casual name

Spirulina. A genus of blue green algae named by Turpin (1827), according to West (1904) and Tilden (1910, p 86) Rabenhorst (1865, p 90) gives the authority as Link (1834)

Cohn (1853, p 125) discusses the organism named by Ehrenberg *Spirochaeta plicatilis* under the name *Spirulina plicatilis* Rabenhorst (1865, p 90) gives the following description "Trichomata articulata spira spiralia, motu proprio spirali praedita, in muco matricali achromatice plus minus liquido indulantia. Propagatio adhuc ignota"

Küchner (1878, p 250) used the name *Spirulina ferruginea* for the iron bacterium generally known as *Gallionella* or *Chlamydothrix ferruginea*

Hueppe (1886, p 146) used the name in an entirely new sense to designate an "Untergattung" of his "Arthrobakteriaceen," and as a synonym of *Proteus* Hauser He gave the following diagnosis "Die

Faden können gerade, wellig gebogen, und schraubig gewunden sein " The genus is regarded in part as a synonym of *Spirillum* by De Toni and Trevisan (1889, p 1006) Sternberg (1892, p 18) ascribed the genus to Hueppe, and defined it as follows "The vegetative cells are sometimes rod-shaped and sometimes spiral, in suitable media they may grow out into long, straight, wavy or spiral filaments These filaments may break up into cocci-like reproductive elements—'arthrospores' " This bacterial genus is also accepted by Baumgarten (1900, p 128)

The genus *Spirulina* is at present generally recognized by algologists as a valid genus of the blue green algae Its use for a genus of bacteria is therefore invalid

Sporonema. A genus of bacteria named by Perty (1852, p 181). He gave as a diagnosis of the genus, "Ein aussert kleiner cylindrischer ungegliedeter, hohler Faden schliesst an einem Ende (selten an beiden) ein, manchmal auch zwei elliptische Körperchen (wohl Sporen) ein " The species was named *Sp. gracile* The organism was so incompletely described that it has not since been recognized with certainty The genus has not been used by subsequent authors, as Migula (1892, 1, p 7)

This name had previously been used by Desmazieres (1851, p 319) to designate a genus of *Discomycetes*, now included usually under *Peziza*, and is invalid as a bacterial designation

Sporosarcina A genus of bacteria proposed by Orla-Jensen (1909, p 477) to include spore bearing packet cocci No species are named He says "Mit demselben Recht, wie man die Gattung *Bacillus* aufstellt, kann man auch die Gattungen *Sporospirillum* und *Sporosarcina* aufstellen, und eigentlich ist es inkonsequent, es nicht zu tun "

Sporospirillum A name proposed by Orla-Jensen (1909, p 340) to include the spiral bacteria which produce spores As a rule these forms are stated not to require organic sources of nitrogen No species are named He says "Mit demselben Recht, wie man die Gattung *Bacillus* aufstellt, kann man auch die Gattungen *Sporospirillum* und *Sporosarcina* aufstellen, und eigentlich ist es inkonsequent, es nicht zu tun "

Staphylococcus. Ogston (1883, p 27) first used this name to designate cocci, observed microscopically in pus, in which the cells were irregularly arranged in grape-like clusters No species name was given by Ogston, the genus should not therefore be credited to him Rosenbach (1884, p 19) cultivated two organisms from pus, which he

described and named *Staphylococcus pyogenes aureus* and *Staph pyogenes albus* respectively, or, as they are later designated in his paper *Staphylococcus aureus* and *Staph albus* Fluegge (1886) defined the genus as made up of spherical or oval non-motile cells in irregular masses. Hueppe (1886) also used this name Trevisan (1889, p 32) included this genus as the second in the tribe *Micrococceae*, with the description "Spore microsome" Nine species are enumerated De Toni and Trevisan (1889, p 1072) ascribe the genus to Ogston and give the following description

Cocci primitus globosi indivisi, aetate proevecta in coccos duos biscocctiformiter geminos, latere fratrem versus plus minus complanato, utrinque ad polos isthmis filamentis tenuissimis insimul nexos, scissi, in turmas racemiformiter consociati Endosporae microsomae in coccis normalibus obvinentes

The genus is also recognized by Cornil and Babes (1890, p 145) Lehmann and Neumann (1901, p 123) state

The genus *Staphylococcus* Ogston has no botanical rights, for the property of forming "grape like" clusters is possessed at times by all varieties described today as micrococci The name *Staphylococcus* does not primarily designate any "new" genus Ogston found (microscopically) two forms of micrococci in pus (without cultivating them), grape cocci and chain cocci, and designated them by the well-chosen names of *Staphylococcus* and *Streptococcus* (Billroth) Rosenbach later cultivated the varieties which Ogston had seen, and gave the name *Staphylococcus* to the bunched cocci, which we may today employ as the ordinary name for species of micrococci causing suppuration, and which we will use, but it must be dropped from the botanical classification

The argument of these authors is not particularly convincing As previously noted Rosenbach definitely and adequately described two species of the new genus The whole question is whether these organisms are worthy of generic separation from the older genus *Micrococcus*

Migula (1897) regarded *Staphylococcus* as a synonym of *Micrococcus* and not worthy of separate generic differentiation Most authors in recent years have accepted Migula's views and have used *Staphylococcus* as a casual name to designate a method of cell grouping or a form genus

The Winslows (1905) have proposed to split the old genus *Staphylococcus* into two genera, *Aurococcus* and *Albococcus* If these organisms are worthy of generic rank one should be *Staphylococcus* and the other *Albococcus*

Buchanan (1917, p 612) emended the generic description as follows (including it as the fifth genus in the tribe *Streptococceae*)

Usually parasitic, cells as a rule in irregular groups or short chains, rarely in true packets, usually Gram-positive Growth fair to good on the surface of artificial media Sugars as a rule fermented with acid Gelatin commonly liquefied Nitrates may or may not be reduced Pigment orange or white

It is probable that the genus should be divided upon the basis of pigment production into two genera, *Staphylococcus* and *Albococcus*, or perhaps these should rank as subgenera

The type species is *Staphylococcus aureus* Rosenbach

Winslow *et al* (Committee Soc Am Bact., 1917, p 558, 1920, p. 207), used practically the same description, as did Bergey *et al* (1923, p 24)

It is probable that *Staphylococcus* is a valid generic designation, with *Staphylococcus aureus* Rosenbach the type

Staphylokokken. A variant spelling of *Staphylococcus* (Heim, 1911, p 249)

Staphylokokkus. A variant spelling of *Staphylococcus* used by various German writers

Stigmatella. A generic name proposed by Berkeley and Curtis (1857, p 313) and described by Berkeley (1874) for an organism grouped with the *Hyphomycetes* Saccardo (1886, IV, p. 679) gives the following description

Sporodochia superficialia, globosa, compacta Hyphae seu sporophora arcte fasciculata, filiformia continua, simplicia vel furcata (flexuosa) Conidia magna, globoso-ellipsoideae, continua, hyalina per caudam subpersistentem cuspidatam sporophoris haerentia Genus distinctum, sed quo ad locum systematicum dubium, hinc Aegerliae hinc Naemateliae subaffine De identitate Sphaerocreatis cum *Stigmatella* nullum mihi est dubium

The genus is included under the Tuberculariae mucedineae, Amerosporae

Thaxter (1892, p 339) concluded that this name is a synonym of *Chondromyces* B and C and included this genus in his family *Myxobacteriaceae* The type species is *Stigmatella aurantiaca* B and C

Stoddardillus. A name proposed by Heller (1922, p 19) for the eleventh genus of the *Clostridioideae* with the following description

Clostridioideae of energetic growth habit that liquefy gelatin, but do not produce H₂S demonstrable by a lead-acetate-paper test in blood broth Produce abundant gas but little acid in meat medium Grow very shyly or not at all in milk Attack a few sugars, but do not produce much acid Short chunky Gram-positive rods which do not form spores Colonies in deep agar large, lenticular and opaque Not easily distinguished from *Welchillius* May invade tissue, causing considerable destruction of muscle but no foul odor One strain, reported from a case of human gas gangrene

Type species *S. egens* (*Bacillus egens* Stoddard) as described by (Stoddard (1919a)) Splits glucose, laevulose, mannose, maltose, dextrin, glycogen, inosite and glycerol Does not sporulate on six days' incubation in inspissated serum

Streblotrichia. A generic name used by Guignard (1890, p 142) for a species of marine microorganism *Streblotrichia Bornetii*, which he placed with the bacteria The organism was found in rock fissures by the ocean It shows certain relationship to *Nostoc* in habitat and to *Rivularia* in manner of growth No heterocysts or sporocysts, however, are formed Migula (1900, p 1039) gives the following description

Diese neue Bakteriengattung bildet kleine farblose bis stecknadel grosse Zoogloen von charakteristischem Bau, In der gemeinsamen Gallerte verlaufen, von dem gemeinsamen Anheftungspunkte ausstreckend, zarte 1 μ dicke Bakterienfaden, die anfanglich dicht gedrangt und gerade, später gewunden sind und namentlich in der Peripherie der Zoogloea dicht und regellos durcheinander gewirrt erscheinen Diese Faden sind aus annahrend isodiametrischen Gliedern mit fein granuliertem Inhalte zusammengesetzt und haben eine ziemlich dicke Membran, die in der gelatinösen Grundmasse mitunter schwer zu erkennen ist

The name is rejected by Erwin F Smith (1905, p 174) This is a doubtful bacterial genus

Streblotrichia. A variant spelling of *Streblotrichia* used by Enlows (1920, p 88) Probably a typographic error

Streptobacillus. This name has occasionally been used as the designation of a form genus Maggi (1886, p 173) designated thus a growth form of *Bacillus* Migula (1900, II, p 374) credits the name *Streptobacillus pseudotuberculosis rodentium* to Preisz (1894, p 231) An examination of this paper fails to show however any use of *Streptobacillus* as a generic name The nearest approach is the following statement

Nous proposons de denominer cette maladie la pseudotuberculose des rongeurs (pseudotuberculosis rodentium) et d'appeler avec M Dor Streptobacille son agent producteur, car ce terme rappelle un des caractères importants de ce bacille

Streptobacillus ulcers mollis was described by Unna (1892, p 485) from soft chancres

Ucke (1898, p 996) named an anaerobe from the soil *Streptobacillus terrae*

The organism which is responsible for acid production in the Egyptian sour milk beverage is termed *Streptobacillus lebenis* by Rust and

Khoury (1902, p 65) also by Guerbet (1906, p 495) Olson-Sopp (1912, p 9) described a *Streptobacillus taette* from Norwegian slimy milk

The name is rejected by Erwin F Smith (1905, p 174)

The nomenclatural status of *Streptobacillus* as a generic name is questionable The various organisms described are for the most part quite unrelated Technically perhaps *Streptobacillus terrae* may be considered the type, as the first organism described for which a binomial nomenclature was used

Streptobacter. A subgeneric name used by Schroeter (1886, p. 157 ff) for seven species of the genus *Bacillus*, among them *Bacillus erythrospora*, *B subtilis* and *B anthracis* All the species so designated are characterized by formation of chains of rods This name has apparently not been used by other authors

Streptobacteria A name used by Billroth (1874, p 18) to designate a growth form of his pleomorphic *Coccobacteria septica*, in which the organism occurred as rods in chains He also used it in the combination *Streptobacteria gigas-pericardii* (1874, p 60) for an organism found in pericardial exudate De Toni and Trevisan (1889, p 945) give the following description of the species "Baculis cylindricis, rectis, rigidis, apicibus rotundatis, 3 25-4 x 1 25, subinde in filamenta breviter dispositis, sporis microsomis" These authors term the species *Bacillus gigas* The name is rejected by Erwin F Smith (1905, p 174) Vuillemin (1913, p 519) stated that the genus should not be retained because the parallelism of planes of division is characteristic of all rod-shaped organisms

Streptobacterium A variant of *Streptobacteria* used by Maggi (1886, p 84) to designate a growth form of *Bacterium aceti* Zopf It was also used by Billet (1890, p 23) as a form genus to designate rod shaped organisms which occurred in chains Later it was used by Jacqué and Masay (1912, p 180) in a generic sense in the combination *Streptobacterium foetidum* The organism described is a short rod, with rounded ends, a "coccobacillus" It is actively motile, and does not form spores In bouillon cultures there is a marked tendency to the production of long chains It stains readily, is Gram negative, and with dilute stains shows polar staining The organism is facultative, grows readily in culture media, produces gas from dextrose and maltose, milk is not coagulated, liquefies gelatine and serum, and produces a foetid odor in culture media The organism described is apparently close to *Proteus*

The nomenclatural status of this name is uncertain. The type species apparently is *Streptobacterium foetidum*.

Streptococceae. The third tribe of the family *Coccogenae* used by De Toni and Trevisan (1889, p. 1051) with the following diagnosis: "Cocci in filamenta moniliformiter concatenati. Arthrospora macrosomae in filamentis vel in eorum apicibus obvenientes." The following genera are included: *Leuconostoc*, *Schuetzia*, *Perroncitoa*, *Babesia*, and *Streptococcus*.

Buchanan (1917, p. 609) included this as the first tribe of the family *Coccaceae* with the description:

Cells spherical when isolated. Usually parasitic, growing well in the absence of oxygen, particularly in the presence of carbohydrates, from which acid is developed. Gas rarely produced. Many forms require special media. Abundant surface growths are rarely developed. Planes of fission are commonly parallel, resulting in the formation of pairs, or of longer or shorter chains of cells, regular packets of cells never formed. With the exception of a few strict parasites, the cells are Gram-positive. Pigment white, orange or none.

The genera *Streptococcus*, *Leuconostoc*, *Diplococcus*, *Neisseria* and *Staphylococcus* are included.

Winslow *et al.* (Committee Soc. Am. Bact., 1917, p. 557 and 1920, p. 205) give the following diagnosis:

Parasites (thriving only or best on or in the animal body) except genus *Leuconostoc*. Grow well under anaerobic conditions. Many forms grow with difficulty on serum-free media, none very abundantly. Planes of fission usually parallel, producing pairs or short or long chains, never packets. Generally stain by Gram. Produce acid but no gas in glucose and generally in lactose broth. Pigment, if any, white or orange.

Practically the same description is used by Castellani and Chalmers (1919, p. 924). These authors include as genera *Ascococcus*, *Diplococcus*, *Streptococcus*, *Aurococcus* and *Albococcus*.

The name is also used by Bergey *et al.* (1923, p. 44).

Streptococceae. A variant of *Streptococceae* used by Trevisan (1889, p. 28) for the third tribe of the suborder *Coccogene*. The description is "Cocci seriatim in filamentis." The genera included are *Leuconostoc*, *Schuetzia*, *Perroncitoa*, *Babesia* and *Streptococcus*.

Streptococcus. An aberrant spelling of *Streptococcus* used by Billroth (1874, p. 10).

Streptococcus. This name in the form *Streptococcus* was first used by Billroth (1874, p. 10) as a designation of a growth form of his pleomorphic species, *Coccobacteria septica*. His use of the term was in

no sense generic, and he should not be cited as the authority, as is often the case (Migula, Winslow, Vuillemin, etc.)

Cohn at first did not use the name *Streptococcus* (1875, p 144) in the designation of a genus. He states "Was Billroth *Streptococcus* nennt, hatte ich selbst als *Torula* form von *Micrococcus* bezeichnet." Later, however, he included the genus in the tribe *Nematogenae*. He described it as containing those organisms whose cells are disposed in filaments, filaments unbranched, free, without phycochrome and made up of spherical cells. However, there seems to be no record of any species being assigned to it. This is essential to the validity of a genus, hence Cohn cannot be quoted as authority for the genus.

In 1883 (p 27), Ogston again used the term *streptococcus* as a form group. He states

Micrococcus is met with in two distinct forms, chains and groups. They are often found together, yet the two are different, and the chain form does not pass into the grouped form, nor the grouped into the chain form. Throughout this paper the term *Micrococcus* is used as embracing both forms, the chain coccus is often called *streptococcus* (Billroth), and I shall call the grouped form *staphylococcus*.

Here again the name is not used in a strict generic sense.

Fehleisen (1883) studied the streptococci of erysipelas, and by Zopf is given credit for the name *Streptococcus erysipelatis*. A careful examination of the paper however shows that not once is any binomial used. The organisms are designated as "kettenbildenden Mikrokokken," "Mikrokokkus des Erysipels" and "Erysipelkokken." He did not use the designation *Streptococcus*.

Apparently the first valid use of the generic name *Streptococcus* was that of Rosenbach (1884, p 22). He states

Wollen wir einen *Coccus*, welcher sich aus mehreren Einzelcoccen zu charakteristischen Reihen, Ketten, Ringeln oder rosenkranzähnlichen Figuren gruppiert mit Ogston, welcher Billroth's Nomenklatur acceptirt hat, *Streptococcus* nennen, so bezeichnet auch hier dieses Wort nur eine Gattung, denn es gibt mehrere Arten, welche sich mikroskopisch in gleicher Weise zur Ketten anordnen.

The first species named, that of wound infections, is *Streptococcus pyogenes*, the second, a species previously described by Fehleisen, *Strept erysipelatos*. The genus should therefore be ascribed to Rosenbach.

In the third edition of *Die Spaltpilze*, Zopf (1885, p 51) credits the genus to Billroth. He characterizes the genus as containing cocci

occurring in chains, division only in one direction of space The two species of Rosenbach are recognized

Fluegge (1886, p 121) regarded the name as one for a growth form, used it in a generic sense in naming some nine species Schroeter (1886, p 149) added to the description that spores were unknown He recognized seven species Forty-seven species of *Streptococcus* are listed by Trevisan (1889, p 29) De Toni and Trevisan (1889, p 1054) differentiated several closely related genera from *Streptococcus*, listing fifty-six species in this genus They stated that arthrospores may be found in the chains

Since 1890 this genus has been more generally accepted by bacteriologists than any other with the possible exception of *Micrococcus* Among those who accept the genus in its original sense may be listed Cornil and Babes (1890, p 142), Baumgarten (1890, p 127), Billet (1890, p 24), Hueppe (1891, p 30), Sternberg (1892, p 17), Ludwig (1892), Fischer (1895, 1897, p 32, and 1903, p 60), Migula (1894, p 97, 236 who says "*Streptococcus* (Billroth) Zopf Die Zellen teilen sich nur nach einer Richtung des Raumes, wodurch, wenn sie nach den Teilung verbunden bleiben, perlschnurartige Ketten entstehen können Bewegungsorgane fehlens (Z B *Streptococcus erysipelatis* Fehleisen, *Str mesenteroides* (Cienk))" Migula (1895, 1897, p 46, and 1900, p 15), Hewlett (1898), Chester (1897, p 62, 1901, p 55), Lehmann and Neumann (1896, p 117, 1901, p 123), Matzuschita (1902), Kendall (1902), A J Smith (1902, p 268), Migula (1904), E F Smith (1905, p 160), Fluegge (1908), Ellis (1909, p 4), Jensen (1909, p 340), Heim (1911, p 248), Frost (1911, p 56), Schneider (1912, p 22), Vuillemin (1913, p 525) and Lohms (1913) In 1905 the Winslows (p 669) emended the characterization of the genus to read as follows

Parasites, cells normally in short or long chains (under favorable conditions sometimes in pairs and small groups—never in large groups or packets) On agar streak effused translucent growth with isolated colonies In stab culture, little surface growth Sugars fermented with formation of acid

Buchanan (1915, p 7) reviewed the nomenclatural status of this genus and later (1917, p 611) included it as the first genus of the tribe *Streptococceae* with the following description

Usually parasitic, though able to maintain a saprophytic existence Cells usually in chains of greater or less length, the elements of the chains sometimes arranged as diplococci, never in packets As a rule Gram-positive Growth on surface of usual laboratory media not luxuriant Growth better in liquid media

containing sugars, which are as a rule fermented with acid production, but rarely gas. Usually do not liquefy gelatin or reduce nitrates. No formation of zoogloal masses in sugar solutions.

The type species is probably *Streptococcus pyogenes* Rosenbach.

Winslow *et al* (Committee Soc. Am. Bact., 1917, p. 558 and 1920, 206) also include it in *Streptococceae* with the following description.

Chiefly parasites. Cells normally in short or long chains (under unfavorable conditions, sometimes in pairs and small groups, never in large packets). Generally stain by Gram. Capsules rarely present, no zoogloal masses. On agar streak, effused translucent growth, often with isolated colonies. In stab culture, little surface growth. Many sugars fermented with formation of large amount of acid, but inulin is rarely attacked. Generally fail to liquefy gelatin or reduce nitrates.

Type species is *Streptococcus pyogenes* Rosenbach.

Castellani and Chalmers (1919, p. 927) give the following description.

Streptococceae parasitic in short or long chains or pairs, forming a large quantity of acid in fermented sugars. Haemolysis present or absent. Without characteristic group serum reactions.

Type species *Streptococcus erysipelatos* Fehleisen, 1883.

Bergey *et al* (1923, p. 46) follow the Committee.

The generic designation *Streptococcus* Rosenbach (not Billroth) would appear to be valid, with *Streptococcus pyogenes* Rosenbach as the type. The genus is proposed as a genus conservandum by Vuillemin (1913, p. 525).

Streptokokken. A variant of *Streptococcus* used by certain German writers, as Baumgarten (1890, p. 127).

Streptokokkus. A variant spelling of *Streptococcus* used by various German writers as Hueppe (1891, p. 30).

Streptomicrococcus. A name applied by Billroth (1874, p. 11) to a growth form of his pleomorphic *Coccobacteria septica* in which the cells appeared as small spheres, in chains. It is not a generic name.

Streptosprillum. The name given by Billet (1890, p. 24) to a form genus for those spiral rod shaped bacteria which occur in chains. It is not a valid generic designation.

Streptothrix. A generic name introduced by Corda (1839, p. 27) for one of the *Hyphomycetes*. The single species described, *Streptothrix fusca*, was also figured. This species has been repeatedly figured and described in floras. The description given by Saccardo (1886,

p. 282) is as follows "Hyphae fertiles erectae virgato ramosae, ramis ramulisque spiraliter tortis Conidia globosa v ovoidea, in ramulis solitaria acrogena, subinde pleurogena, sessilia v stipitellata, fusca." Several species of this genus are listed in Saccardo's *Sylloge Fungorum*.

Cohn (1875, p 186) used the generic name *Streptothrix* for a totally distinct organism, one of the thread bacteria producing concretions in the lachrymal duct, with the name *Streptothrix Foersteri*. This species is described by De Toni and Trevisan (1889, p 928) as follows "Filamentis tenuissimis, hyalinis, parallele insimul stratiformi-coalitis rectis vel incurvis, sparse irregulariterque ramosis, in fragmenta inaequalia secedentibus."

Cohn (1875) confused somewhat this genus with *Cladothrix* by his statement that the filaments are cylindrical, colorless, and with *false branching*. He included it in his tribe *Nematogenes*.

Trevisan (1879, p 139) gave the generic description "Somatia cylindrica, leptotrichoidea, indistincte articulata, tenerrima, stricta vel auguste spirale parce ramosa." He included this genus in his subtribe *Cladothriceae*.

Magnin (1880, p 97) states the filaments to be in the form "of *Leptothrix*, very slender, colorless, not articulated, straight or slightly spiral, a little branched."

Much confusion has arisen in the literature of recent years due to the confusion of *Streptothrix* Corda, *Streptothrix* Cohn, *Oospora* Wallroth, *Actinomyces* Harz, *Nocardia* Trevisan, *Cladothrix* Cohn.

It is probable that all uses of the name *Streptothrix* not in accord with the organism named by Corda are invalid. Many authors have used *Streptothrix* in place of *Actinomyces* (q v).

De Toni and Trevisan (1889) used the generic name *Nocardia*.

Gasperini (1889, p 449) identified the *Streptothrix Foersteri* of Cohn as a common air form.

Sauvageau and Radais (1892, p 246) called attention to the use of *Streptothrix* by Corda, and urged that the *Streptothrix* of Cohn was a synonym of the older name *Oospora* Wallroth.

Migula (1894, p 237) says "*Streptothrix* Cohn Zellen zu einfachen unverzweigten Faden vereinigt. Teilung nur nach einer Richtung des Raumes. Endosporenbildung oder eine sonstige besondere Fortpflanzungsform fehlt."

Migula (1897, p 47) gives the following diagnosis of *Streptothrix*. "Zellen zu einfachen, unverzweigten Faden vereinigt. Teilung nur nach einer Richtung des Raumes. Fortpflanzung durch bewegungslose

Conidien" He has been followed by Chester (1899, p 64) Hewlett (1898) groups all thread bacteria under *Streptothrix* Berestnew (1899, p 390) rejected the name as invalid Chester (1901, p 360) uses the following definition:

Cells in their ordinary form as long branched filaments Cultures on solid media raised Growth coherent, dry, rough or crumpled, often with a moldy appearance due to the formation of aerial hyphae Without endospores, but by a multiple segmentation of a filament, the production of short, gonidia like bodies

Lehmann and Neumann (1901, p 128) definitely discard *Streptothrix* in favor of *Actinomyces*

A. J Smith (1902, p 270) and Kendall (1902) agree with Chester.

Petruschky (1903, 2, 832) includes *Actinomyces*, *Streptothrix*, *Cladothrix* and *Leptothrix* as "species" of the *Trichomycetes* *Streptothrix* is used as the generic designation of the organisms causing mycelomata.

E F Smith (1905) rejects the name as invalid for a genus of bacteria. Foulerton (1906, p 302) according to Musgrave and Clegg (1908, p 459) uses the name *Streptothrix* for the following reasons

1 He does not consider that Corda's indefinite use of the term in 1883 should make *Streptothrix* Cohn (1875) untenable

2 This being acceptable, *Streptothrix* becomes the correct botanical name by priority of description

3 *Streptothrix* was adopted by the committee of the Pathological Society

4 It is by far the most generally used name

Musgrave and Clegg (1908, p 471) while agreeing that probably *Streptothrix* Cohn is not tenable, nevertheless adopt it tentatively for lack of a better designation Jordan (1908) holds *Streptothrix* to be invalid Heim (1911) includes *Actinomyces* under *Streptothrix* Petruschky (1913, p 270) believes that no confusion can arise through the use of *Streptothrix* to designate two such unlike organisms as the hyphomycete of Corda and the thread bacterium of Cohn *Streptothrix* is to be differentiated through its abundant true branching, its ability to fragment into segments, its wavy filaments and its chains of conidia



The name *Streptothrix* is probably invalid as a generic designation among the bacteria

Streptovibrio. A name given by Maggi (1886, p 176) to a growth form of *Vibrio* in which the cells occur in chains

Sulfomonas. A genus of sulphur bacteria proposed by Orla-Jensen (1909, p 313) to include those forms which are not spiral, autotrophic.

The genus is included as the first in his family *Thiobacteriaceae* Orla-Jensen's statement follows

Ausser den erwähnten Schwefelbakterien, die also zur vollständigen Ausnutzung des Schwefelwasserstoffes ausgerüstet sind, gibt es nach Natanssohn und Beijerinck auch Bakterien, die Schwefelwasserstoff nur zu Schwefel oxydieren, und somit in demselben Verhältnis zu den ersteren stehen wie der *Nitrosomonas* zum *Bacillus nitrator*. Ein solches von Beijerinck näher beschriebenes Kurzstabchen, *Thiobacillus thioparus*, benutzt die Kohlensäure der Luft als Kohlenstoffquelle und musz deshalb als eine der allerältesten Schwefelbakterien betrachtet werden. Wir wollen ihm den Gattungsnamen *Sulfomonas* geben. Von diesem mikroskopisch nicht zu unterscheiden soll *Sulfomonas* (*Thiobacillus*) *denitrificans* sein, der auch autotroph lebt, und welcher Schwefel zu Schwefelsäure oxydiert. Dies geschieht jedoch nicht mittels des Luftsauerstoffes, sondern mit Salpeter als Oxydationsmittel.



Dieser Organismus bildet also das natürliche Verbindungsglied zwischen den Schwefelbakterien und den denitrifizierenden Bakterien, welche wir jetzt besprechen werden. Hiermit gehen wir zu den bakteriellen Oxydationen über, die mittels gebundenen Sauerstoffes zustande kommen.

Termobacterium. This generic name has been used with species not closely related. Zeidler (1896, p. 739) described an acetic acid organism from beerwort as *Termobacterium aceti*. He later states that this organism is probably identical with *Bact. oxydans* Henneberg. Other species as *T. lutescens*, *T. fuscescens*, *T. iridescens*, *T. album* and *T. erythrinum* have been named by Lindner (1909, p. 508). Henneberg (1909, p. 123) states that organisms called *Termobacterium* are small actively motile wort bacteria. Spores are not produced. It is a group designation for such forms found in wort as do not belong either to the lactic or acetic bacteria. They are suppressed by the growth of yeasts. They are organisms closely related to *Bacillus vulgaris*, the old *Bacterium termo*. Zeidler's organism is not a typical *Termobacterium*.

Delbruck (1910, p. 66) states that this is a generic designation of organisms of the type of *Bact. termo*, that the term was first used by Lindner. The name has been written *Thermobacterium* (*qv*) by some writers. The nomenclatural status of the genus is uncertain.

Termobakterien. A variant spelling of *Termobacterium* used by Henneberg (1909, p. 124) to designate the wort bacteria.

Tetrabacterium. A casual name used by Maggi (1886, p. 85) to designate a growth form of *Bacterium*.

Tetracoccus. A form genus used by Billet (1890, p 24) to designate cocci which occur in fouis The name is rejected by E F Smith (1905). Miquel (1891) also used this generic designation

Apparently the first use in a binomial was by Kleckı (1894, p. 354) who described a species *Tetracoccus butyri* from rancid butter

An independent introduction of a genus name *Tetracoccus* is that of Orla-Jensen (1919, p 154) He says

In this genus (Table XXVII) I include all sugar-fermenting micrococci and sarcinae From the sugar they form, besides lactic acid, smaller or greater quantities of acetic acid The quantity of lactic acid was in many cases so small that we were not able to determine with certainty of what sort it was As they thus stand at the limit of what we will term lactic acid bacteria, we have not sought for them systematically, as for the cocci already described, and thus make no claim to have found, even approximately, representatives of all species belonging thereto, but merely of some of those most frequently met with in the dairy

The first species named is *Tetracoccus liquefaciens* (*Micrococcus casei liquefaciens* Orla-Jensen) Others named are *Tc casei*, *Tc mastitidis*, *Tc pyogenes aureus*, *Tc pyogenes albus* and *Tc mycoder-matus*

Later (1921, p 268) he states

The acid forming micrococci and sarcinae I have brought together in the genus *Tetracoccus*, as I believed it to be quite as wrong to draw a limit between the micrococci and the sarcinae as between the short- and long-chained streptococci

Tetradiplococcus. Bartoszewicz and Schwarzwasser (1908, p 614) proposed the name *Tetradiplococcus filiformans lodzensis* The organism was isolated from the Lodz tap water by means of plate culture In hanging drop the organism appears always in the form of tetrads so constructed that each element of the tetrad is a diplococcus Each cell has the form of the gonococcus The tetrads are actively motile in young cultures, but flagella could not be demonstrated The tetrads are 4 to 6 μ in diameter The organism grows well at room temperature, but much better at 30 to 37° Gelatin is not liquefied The most conspicuous growth character is the formation of slender filaments in broth which rise from the bottom of the tube toward the mouth The filaments do not develop well at room temperature The broth remains clear, but there is a gelatinous deposit In stab cultures in agar a fir tree growth develops Plate colonies are small, white granular known on the surface Gelatin kept at 30° shows the filament formation well Practically an obligate aerobe The cells are Gram-positive

Apparently this name has not been used in a binomial, and is therefore invalid

Tetragenus A name apparently first used as a pseudogeneric designation by Kruse (1896, p 94) As a binomial apparently it was first used by Vincenzi (1897, p 758) for an organism isolated from a lymph node and named *Tetragenus citreus* A second species *Tetragenus tardissimus* was named by Altana (1909, p 44)

If the first species (type) *Tetragenus citreus* Vincenzi is worthy of generic designation, the genus is valid It probably is a synonym

Tetrakokkus. A variant spelling of *Tetracoccus* used by Klebs (1887, p 337) for the species *Tetrakokkus variolae*, the cause of variola

Thermoactinomyces. A genus described by Tsiklinsky (1899, p 500) to include two species of thermophilic thread bacteria, *Thermoactinomyces vulgaris* and *Th lanuginosus* These organisms grow at temperatures between 48° and 68°C

Morphologically these species do not differ from the soil species of *Actinomyces* (qv), of which this name may be regarded as a synonym

The name is rejected by Erwin F Smith (1905, p 174)

According to Enlows (1920, p 92) the type species is *Thermoactinomyces vulgaris* Tsiklinsky

Thermobacillus A generic name proposed by Orla-Jensen (1909, p 33) to include thermophilic rod shaped bacteria, requiring as a rule organic sources of nitrogen The genus is included in the family *Alkalibacteriaceae* and the order *Perithrircinae* The description given by Jensen is as follows

Eigentümlich, für die aeroben Bacillen ist das grosz Temperaturintervall, innerhalb dessen sie gedeihen können Die Heubacillen z B wachsen zwischen 6-50° Eine weitere Anpassung an die höheren Temperaturen und damit Abgewohnung von den niedrigen führt zu den thermophilen Bakterien Diese bilden Sporen, welche hinsichtlich ihrer Widerstandsfähigkeit sogar diejenigen der Kartoffelbacillen übertreffen So erklärt sich das Phanomen, dass hochsterilisierte Milch, welche Wochen lang bei 35° unverändert blieb, bisweilen nach wenigen Tagen bei 70° gerinnt Es wäre vielleicht angezeigt, *Thermobacillus* als selbständige Gattung aufzustellen

Erwin F Smith (1905, p 174) includes *Thermobacillus* in a list of rejected genera A use of the name earlier than that of Jensen has not been found

It is probable that this generic name is synonymous with *Bacillus*

Thermobacterium. Probably a misprint for *Termobacterium* Fuhrmann (1906, p 8) refers to *Thermobacterium aceti* of Zeidler. The name is listed as invalid by E F Smith (1905, p 174).

Benecke (p 442) makes *Thermobacterium Zeidlerii* a synonym of *Bact acetigenum*

As a generic name *Thermobacterium* was proposed by Orla-Jensen (1919) for the thermophilic lactic acid organisms The first species mentioned is *Thermobacterium bulgaricum* In addition he names *Thermobacterium cereale*, and *Tbm lactis* It would seem that this is a synonym of *Lactobacillus* Probably the type species may be designated as *Thermobacterium bulgaricum* Orla-Jensen

Thiobacillus. A generic name used by Beijerinck (1904, p. 593) for organisms oxidizing sulphur, but not containing sulphur granules or bacteriopurpurin The first species named, *Thiobacillus thioparus* oxidizes H_2S , $Na_2S_4O_6$ and other sulphur compounds for energy, and is oligocarbophilous, utilizing carbonates The organism is motile and sporeless The second species *Thiobacillus denitrificans* when grown in the presence of free sulphur, carbonates and nitrates, reduces the last with evolution of free nitrogen, oxidizes the sulphur and utilizes the inorganic carbon

Other species of *Thiobacillus* have been described by other workers as Waksman and Joffe (1922, p 239) who named *Thiobacillus thiooxidans*

Apparently the genus is valid, and the type species *Thiobacillus thioparus* Beijerinck

Bergey *et al* (1923, p 37) include the genus as the seventh in the tribe *Nitrobacteriaceae* with the following description

Small rod-shaped organisms deriving their energy from the oxidation of sulfides, thiosulfates or elementary sulfur, forming sulfur, persulfates, and sulfates under acid or alkaline conditions and deriving their carbon from carbon dioxide or from bicarbonates and carbonates in solution

The type species apparently is *Thiobacillus thioparus* Beijerinck

Thiobacteria. An ordinal name introduced by Migula (1900, p 1039) to include the sulphur bacteria The diagnosis given is "Zellen ohne Centralkörper, aber Schwefeleinschlüsse enthaltend, farblos oder durch Bacteriopurpurin rosa rot oder violett gefärbt, niemals grün" This order is recognized by E F Smith (1905, p 162), Frost (1911, p 60) and others

Vuillemin (1913, p 522) excludes the entire group from the bacteria with the statement that it belongs with the *Schizophyceae*

Thiobacteriaceae A family of the order *Cephalotrichinae* created by Orla-Jensen (1909, p 303) for organisms which contain sulphur but

no bacteriopurpurin, and with cells not in thread-like filaments. The following genera are included *Sulfomonas*, *Thiomonas*, *Thiococcus* and *Thiospirillum*. There is no genus *Thiobacter* or *Thiobacterium* hence the family name is incorrectly constituted.

Thiobacteriales. An order of bacteria named by Buchanan (1917) to include the sulphur bacteria. The following is his description (1918, p 461)

Cells various, typically containing either granules of free sulphur, or bacteriopurpurin, or both, usually growing best in the presence of hydrogen sulphide. The cells are plant-like, not protozoan-like, not producing a pseudoplasmodium or a highly developed encysted resting stage. Spores are rarely or never formed.

Classification within this group is in a very unsatisfactory and very superficial state. Few investigators have studied these forms, and most of the work is old, and in need of careful revision. Undoubtedly many of the genera are to be regarded as growth forms, merely.

Three families were recognized, *Achromatiaceae*, *Beggiatoaceae* and *Rhodobacteriaceae*. The order was recognized by Winslow *et al* (Committee Soc. Am. Bact., 1917, p 549, and 1920, p 196) by Castellani and Chalmers (1919, p 924), Heller (1921, p 449) and Bergey *et al* (1923, p 393).

Thiocapsa. A genus of sulphur bacteria described by Winogradsky (1888, p 84) with the species *Thiocapsa roseo-persicina*. The genus resembles in the grouping and multiplication of the cells the algal genus *Aphanocapsa*. Cell division occurs in all directions of space, the cells are spherical, with thick confluent membranes, which unite to form a structureless gelatinous layer. The cells are a bright rose red color and contain numerous sulphur granules.

Trevisan (1889, p 28) includes *Thiocapsa* as the fourth genus of the tribe *Sarcinee*. The description reads "Cocchi lassamente aggregati in famiglie irregolari nella ganga mucosa disugualmente spianata senza forma definita. Temnogenesi in tre direzioni."

De Toni and Trevisan (1889, p 1049) recognize this genus, as do Migula (1900, p 1043) (1904, p 146), E. F. Smith (1905, p 163) and Frost (1911, p 61). Smith gives the following description.

Non-swarming, globose cells spread out upon the substratum in flat families, which are loosely enveloped in a common gelatin. The membrane is split by the growth of the family, and the cells are separated as if by the swelling of an intermediate substance.

Orla-Jensen (1909, p 334) incorrectly made this a synonym of *Rhodocapsa*.

Buchanan (1918, p 467) gave the following generic diagnosis (including it as the fourth genus of the tribe *Thiocapsaeae*)

Cell families resembling in grouping and multiplication the cells of the algal genus *Aphanocapsa* Cell division occurs in all directions of space, the cells are spherical, with thick confluent membranes, which unite to form a structureless, gelatinous layer The cells are of a bright rose red color and contain numerous sulphur granules The cells do not swarm

The type species is *Thiocapsa roseo-persicina* Winogradsky

Bergey *et al* (1923, p 396) followed Buchanan

Thiocapsaceae. A subfamily described by Migula (1900, p 1042). It includes those sulphur bacteria in which the cells are united into families, and division of the cells takes place in three directions of space The following genera are included, *Thiocystis*, *Thiocapsa* and *Thiosarcina* The subfamily is recognized by Migula (1904, p 146) (1905, p 163) by Frost (1911, p 61) and by Molisch (1910)

Thiocapsaeae A tribal designation proposed by Buchanan (1918, p 467) for the first tribe of the subfamily *Chromatioidae* with the following description "Bacteria containing both sulphur granules and bacteriopurpurin Cells divide in three directions of space, united into families"

The genera included were *Thiocystis*, *Thiosphaera*, *Thiosphaerion*, *Thiocapsa* and *Thiosarcina*

Bergey *et al* (1923, p 395) used the same description

Thiococcus. The third genus of spherical sulphur bacteria of the family *Thiobacteriaceae* used by Orla-Jensen (1909, p 330) He mentions as a type a colorless sulphur coccus described by Warming (1876)

Thiocystis A genus of sulphur bacteria described by Winogradsky (1888, p 60) The spherical cells are massed in small compact families, enveloped singly or several together in a gelatinous cyst, and capable of swimming Usually 4 to 20 or 30 cells are united into a single family The cells are light colored, single cells almost colorless In masses the cells show a beautiful violet or red-violet color The cells are frequently filled with sulphur granules *Thiocystis violacea* and *Th. rufa* were described

The genus was included by Trevisan (1889, p 27) as the sixth of the tribe *Ascococceae*

De Toni and Trevisan (1889, p 1040) give the following description

Cocci globosi vel divisionis tempore globosovoides, modice numerosi, in mucosae matriciali nidulantes, in familia forma indefinitas, irregulares, parvulas, intus

solidas, singulas vel plures simul cystidibus universalibus amplis, crassis homogenis, non lamellosis, gelatinoso-subcartilagineis, involutas, arcte consociatis Cystides speciales nullae Coccorum divisio alternatim in tres directiones

The genus is discussed by Migula (1900, p 1042) (1904, p 146) E F Smith (1905, p 163) and Frost (1911, p 61) Smith gives the following description:

Families small, compact, enveloped singly or several together in a gelatinous cyst, capable of swarming When the families have reached a definite size they escape from the gelatinous cyst, the latter either swelling and softening uniformly or at some particular spot The escaped cells either pass into the swarm stage or unite into a larger fused complex of families, the individual cells of which separate and swim away only after a long time, and by means of much vigorous struggling.

Orla-Jensen (1909, p 334) incorrectly made this generic name a synonym of *Rhodocystis*

Buchanan (1918, p 466) include this as the first genus of the tribe *Thiocapsee* with the following description

Usually 4 to 20 or 30 cells massed into small, compact families, enveloped singly or several together in a gelatinous cyst, capable of swarming When the families have reached a definite size they escape from the gelatinous cyst, the latter either swelling and softening uniformly or at some particular spot The escaped cells either pass into the swarm stage or unit into a larger fused complex of families Cells are light colored, single cells almost colorless In masses the cells show a beautiful violet or red violet color The cells are frequently quite filled with sulphur granules

The type species is *Thiocystis violacea* Winogradsky

Bergey *et al* (1923, p 395) used the same description

Thoderma. A name proposed by Miyoshi (1897, p 158) to include certain sulphur bacteria Two species were described, *Thoderma rubrum* and *Th roseum* The later is described as follows *Thoderma roseum* nov gen e spec Cells spheroidal, 2.5μ long and 1.5μ wide of a light rose color, containing small, inconspicuous sulphur granules United into thin, removable, smooth purple red skin On grass culms, leaves, etc, on soil Migula (1900, p 1044) concluded the generic name to be a synonym of *Lamprocystis*

Buchanan (1918, p 468) included this as the second genus of the tribe *Thiopediaceae* with the following description

Cells spheroidal, light rose in color, containing small, inconspicuous, sulphur granules United into thin purplish membrane

The type species is *Thoderma roseum* Miyoshi

Bergey *et al* (1923, p 398) used the same description

Thiodictyon. A genus of sulphur bacteria described by Winogradsky (1888, p 80) The organisms are rod-shaped, with ends united to form a net work The species *Thiodictyon elegans* was described by this author Winogradsky gives the following description of this organism

Die Stäbchen sind schlank spindelförmig, mit scharf zugespitzten Enden, 5μ lang, $1,7\mu$ dick, vor der Theilung fast doppelt so lang, nach der Theilung ist das eine Ende in der Einschnürung abgerundet, bei weiterem Wachstum spitzt es sich wieder zu Auch in gut wachsendem Zustande zeigen die Zellen in ihrem Inhalte die charakteristische "Vacuole," deren Umrisse denen des Stäbchens parallel sind Das Plasma bildet nur eine sehr schmale wandständige Schicht, in der allein (nie in der "Vacuole") winzige, punktförmige Schwefelkörnchen eingelagert sind Die Schwefelaufspeicherung ist immer, auch unter den dafür günstigsten Bedingungen relativ unbedeutend Die Färbung ist so schwach, dass die Stäbchen bei gewöhnlicher Beleuchtung manchmal farblos erscheinen, wenn man aber dieselben bei starker Beleuchtung mit dem Abbe'schen Condensor ohne Diaphragma betrachtet, merkt man immer einen schwachen, aber deutlichen violetten Ton (p 82)

De Toni and Trevisan (1889, p 941) gave the following generic description

Baculi fusiformes vel ellipsoideo-cylindracei, recti, apicibus conformibus, in coenobium reticulatum, Hydrodictyū utriculati coenobia simulans, connexi Multiplicatio cingulis () 5-15-cellularibus qui a colonia matricali movimento peculiari secedunt

The genus is described by Migula (1900, p 1046) as follows

Familien aus stäbchenförmigen, mit ihren Enden zu einem Netz verbundenen Zellen bestehend Aus einer anfanglich kompakten Stäbchenmasse gehen durch allmähliche Umlagerung hydrodictyonartige Zellverbände hervor, die sich unter ungünstigen Verhältnissen wieder in solche kompakte Stäbchenmassen zusammenziehen können Die Vermehrung der Familien erfolgt durch Teilung oder durch Ablösung langsam beweglichen kleiner Zellverbände

The genus was noted by Migula (1904, p 146), E F Smith (1905 p 164) and Frost (1911, p 61)

Smith gives the generic description as follows

Families consisting of rod-shaped cells having their ends united into a net Out of an originally compact mass of rods there gradually results from re-arrangement a *Hydrodictyon*-like cell-union, which under unfavorable conditions, may again draw together into a compact mass of rods The multiplication of the families results from division or by loosening of slowly motile small cell-colonies

Orla-Jensen (1909, p 334) incorrectly termed this genus a synonym of *Rhododictyon*. Buchanan (1918, p 469) included this as the second genus of the tribe *Amoebobacterineae* with the following description

Cells rod-shaped or spindle-shaped, with sharply pointed ends, united into a net. The compact mass of rods finally assumes an appearance like that of *Hydrodictyon*. Slight violet color

The type species is *Thiodictyon elegans* Winogradsky

Bergey *et al* (1923, p 399) used the same description

Thiomonas. A genus of sulphur bacteria belonging to the *Thiobacteriaceae* named by Orla-Jensen (1909, p 330). The organisms are not autotrophic, cells rod shaped, not spiral. The organisms described by Hinze (1903, p 309) and Jegunow (1896, p 11) are assigned to this genus

Thiopedia. A genus name used by Winogradsky (1888, p 85) to include *Merismopedia*-like organisms with bacterio-purpurin and sulphur granules. The species described is *Th. rosea*. De Toni and Trevisan (1889, p 1048) regard this as a synonym of *Lampropedia* Schroeter. The genus is discussed by Migula (1900, p 1045) (1904, p 146), E. F. Smith (1905, p 163) and Frost (1911, p 61). Smith gives the following description of the genus "Families tabular, formed of cells arranged in fours, and capable of swarming"

Orla-Jensen (1909, p 334) incorrectly makes this the genus *Rhodococcus*.

It seems that *Lampropedia* has priority over *Thiopedia*

Thiopediaceae A subfamily name used by Migula (1900, p 1044) to include those sulphur bacteria in which the cells are united into families and cell division is in two directions of space. The only genus is *Thiopedia* Winogradsky. The subfamily is recognized by Smith (1905, p 163) and Frost (1911, p 61)

Thiopediaceae. The third tribe of the subfamily *Chromatiordeae* as named by Buchanan (1918, p 468) with the description "Sulphur bacteria in which the cells are united into families, and cell division is in two directions of space, resulting in the development of plates of cells"

The genera included are *Lampropedia* and *Thioderma*

Bergey *et al* (1923, p 397) use the same description

This designation probably should be changed to *Lampropediaceae*, as the tribe does not contain a valid genus *Thiopedia*.

Thiophysa. A generic name given by Hinze (1903, p 310) to a sulphur bacterium from the Bay of Naples

In der typischen Form kugelige, mit schwefeltröpfchen beladene Zellen, welche von einer die Reaktionen der Pektinstoffe gebenden Membran umgrenzt sind. Der protoplasmatische Wandbeleg umschliesst eine grosse centrale Vakuole, ein Zellkern ist nicht nachweisbar. Geisseln fehlen. Vor der Teilung streckt sich die Zelle in die Länge, schnürt sich biscuitförmig ein und zerfällt in zwei sich später gebundene kalottenförmige Zellen.

One species, *Thiophysa volutans* was named. The cells vary from 7 to 18 μ in diameter. Cells possess power of very slow motion. The central portion or "vacuole" contains sulphur.

This genus is recognized by Buchanan (1918, p. 463) as the second of the family *Achromataceae* with the description:

Spherical cells laden with sulphur. The protoplasmic layer surrounds a large central vacuole. Cell nucleus not recognized. Flagella lacking. Cells elongate before division, divide to biscuit-shaped cells. Cells 7 to 18 μ in diameter.

The type species *Thiophysa volutans* Hinze was secured from the Bay of Naples.

Bergey *et al* (1923, p. 411) gives the following generic description:

Spherical cells, the cell membrane of which is laden with sulphur granules. The protoplasmic layer surrounds a large central vacuole. The ovalate is contained in the vacuole. Cell nucleus not recognized. Flagella lacking. Cells elongate before division, divide into biscuit-shaped cells. Cells 7 to 18 microns in diameter.

In the presence of an excess of oxygen the sulphur drops disappear and only the ovalate remains. With a lack of oxygen, in the presence of H_2S the ovalate disappears and sulphur drops fill the cell.

Thioploca A generic name given by Lauterborn (1907, p. 238) to one of the sulphur bacteria. The genus is placed in the family *Beggiatoaceae*. The generic description is:

Faden von Beggiatoaartigem Habitus, mit reichlichem Schwefelkornern, beweglich, in oft beträchtlicher Zahl parallel nebeneinander verlaufend, zu seilartigen Bündeln vereinigt und verflochten. Nach aussen umschlossen von weit abstehenden farblosen Gallertrohren, meist mit Schlammteilchen inkrustiert und bisweilen mit ringförmigen Einschnürungen versehen.

The species is described as follows:

Zellen der Faden 5-9 μ dick, 1-1½ mal so lang als breit, Gallertschlauche 50-160 μ dick, bis mehrere Centimeter lang. Vorkommen: Untersee des Bodensees in der Gegend von Ermatingen, in 15-20 m Tiefe des Innern des kalkreichen Grundschlicks durchziehend.

Buchanan (1918, p. 464) gives the following description: The genus is included as the third in the family *Beggiatoaceae*.

Filaments *Beggiatoa*-like, with numerous sulphur granules, motile, lying parallel in considerable numbers, or united in bundles enclosed in a colorless layer of gelatin

The type species, *Thioploca schmidleri* Lauterborn has filaments 5 to 9 μ thick, and gelatinous sheath 50 to 160 μ thick From the ocean bed

Bergey *et al.* (1923, p. 410) uses the same description and position of the genus

Thioplococcus. A genus of sulphur bacteria named by Winogradsky (1888, p. 79) with one species *Thioplococcus ruber* The description given follows

Die gestaltlosen dichten Aggregate von kleinen, runden, lebhaft gefärbten Zellen, welche man zusammen mit *Amoebobacter*, *Lamprocystis* u s w findet, gehören manchmal zu dem in der Ueberschrift genannten Organismus Nur die Beobachtung der weiteren Entwicklung dieser soliden Häufchen kann entscheiden, ob sie nicht vielleicht zu anderen rothen Zoogloea-Formen, deren Entwicklung oben geschildert ist gehören Die *Thioplococcus*-familien sind, im Gegensatz zu *Amoebobacter*, vollkommen unbeweglich und bilden bei ihrem weiteren Wachstum keine hohlkugeligen Zoogloeen wie *Lamprocystis*, sondern stellen immer grössere oder kleinere solide und ganz unregelmässige Coccen-Aggregate dar

De Toni and Trevisan (1889, p. 1044) recognized the genus and gave the following description

Cocci globosi vel divisionis tempore ovoidei, numerosissimi, in familias solidas, immobiles, cumuliformes, sine forma definita, tegumento gelatinoso tenuissimo vix conspicuo praeditas, densissime aggregati Coccorum divisio in unam directionem Sporae ignotae

The genus was also described by Migula (1900 II, p. 1046)

Familien solid, unbeweglich, aus kleinen, dicht zusammen gepressten Zellen bestehend Vermehrung der Kolonien erfolgt dadurch, dass die Kolonien sich an der Oberfläche auflöcker und zahlreiche kurze Fitzen und Lappen entstehen und diese in immer kleiner Häufchen zerfallen

E. F. Smith (1905 p. 164) uses the following description

Families solid, non-motile, consisting of small cells closely pressed together Multiplication of the colonies by the breaking up of the surface into numerous short shreds and lobes which continue to split up into smaller heaps

Orla-Jensen (1909, p. 334) incorrectly makes this a synonym of *Rhodoplococcus*

Buchanan (1918, p. 470) made this the fourth genus of the tribe *Amoebobacterieae* with the description

Families solid, non-motile, consisting of small cells closely appressed. Multiplication of the colonies by the breaking up of the surface into numerous short shreds and lobes which continue to split up into smaller heaps. Cells red.

The type species is *Thiopolycoccus ruber* Winogradsky.

Bergey *et al.* (1923, p. 400) followed Buchanan.

Thiosarcina. A generic name proposed by Winogradsky (1888, p. 104) for the species *Thiosarcina rosea*, named *Sarcina rosea* by Schroeter (1886, p. 154). The genus is regarded as a synonym of *Sarcina* by De Toni and Trevisan (1889, p. 1044). The genus has been described by Migula (1900, p. 1043) (1904, p. 146), E. F. Smith (1905, p. 163), and Frost (1911, p. 61). Smith's description follows: "Non-swarming cells arranged in packet shaped families, corresponding to the genus *Sarcina* in the *Eubacteriaceae*."

Orla-Jensen (1909, p. 334) incorrectly makes this genus a synonym of *Rhodospirillum*.

Buchanan (1918, p. 467) describes this as the fifth genus of the tribe *Thiocapseeae* as follows:

Non-swarming cells arranged in packet shaped families, corresponding to the genus *Sarcina*. Cells red, with sulphur granules.

The type species is *Thiosarcina rosea* (Schroeter) Winogradsky.

Bergey *et al.* (1923, p. 396) followed Buchanan.

Thiosphaerella. A generic name given by Nadson (1914, p. 52). Enlows (1920, p. 94) gives the following description:

Type species (monotypy). *T. amyliifera*. A sulfur bacterium. Cells are round or slightly elliptical, measuring 4.8 by 6 μ . A very thick cell membrane enveloped in a colorless gelatinous layer. Protoplasm sometimes has a gray-green color, and in it are found sulfur granules, and a substance resembling starch. Motile. Multiplication by transverse division. Found frequently associated with *Thiophysa* Hinze and *Achromatium* Schewiakoff.

Thiosphaera. A genus of sulphur bacteria named by Miyoshi (1897, p. 170) with the species *Thiosphaera gelatinosa*. The description given by Migula (1900, II, p. 1044) is: "Zellen sphaeroellipsoidisch, 7 μ und 8 μ gross, leicht violett gefarbt, mit einer farblosen Gallertsubstanz zu einer Familie locker verbunden. Schwefeileinschlüsse ziemlich Zahlreich. Schwarmfähig."

Migula changed the name (p. 1044) to *Lamprocystis gelatinosa* (Miyoshi) Migula.

The genus is rejected by E. F. Smith (1905, p. 174).

Buchanan (1918, p. 466) included this as the second genus of the tribe *Thiocapsee* with the description

Cells spherical-ellipsoidal, relatively large (7 to 8μ) light violet in color, bound into loose families by a colorless gelatin Capable of swarming Sulphur inclusions relatively abundant

The type species is *Thiosphaera gelatinosa* Miyoshi

Bergey *et al* (1923, p 396) followed Buchanan

Thiosphaerion. A genus of sulphur bacteria named by Miyoshi (1897, p 170) for the species *Thiosphaerion violaceum*. The description given by Migula is. "Zellen spharo-ellipsoidisch, ca 2.5 und 1.8 μ groß, violett gefärbt, mit winzigen Schwefeleinclusionen, durch eine gallertartige Substanz zu soliden rundlichen Familien verbunden Schwarmfähig "

Migula calls it *Lamprocystis violacea* (Miyoshi) Mig and says

Ich halte diese wie die 3 folgenden Arten nicht für so verschieden von *Lamprocystis roseo-persicina*

An Faden von *Thiothrix nivea*

The genus is not accepted by E. F. Smith (1905, p 174)

Buchanan includes this as the third genus of *Thiocapsee* with the following description

Cells spherical-ellipsoidal, small (1.8 to 2.5 μ) violet in color, with delicate sulphur inclusions United by means of gelatin into solid spherical families Capable of swarming

The type species is *Thiosphaerion violaceum* Miyoshi

Bergey *et al* (1923, p 396) followed Buchanan

Thiospira. A generic name proposed by Wislouch (1914, p 50) Enlows summarizes the description as follows

Motile, colorless, slightly curved sulfur spirilla with pointed ends Sulfur granules present A few polar flagella *T. winogradskii* (Omélieński) A giant sulfur spirillum 3.5 μ by 50 μ *T. bipunctata* (Molisch) Small, very delicately curved sulfur spirilla 1.7 to 2.4 μ by 6.6 to 14 μ long

Bergey (1923, p 411) includes this genus as the third in the family *Achromatiaceae* with the description "Colorless, motile, slightly bent, somewhat pointed at the ends, with drops of sulphur within the cells and a small number of flagella at the ends "

The type species may be designated as *Thiospira Winogradskyi* Wislouch

Thiospirillum. A genus of red sulphur bacteria named by Winogradsky (1888) to include the spiral sulphur bacteria. The genus is apparently a synonym of *Ophiodomonas* Ehrenberg (*q v*). Two species, *Thiospirillum sanguineum* and *Th. jenensis* are included.

The genus is reduced to subgeneric rank under *Spirillum* by De Toni and Trevisan (1889, p. 1012). It is recognized as valid by Migula (1900, II, p. 1049), "Zellen frei, zeitlebens schwarmfähig, spiralig gewunden" (1904, p. 146), E. F. Smith (1905, p. 164) and Frost (1911, p. 62).

Oila-Jensen (1909, p. 330) used this generic name to include colorless sulphur bacteria of the type described by Omelianski (1905, p. 769) changing the name of the corresponding red forms (incorrectly) to *Rhodospirillum*.

Enlows (1920, p. 94) designates the type species as *Thiospirillum sanguineum* (Ehrenberg) Winogradsky.

Buchanan (1918, p. 471) gives the following description:

Spiral motile bacteria containing sulphur granules and bacterio-purpurin. The type species is *Thiospirillum sanguineum* (Ehrenberg) Winogradsky.

Bergey *et al.* (1923, p. 402) followed Buchanan.

Thiothece. A genus of sulphur bacteria named by Winogradsky (1888, p. 82) with the species *Thiothece gelatinosa*. Winogradsky gives the following description:

Dieser Organismus zeigt in seinem Wachstumsmodus eine vollkommene Uebereinstimmung mit der *Chroococcaceen*-Gattung *Aphanothece* Nageli. Indessen wäre eine Vereinigung dieser neuen Gattung mit *Aphanothece* unstatthaft oder wenigstens verfrüht. Von dieser unterscheidet sie sich durch ihr Pigment und den Schwefelkornergehalt, hauptsächlich aber durch die Fähigkeit der Zellen, in den Schwarmzustand überzugehen, was bis jetzt unter allen Chroococcaceen, soviel ich weiss, nicht bei *Merismopedia* beobachtet worden ist.

De Toni and Trevisan (1889, p. 1040) recognize the genus with the following diagnosis:

Cocci globosi vel divisionis tempore globosovoides, pauci, in muco matriciali sparsi, remoti, cystidibus universalibus amplissimis, crassissimis, persistentibus, homogeneis, non lamellosis, gelatinosubcartilagineis, persistentibus, involuti. Cystides speciales nullae. Coccorum divisio in unam directionem.

Migula (1900, II, p. 1046) (1904, p. 146), E. F. Smith (1905, p. 164) and Frost (1911, p. 61) discuss the genus. The description given by Smith follows:

Families enclosed in a thick gelatinous cyst Cells capable of swarming and very loosely embedded in a common gelatin When the swarm stage supervenes, the cells lie more loosely, the gelatin is swollen, and the cells swarm out singly and rather irregularly

Orla-Jensen (1909, p 334) renamed this genus *Rhodotheca*

Buchanan (1918, p. 469) included this as the third genus of the tribe *Amoebobacterieae* with the description

Cells spherical, in families enclosed in a thick, gelatinous cyst Cells capable of swarming and very loosely embedded in a common gelatin When the swarm stage supervenes, the cells lie more loosely, the gelatin is swollen, and the cells swarm out singly and rather irregularly

The type species is *Thiotheca gelatinosa* Winogradsky

Bergey *et al* (1923, p 399) followed Buchanan

Thiothrix. A genus of sulphur bacteria proposed by Winogradsky (1888, p 39) with the following diagnosis

Faden unbeweglich, gegliedert, mit einer zarten Scheide, einen deutlichen Gegensatz von Basis und Spitze zeigend, durch ein Gallertpolster an feste Gegenstände befestigt, unter normalen Wachstumsbedingungen dicht mit Schwefelkörnchen gefüllt, Reproduction durch Stabchengeronidien, welche auf festen Gegenständen kriechend sich langsam bewegen, nach kurzer Bewegungsdauer sich auf verschiedene Gegenstände festsetzen und zu Faden auswachsen

The genus is recognized by many subsequent authors, among them Migula (1897, p 40), Fischer (1897, p 32), Migula (1894, p 238) says "*Thiothrix* Winogradsky Unverzweigte in feine Scheiden eingeschlossene, unbewegliche Faden mit Teilung der Zellen nach einer Richtung des Raumes Die Zellen enthalten Schwefelkörnchen)," Chester (1899, p 64), Migula (1900, p 1039), Chester (1901, p 378), Fischer (1903, p 61), E F Smith (1905, p 162), Orla-Jensen (1909), Heim (1911, p 254), Frost (1911, p 60) The generic description by Smith is as follows

Threads attached, not uniformly thick, enveloped in a delicate sheath which is not easily demonstrable, non-motile, contents containing sulphur granules The threads produce rod-shaped conidia at their end These conidia, which are self-motile by means of a slow, creeping motion, attach themselves by one end to any sort of substratum, extrude a slime cushion at the base, bend over ordinarily in their middle to a nearly right angle and grow into a new thread Habitat, hot sulphur springs

Buchanan (1918, p 463) included this genus as the first in the family *Beggiatoaceae* with the following description

Filament non-motile, segmented, a definite differentiation into base and tip, attached, usually filled with sulphur granules. The threads produce rod shaped conidia at their ends. These conidia are self motile by means of a slow creeping motion, attach themselves and develop into new threads. The habitat is hot sulphur springs.

The type species is *Thiothrix nivea* (Rabenhorst) Winogradsky.

Bergey *et al* (1923, p 408) followed Buchanan.

Thiotrix. A variant spelling of *Thiothrix* used by Schmidt and Weis (1902, p 92). It is spelled thus in a key. Probably a misprint, as *Thiothrix* is used later in the text.

Tilletia. A generic name given by Tulasne (1847) to a group of the smut fungi. According to Trevisan (1879, p 60) several bacteria were incorrectly placed in the genus by Halber (1869) under the names of *Tilletia scarlatinosa* and *Tilletia diphtherica*.

Tissierillus. A name proposed by Heller (1922, p 27) for a genus of *Putrificoideae* with the description

Putrificoideae that attack sugars and clot milk. Slender Gram-negative or Gram-positive rods with oval terminal spores. Colonies in deep agar have radiate periphery and opaque center. Frequently intestinal saprophytes.

Type species *T. paraputrificus* (*Bacillus paraputrificus* defined by Bienstock) as described by McIntosh under the name *Bacillus putrificus*. Ferments glucose, maltose, lactose, sucrose and starch.

Treponema. A generic name proposed by Schaudinn (1905, p 1728) for the species *Treponema pallida*, the causal organism of syphilis. This organism was first named *Spirochaete pallidum* by Schaudinn and Hoffmann (1905), later the generic name *Spironema* (*q v*) was proposed by Vuillemin (1905). Schaudinn noted that *Spironema* was pre-occupied, and proposed *Treponema*. A short time later *Microspironema* was proposed by Stiles and Pfender (1905). Blanchard (1906, p 4) gives the following generic diagnosis:

Corps spirale, non aplati, à section cylindrique, effilé aux extrémités. Un flagelle à chaque extrémité, pas de membrane ondulante. Multiplication par division longitudinale, le stade initial pouvant être marqué par la duplicité du flagelle à l'une des extrémités. Une seule espèce certaine. *Tr. pallidum* (Schaudinn (1905)).

This generic name was also used by Swellengrebel (1907, p 582). Dobell (1910, p 536) extended the definition to include all the spirochetes parasitic in animals. Doflein (1911), treats *Treponema* as a subgenus under *Spirochaeta*. Gonder (1914, p 687) gives the following description:

Der einzellige Organismus ist stark flexibel, wodurch neben der schrauben oder korkzieherartigen Bewegung auch eine schlängelnde, peitschende Bewegung hervorgerufen werden kann. Die eng gewundene Spirale bleibt dabei fast unverändert, ist praformiert. Besonders bei den pathogenen Formen ist die Zahl der steilen Windungen auffallend gross. Eine feine plasmatische Hülle (periplast), die an den Enden in feinste geiszelartige Fortsätze auslaufen kann, ist jedenfalls vorhanden. Die Fortpflanzung geschieht, soweit bis heute festgestellt werden konnte, durch Längs- oder Querteilung.

Winslow *et al* (Committee Soc Am Bact, 1917, p 563) include this as genus four of the family *Spirochaetaceae* with the description

Treponema Schaudinn. Parasitic and frequently pathogenic forms with undulating or rigid spirilliform body. Without crista or columella. With or without flagelliform tapering ends.

The type species is *Treponema pallidum* Schaudinn.

Buchanan (1918, p 544) proposed the following description

Cells slender, spiral, not flattened, attenuated at tips, without crest. Multiplication by longitudinal or by cross division. Parasites in warm-blooded animals. Motile.

The type species is *Treponema pallida* Schaudinn.

Castellani and Chalmers (1919, p 454) include this as the single genus of the family *Treponemidae* *q v*

Bergey *et al* (1923, p 426) include this genus as the fifth in the family *Spirochaetaceae* with the diagnosis "Parasitic and frequently pathogenic forms with undulating or rigid spirilliform body. Without crista or columella. With or without flagelliform tapering end."

Treponemidae. A family name proposed by Schaudinn (1905) according to Castellani and Chalmers (1919, p 454). The latter authors give the following description

Spirochaetacea with a minute threadlike body twisted into numerous fine coils, with pointed tapering extremities. The body is cylindrical on section, and not flattened, and the spirals appear preformed. There is no undulating membrane. Transverse and longitudinal division have been observed, the latter type being the only one found in cultures.

One genus only, *Treponema*, is included.

Trichobacteria. A group name proposed by Messea (1891) and used by many authors for the filamentous bacteria.

Trichobacteriaceae. A family of bacteria named by Fischer (1895, p 138) to include the filamentous bacteria. Fischer's statement follows

Samtliche Bakterien mochte ich zunachst in zwei Orduungen eintheilen, die Haplobakterien, Einzelbakterien, und die Trichobakterien, Fadenbakterien. Das unterscheidende Merkmal liegt in dem Bau des Vegetationskörpers, der bei den Trichobakterien immer ein unverzweigter oder verzweigter aus einzelnen aufgebafter Faden ist, während die Haplobakterien einen einzelligen Vegetationskörper besitzen, der kugelig, stabartig oder schraubig ist. Wenn bei den Trichobakterien die Faden in ihre einzelnen Glieder zerfallen und diese zuschwärmen, so haben wir hier eine Fortpflanzungserscheinung vor aus, die einzelnen Glieder werden zu Schwärmen, zu Gonidien. Der vollendet erwachsene Trichobakterien "im Faden"

The genera included are *Crenothrix*, *Thiothrix*, *Cladothrix* and *Beggiatoa*

The same family name was used by Orla-Jensen (1909, p 303) to designate the typically filamentous water bacteria showing no tendency to the formation of a mycelium. The family is included in the *Cephalotrichinae*, and includes the following genera, *Cladothrix*, *Leptothrix*, *Crenothrix*, *Spirochaeta*, *Spirophyllum*, *Beggiatoa* and *Thiothrix*

Trichobacterinae. An ordinal name used by Fischer (1897, p 32) for bacteria whose "vegetative phase consists of an unbranched or branched filament or chain of cells, the individual members of which break off as swarm spores (gonidia)." The order contains the single family *Trichobacteriaceae*

This name is also used by Schmidt and Weis (1902, p 373) and by Frost (1911, p 59) as a name for a suborder

Trichobakterien A variant (German) spelling of *Trichobacteriaceae* used by Fischer (1895, p 138) and others

Trichogenae A sub-family of *Schizomycetaceae* named by Trevisan (1888). De Toni and Trevisan (1889, p 924) give the following diagnosis

Adsunt tres status evolutionis, h e filamenta (individua), baculi, cocci. Filamenta (status primarius, initialis seu archetypus) tum vaginata tum evaginata, ut plurimum basi ab apice superiori distincta praedita, affixa vel e puncto centrali radiantia, rarius utrinque (apicibus conformibus) libera. Baculi et cocci in filamentis inclusi

This subfamily together with *Baculogenae* and *Coccogenae* constitute the family *Schizomycetaceae*. *Trichogenae* is divided into the following tribes, *Crenotrichae*, *Cladotrichae*, *Kurthiae* and *Leptotrichiae*

Trichomycetes. A group name used by various writers to include various genera of the thread bacteria, probably first by Petruschky (1903). Jordan (1910, p 411) gives the following discussion

The organisms included under this head are filamentous forms and possess many characters relating them to the group of true molds (Hyphomycetes) rather than to the bacteria proper (Schizomycetes). They may be said perhaps to stand in an intermediate position between the bacteria and the higher fungi. All have a more complicated cycle of development than has been observed in any of the true bacteria. On the other hand, they are distinguished from the higher fungi by the slenderness of their filaments and their more simple life-history. Among the so-called Trichomycetes themselves great confusion has prevailed both in classification and in nomenclature, and there is still far from being any universal agreement concerning the subdivision of the group and their designations.

The genera *Leptothrix*, *Cladothrix*, *Nocardia*, and *Actinomyces* are included.

Petruschky (in Kolle and Wasserman, Vol 5, p 267) includes *Actinomyces* and *Streptothrix* under the heading of *Trichomycetes* but places *Cladothrix* and *Leptothrix* with the *Trichobacteria*.

Heim (1911, p 253) has modified the conception somewhat. He says

Der Name ist von J. Petruschky für eine Gruppe von Microorganismen vorgeschlagen worden, die man bisher entweder unter die Hyphomyceten eingereiht oder denen man eine besondere Stelle zwischen diesen und den Schizomyceten zuerkannt hat. Es sind fadenförmige Gebilde mit echten Verzweigungen, die entweder spitz oder mit einer kolbigen Anschwellung endigen und in ihren Breitendurchmesser die Bakterien nicht beträchtlich übertreffen.

Diese Eigentümlichkeiten kommen vielen Bakterien unter gewissen Ernährungsbedingungen zu, darum ist man geneigt, solche, und zwar insbesondere die Tuberkel-, Rotz-, Diphtheriebazillen und ihre Verwandten hierher zu rechnen.

Castellani and Chalmers (1919, p 1040) make this a synonym of *Nocardaceae*.

Trichomyceten A variant of *Trichomycetes* used by Heim (1911, p 253).

Tricogene A variant of *Trichogenae* used by Trevisan (1889, p 9) for the first suborder of bacteria. He states "Tre stati di sviluppo vegetativo 1 Filamenti, 2 Baculi, 3 Cocchi Filamenti, (state primario, iniziale, archetipo) ora rivestiti di vagina propria ambiente, ora non vaginati, per lo più forniti di base distinta dalla estremità superiore, affisi o irradianti da un punto centrale, di rado liberi colle due estremità perfettamente eguali. Baculi e cocci provenienti dai filamenti."

Tyrothrix. A name proposed by Duclaux (1882, p 79) as a generic name for bacteria found in cheese. The first species named is *Tyrothrix tenuis*. The description is summarized by Migula (1900, p. 587 under *Bacillus tenuis*) as follows

Diese Bakterienart bildet kleine, cylindrische, lebhaft bewegliche, im Innern scheinbar fein granulirte, 0 6 μ dicke und 3 0 μ lange Stäbchen, die oft zu langen Fäden auswachsen. Dieselbe ist sporenbildend, ihre Sporen können in alkalischen Nährmedien ein Temperatur von 115°C ertragen. Von besonderem Interesse ist ihr Verhalten in Milch. Vor allem wird das Casein gefällt, das Coagulum ist jedoch weicher als das durch Lab verursachte. Späterhin beginnt in den oberen Schichten die Lösung desselben, ist dieselbe beendet, so zeigt das Ganze eine opaleszierende Flüssigkeit. Duclaux schliesst aus dieser Erscheinung, dass *Tyrothrix tenuis* eine labähnliche, Caseinfallende Substanz (presure) und einen Caseinlösenden Stoff (Casease) abscheidet. Diese Substanz wird von dieser Species in bestimmten Zeitpunkten in sehr grosser Menge produziert und kann durch Alkohol isoliert werden. Weigmann stellte die Casease in Substanz dar und fand, dass dieselbe, frischer Kaseinmasse zugesetzt, die Reifung des Kases ausserordentlich beschleunigt.

Nach Duclaux' Ansicht dient die Casease dazu, um das Casein in eine für die Bakterien assimilierbare Verbindung (Caseone) zu verwandeln. Dieselbe tritt in den Lebenskreislauf der Zelle ein, um in neue Verbindungen (Extraktivstoffe?) verwandelt zu werden. Das Endprodukt der Zersetzung dieser "Caseone" besteht aus einem Gemenge von Ammoniumsalzen, Leucin und Tyrosin. *Tyrothrix tenuis* erzeugt in Milchkulturen speciell valeriansaures Ammoniak. Glycerin wird von demselben vollkommen verbrannt und der milchsaure Kalk in Calciumcarbonate verwandelt. Die nachfolgenden Tyrothrixarten liefern auch Casease, jedoch nicht in solcher Menge wie *Tyrothrix tenuis*.

In addition to this species, six other aerobic forms were described, *Tyrothrix filiformis*, *T. distortus*, *T. geniculatus*, *T. turgidus*, *T. scaber* and *T. virgula*, also three anaerobes *T. urocephalum*, *T. claviforme* and *T. catenula*. The various species described have little in common other than origin.

Maggi (1886, p 17) listed Duclaux's species with brief descriptions, including the genus as valid in his classification.

De Toni and Trevisan (1889, p 963) renamed *Tyrothrix tenuis* as *Bacillus tenuis* (Duclaux). Trevisan. The other species are distributed in *Bacillus* and *Cornelia*.

Macé states (1897, p 486)

Duclaux (2) a groupé sous la designation de *Tyrothrix*, sans toutefois vouloir lui attribuer une valeur generique, toute une serie d'interessantes especes qu'il a rencontrees dans les fermentations des albuminoides, en particulier de la caseine du lait. Ces *Tyrothrix* appartiennent tous au genre *Bacillus*, tel que nous le comprenons ici.

Migula (1900, II, pp 322-588) has apportioned some ten species between the genera *Bacillus* and *Bacterium*. The genus name is rejected by E F Smith (1905, p. 174)

Lehmann and Neumann (1907, p 396) make the following statement

Das Genus *Tyrothrix* Duclaux fällt mit *Bacillus* zusammen, es bezeichnet ursprünglich aus Milch und Kase stammende, sporentragende, längere Faden bildende Arten. Zwei species sind unten beschreiben. Die *Tyrothrix* arten sind neuerdings von Niede (C B L XII 344) mit einem von ihm genau beschriebenen *Bacillus parvus* für synonym erachtet worden

The lack of adequacy in the original description make determination of the nomenclatural status of this genus difficult. It is probably a synonym of *Bacillus* Cohn. The type species may be designated as *Tyrothrix tenuis*.

Ulvina. A generic name applied as *Ulvina aceti* to mother of vinegar by Kuetzing in the 11th decade of his *Algae aquae dulcis* (1837). In the same year (1837, p 26) the organism is mentioned by name and later (p 30) illustrated.

Later (1843, p 149) the following generic description is given: "Stratum compactum lubricum ex granulis minutissimis compositum". Four species are listed, *U. myxophila* from sour quince slime, *U. rubridaei* from fermenting raspberry juice, *U. aceti* from mother of vinegar, and *U. sambuci* from the aqua sambuci flor., of the apothecary.

Naegeli (1849, p 5) mentions *Ulvina* as a genus to be included among the fungi and excluded from the unicellular algae.

In his *Species Algarum* (1849, p 147) the generic description is "Globuli gonimici minutissimi in stratum compactum cohaerens fluitans coaliti". The same four species previously described are listed. *Ulvina aceti* is again mentioned in 1852, (p 269).

Rabenhorst (1864, p 8) in a footnote under Subfamilia II, *Leptothricheae* says

Fungorum genera *Leptothricheae* et *Oscillariens* quasi affinia

a *Cryptococcus* Cellulae sphaericae, in stratum mucosum aggregatae

b *Ulvina* Cellulae minutissimae, in stratum membranaceum coalitae

According to De Toni and Trevisan (1889, p 1021) (*g v*) the *Ulvina aceti* was termed *Mycoderma aceti* by Pasteur (1868, p 106), *Bacterium aceti* by Lanzi (1876, p. 257), *Torula aceti* by Saccardo (1878, p 315), *Bacillus aceti* by Schroeter (1886, p 161).

Migula (1900, p 399) gives an incorrect reference to the name *Ulvina* (Kuetzing 1837, p 385)

The name has not been used by recent writers It is rejected by E F Smith (1905, p 174)

It is probable that *Ulvina* has priority as a separate generic name for the acetic acid bacteria. However, the author of the name undoubtedly was dealing with mixed cultures of such a nature that a reasonably certain identification of his species (type) *Ulvina aceti* is probably not practicable To eliminate this difficulty, Winslow *et al.* (Committee Soc Am Bact, 1917) have accepted the later name *Acetobacter*

Umbina. According to Migula (1897, p 10) this name was used by Naegeli (1848) (should read 1849) in the combination *Umbina aceti* for the organism forming mother of vinegar

An examination of Naegeli's work shows no use of this name, but instead (p 5) a statement that *Ulvina* belongs with the unicellular fungi and not the algae

In 1857 (p 760) he includes *Umbina aceti* with *Nosema*, *Bacterium*, *Vibrio*, *Spirillum*, *Hydrococcus* and *Sarcina* in his new group, *Schizomycetes* He says "*Umbina aceti* die Essigmutter, ist mit *Nosema* sehr verwandt, jedoch fallen die Zellen nicht auseinander, sondern bleiben vereinigt "

Umbina is a synonym of the older *Mycoderma* and *Ulvina*

Urobacillus A physiological genus proposed by Miquel (1889, p 517) for certain rod shaped bacteria found in decomposing urine The species described by Miquel were for the most part spore-bearing motile rods The first species described is *Urobacillus Pasteuri* and may be designated (Enlows, 1920, p 96) as the type

The name has not apparently been recognized by many authors as a true generic designation It is used by Beijerinck (1901, p 40) also (1902, p 30) It is rejected by E F Smith (1905) as invalid It is proposed as the third genus of the family *Alkalibacteriaceae* by Orla-Jensen (1909, p 339) He says,

Einen besonderen Zweig der aeroben Bacillen bildet die Gattung *Urobacillus* Diese ist jedoch ohne Zweifel desselben Ursprungs wie die Gattung *Bacillus*, denn die meisten *Proteus* Bakterien (auch *Lb prodigiosum*) hydrolysieren den Harnstoff Diese Fähigkeit wird auch bei einzelnen Coli-Arten, vielen Kokken und den meisten anaeroben Faulnisbakterien angetroffen und kann deshalb nicht für sich allein als ein Gattungsmerkmal dienen Eine Urobakterie ist erst dann typisch, wenn sie sich derart an die durch die Hydrolyse des Harnstoffes hervorgerufene hohe Alkalität gewöhnt hat, dass sie für die gewöhnlichen Nahrsubstrate

zu säureempfindlich geworden ist. Die ausgesprochensten Urobakterien verflüssigen die Gelatine nicht, denn sie bedürfen keiner anderen Stickstoffquellen als des normalerweise reichlich vorhandenen Harnstoffes, und proteolytische Enzyme wurden wegen der gebildeten grossen Ammoniakmenge überhaupt nicht wirken können. Die Entwicklung zu den, was sonst der Fall ist—von gelatineverflüssigenden Formen. Nach Miquel sollen die wichtigsten stabchenförmigen Urobakterien die folgenden sein:

Nichtsporenbildend	$\left\{ \begin{array}{l} \text{Urobacillus Miqueli} \\ \text{Urobacillus Schutzenbergii I} \\ \text{Urobacillus Schutzenbergii II} \\ \text{Urobacillus Freudenreichii} \end{array} \right\}$	Gelatineverflüssigend
Sporenbildend	$\left\{ \begin{array}{l} \text{Urobacillus Leubei} \\ \text{Urobacillus Maddoxii} \\ \text{Urobacillus Pasteurii} \\ \text{Urobacillus Duclauxii} \end{array} \right\}$	Nichtgelatineverflüssigend

Da die drei ersten Arten keine Sporen bilden, sind sie laut unserer Definition—einfach den Proteus-Bakterien zuzuzählen, *Liquidobacterium Miqueli* bildet auch in der Gelatine die für diese Gattung charakteristischen bildende Art, *Urobacillus Freudenreichii*, und sie nähert sich dadurch wie auch durch die Gelatineverflüssigung den ältesten Arten der Gattung *Bacillus*.

Urobacillus is probably best regarded as a synonym of *Bacillus*, or possibly as a subgenus.

Urobacter. According to Cohn (1875, p. 188), Trecul used the name *Urobacter* as the name of a bacterial genus. The original paper has not been found. Cohn says "Ebenso hatte Trecul in den geschwanzten Bakterien eine selbständige Gattung erblickt, die er *Urobacter* nannte."

The name is rejected by E. F. Smith (1905, p. 174).

Urocephalum. A name used by Trecul (1865, p. 432) to designate certain granules of uncertain affinities found in decaying plant cells. The name has apparently never been used in a strict generic sense, although Trecul undoubtedly saw bacteria. He says:

Malgré la variété des formes de ces petits végétaux, ou plutôt à cause de cette variété même, puisque l'on passe d'une forme à l'autre par toutes les gradations, en raison aussi de l'amidon qu'ils contiennent, et pour rappeler la ressemblance des formes cylindriques avec les Bactéries, je crois utile de les réunir sous le nom d'*Amylobacter*. Si on voulait les diviser d'après les formes les plus disparates, on pourrait établir trois sous-genres: 1° l'*Urocephalum*, qui comprendrait les formes en têtard, 2° l'*Amylobacter* vrai, auquel seraient attribuées les formes cylindriques, 3° le *Clostridium*, qui renfermerait les formes en fuseau.

What Trecul observed were probably butyric acid bacteria. Migula (1897, p. 324) gives this as a synonym of *Clostridium butyricum* Prazmowski.

The name is rejected by Erwin F. Smith (1905, p. 174)

Urococcus. *Urococcus* was proposed as a generic name by Kuetzing (1849, p. 206), for a group of algae. This author ascribed the name to Hassall (1845, p. 322) but this latter author used the name *Ourococcus* as a subgeneric designation under the genus *Haematococcus*.

Miquel (1888, p. 518) used this as a generic designation for cocci bringing about ammoniacal fermentation of urea.

Bejerinck (1902, p. 37) says

Pour reconnaître et distinguer les urobactéries, M. Miquel les a classées en coques qu'il désigne par le nom d'*Urococcus*, et en bacilles qu'il appelle *Urobacillus*. Dans le genre physiologique *Urococcus* il classe 9, dans le genre *Urobacillus* 8 espèces.

The description and designation of species by Miquel have not been found. However, Macé states (1912, p. 590) that the *Micrococcus ureae* Van Tieghem is probably the *Urococcus Van Tieghemi* Miquel.

Bejerinck (1901, p. 54) renamed the *Micrococcus ureae* Cohn as *Urococcus ureae*.

The nomenclatural status of the genus and the designation of type species are uncertain.

Urosarcina. A generic name used by Miquel (1888, p. 517) for a motile sarcina form in urine. It is listed as a synonym of *Sarcina* by De Toni and Trevisan (1889, p. 1044) but no species is listed in which the name was used. Bejerinck (1901, p. 43) states that this organism is a bacillus. It is rejected as a generic name by E. F. Smith (1905, p. 174).

Vallorillus. A generic name proposed by Heller (1922, p. 16) as the seventh in the *Clostridiaceae*. The description given is

Clostridiaceae that do not liquefy gelatin. They produce gas and acid in meat medium but no digestion. They clot milk slowly and attack various sugars. Rather slender Gram-positive rods with little or no tendency to form spores. Form lenticular colonies, "coeurs jaunes," in deep agar. Many invade tissue, producing oedema and gas. Pathogenicity transitory.

Type species *V. fallax* (*Bacillus fallax* Weinberg and Séguin) as described by the Committee (p. 27), the type which ferments glucose, laevulose and maltose.

Vibrio. A generic name proposed by Muller (1773, p. 39) in the group *Infusoria crassiuscula* with the generic description "Vermis inconspicuus simplicissimus, teres elongatus." Other descriptions of the genus as taken from the keys are "*Tentaculis destitute, vagantes*,

organis externis nullis, teretes, corpus elongatum," "Infusoria" "Aquatici," "Hyalini," "Gelatinosi, Vibrionis species" "Tentaculis nullis, inconspicuis, teretes" "Vacillantes pellucidi" The species first described is *Vibrio lineola*, which would thereby become the type. Other species named are *Vibrio bacillus* and *Vibrio undula*.

Mueller (1786, p. 43) described *Vibrio lineola* as follows:

Linearis minutissimus. Animalculum omnium minutissimum, monadem terminem exiguitate, fere superans, Vibrioneque Bacillo tricies minus et prorsus diversum. Motus tremulus myriadum punctulorum oblongorum obscuriorumque in unica guttula, seu undulatio oculo, lenticula maxime amplificante, exhibetur. In infusione vegetabili substantiam aquae post plures dies fere adimplet, in alia foetente ultra trimestre servata, et in non foetente post mensem Lemna cooperta cum Cycloidio glaucomate.

Other species named are *Vibrio rugula*, *V. bacillus*, *V. undula*, *V. serpens* and *V. Spirillum* which quite certainly were forms now included among the bacteria. Altogether some 31 species were described. The fact that the first named organism probably belonged to the bacteria, though many of the others were protozoa, makes its status in bacteriology a matter of concern.

Two lines of study are necessary in determination of the present nomenclatural status of this genus. First, it is of interest to note the gradual change in the concept of this genus, and second, to trace the fate of Muller's type species.

Bory de St. Vincent (1826, p. 533) included *Vibrio* as the third genus of his family *Vibrionides*. The organisms are described as linear or worm-like, body diaphanous, attenuate at both tips.

Ehrenberg (1828, p. 24) gave the following generic diagnosis: "Animal Polygastricum, Anenterum, elongatum, monomorphum (contractione non incrassatum), transverse in multas partes sponte dividuum, ore terminali, corpore filiformi cylindrico undatum flexili." He included two species of Mueller, *V. lineola* and *V. rugula*.

Ehrenberg (1838) split this genus into three, *Vibrio*, *Spirillum* and *Spirochaeta*. *Vibrio* included the organisms which are straight rods, but are flexuous when in motion. *Spirillum* included the spiral non-flexuous forms, and *Spirochaeta* the spiral flexuous types. He described six species in the genus *Vibrio*, *V. lineola*, *tremulans*, *subtilis*, *rugula*, *prolifer* and *bacillus*. In 1840 he named two organisms which can be recognized even today, *Vibrio syncyaneus* from blue milk and *V. synzanthus* from yellow milk.

Fuchs (1841, p 190) also described *Vibrio cyanogenus* and *V xanthogenus*

Dujardin (1841, p 216) characterized the genus as follows "Corps filiforme, plus ou moins distinctement articulé par suite, d'une division spontanée imparfaite, susceptible d'un mouvement ondulatoire comme un serpent" The family of *Vibrionia* he characterized as "Animaux sans organes locomoteurs visibles, se meuvent par l'effet de leur contractibilité générale" Four species of *Vibrio* are recognized, *V Lineola*, *rugula*, *bacillus* and *serpens*

Leidy (1849, p 228) was apparently the first to suggest the placing of this genus among plants rather than animals He discussed two species which he believes belong to the genus *Vibrio*, from the coecum of *Julus* He says

Even those moving filamentary bodies belonging to the genus *Vibrio*, I am inclined to think, are of the character of algaes vegetation Their movement is no objection to this opinion, for much higher confervae, as the *Oscillatorias*, are endowed with inherent power of movement, not very unlike that of the *Vibrio*, and indeed the movement of the latter appears to belong only to one stage of its existence Thus, in the toad (*Bufo americanus*), in the stomach and small intestine, there exist simple, delicate, filamentary bodies, which are of three different kinds One is exceedingly minute, forms a single spiral, is endowed with a power of rapid movement, and appears to be the *Spirillum undula* of Ehrenberg, the second is an exceedingly minute, straight and short filament, with a movement actively molecular in character, and is probably the *Vibrio lineola* of the same author, the third consists of straight, motionless filaments, measuring 1/1125th inch long, by 1/15,000th broad, some were, however, twice, or even thrice this length, but then I could always detect one or two articulations, and these, in all their characters, excepting want of movement, resemble the *Vibrio* In the rectum of the same animal, the same filamentary bodies are found with myriads of *Bodo intestinalis*, but the third species, or longest of the filamentary bodies, have increased immensely in numbers, and now possess the movement peculiar to the *Vibrio lineola* which, however, does not appear to be voluntary, but reactionary, they bend and pursue a straight course, until they meet with some obstacle, when they instantly move in the opposite direction, either extremity forward "

Perty (1852) divides the family *Vibrionida* into two groups, *Spirulina*, containing organisms in chains or filaments spirally wound and *Bacterina* in which the filaments are flexuous or straight In the group *Bacterina* three genera were included, *Vibrio*, *Bacterium* and *Metallacter* He retained only two species in *Vibrio*, *V rugula* and *V lineola* but recognized *V syncyanus* and *V synxanthus* Davaine (1859) used practically the same definition of the genus

According to Trevisan (1885, p 83), Pacini (1854, p 40) used the designation *Vibrio cholerae* for the organism causing Asiatic cholera, but a search of the original paper fails to show it. The nearest approach is the following statement:

Finalmente notero che nel fluido intestinale trovasi ancora una grandissima quantità di *vibroni*, i quali, attesa la loro estrema tenuità, possono facilmente passare inosservati, quando siano dispersi in una certa quantità di fluido. Questi vibroni avevano una lunghezza di 0.0020 a 40^{mm}, ed un diametro di 0.0005 a 7^{mm} mentre avevano qualche simiglianza al *Bacterium termo* di Dujardin.

Cohn (1854) concluded the *Bacterium termo* of Dujardin was the *Vibrio lineola* of Ehrenberg, and as a result of his study of the organism, placed it in the plant kingdom.

Rabenhorst (1865, p 71) gives the following generic diagnosis: "Trichomata abbreviata filiformia, plus minus distincte articulata, motu proprio, plerumque flexuoso praedita, cytoplasma pallidum, interdum dilute seruginosum, coeruleum vel caesium." He lists seven species, three described by Ehrenberg and four by Mueller.

Cohn (1872, p 178) emended the definition of the genus. The organisms are characterized by the wavy motion of the filaments, which by their rotation give the appearance of sinuous motion, and constitute a transition to the spiral bacteria. The two species named are *Vibrio rugula* and *V. serpens*.

Cohn (1875) included the genus in his tribe *Nematogenes*. Luerssen (1879, p 25) used the same definition and species. Trevisan (1879, p 148) gave the following definition: "Somatia cylindrica, articulata, valida, abbreviata, undulato-flexuosa." The two species listed are *V. serpens* and *V. rugula*. This genus is included in the subtribe *Eumembrionaeae*. Magnin (1880, p 92) lists the same species. Lehmann and Neumann (1901) credit *Vibrio cholerae* to Buchner (1884). Van Tieghem (1884) defined the genus as made up of spiral cells, promptly dissociating. Zopf (1885, p 61) states "Faden schraubig, in den längeren oder kürzeren Gliedern Sporenbildung." Trevisan (1885, p 83) replaced this with *Pacima* with Koch's comma bacillus as the type. Hueppe (1886) included forms with spiral cells, which produce endospores and change the cell form when spores are produced. Flugge (1886) states that this genus differs from *Spirillum* in that the spirals are not as pronounced. Maggi (1886, p 176) included *Vibrio* as the first genus of his order *Ophidobacteries*. Schroeter (1886, p 166) introduced *Microspira* with Koch's comma bacillus as the type. Hansgirt (1888, p 264) makes this the first subgenus of the genus *Spirillum*.

with the definition "Zellen farblos, meist ohne grossere Schwefelkornchen in plasmatischen Inhalte" De Toni and Trevisan (1889, p 1005) give the following diagnosis

Baculi cylindrici, leviter curvuli v undulatoflexuosi (nunquam spirales), plasmate uniformiter diffuso instructi Sporae (endosporae) macrosomae, ovoideae in baculorum specialium apice claviformi-intumescenti ortae Baculi speciales, qui ad sporas ferendas praeparantur, fiunt breviores, crassiores, plasmate tenuissime granuloso

They list as species *Vibrio rugula*, *V. urinae*, *V. amphibolus* and *V. typhosus*

Billet (1890, p 24) defined the genus as made up of curved organisms with a single curve

Sternberg (1892, p 18) makes the following statement

Filaments flexible, straight or sinuous, movements sinuous A considerable number of bacteria which are usually seen as short, curved rods, but which may grow out into long, spiral filaments, are described by some authors under the generic name *Vibrio*, e g, the so-called "comma bacillus" of Koch—"Spirillum cholerae Asiaticae," the spirillum of Finkler and Prior—"Vibrio proteus," the spirillum described by Gameleia—"Vibrio Metschnikovi," etc These microorganisms have not the characters which distinguished the genus *Vibrio* as established by Ehrenberg, and we prefer to follow Flugge in describing them under the generic name *Spirillum*

Among those referred to by Steinberg who used *Vibrio* for the short spirals are Firtsch (1888, p 369) who discussed *Vibrio proteus* (the *Spirillum* of Finkler and Prior), Gameleia (1888, p 482) According to De Toni and Trevisan (1889, p 1018) the name *Vibrio cholerae* was used by Pacini (1854) Pouchet (1849) five years before had observed vibrios in cholera stools It is by no means certain, however, that these observers actually saw the true comma bacillus, but it is not improbable

Lehmann and Neumann (1896, p 316) give the following description

Zellen kurz, schwach bogig, starr, kommaartig gekrümmt, zuweilen in schraubenartigen Verbanden an einander hangend, stets nur mit einer, ausnahmsweise 2 endständigen Geisseln Endosporen fehlen, nach Huppe Bildung von Arthrosporen

Macé (1897, p 485) makes the following statement concerning the use of *Vibrio*

Le genre *Vibrio* d'Ehrenberg etait trop peu homogene pour être conserve Le plus grand nombre de ses espèces sont de vrais *Bacillus*, le restant des *Spirillum*

Il en reste la denomination française de Vibrion, que l'on ne doit pas considerer comme un terme de classification, mais comme une simple expression de valeur generale Elle a été tres employee par l'école de Pasteur, qui l'a appliquee a bien des Bactéries mobiles, *Micrococcus* ou *Bacillus* Le Vibrion pyogene est un *Micrococcus*, le Vibrion lactique, le Vibrion butyrique, le Vibrion septique, sont des *Bacillus*

Fischer (1897, p 32) ascribes an emendation of *Vibrio* to Loeffler He includes in the genus the very slightly curved rods which are "comma" shaped, motile, monotrichous, with the type species *Vibrio cholerae-asiaticae* Lehmann and Neumann (1901, p 126) use this same definition These authors credit the name *Vibrio cholerae* to Buchner (1884) Migula (1897 and 1900) follows Schroeter (1886) in the use of *Microspira*.

Matzuschita (1902) defines a *Vibrio* as a short spiral

E F Smith (1905, p 173) agrees that the original description of *Vibrio* by Muller does not define the organism, and accepts Loeffler's emendation of *Vibrio* with Koch's comma bacillus as the type

Blanchard (1906, p 1) credits *Vibrio* to Ehrenberg (1838) with *V comma* as the type

Flügge (1908) includes in *Vibrio* all comma shaped organisms with polar flagella, all gram negative

Vuillemin (1913, p 518) concludes that *Vibrio* as a generic name should be suppressed because of its varied uses and believes that if the short spirals are worthy of generic recognition, the name *Microspira* should be used

Lohns (1913) accepts the definition of *Vibrio* as a comma shaped rod

Winslow *et al* (Committee Soc Am Bact, 1917, p 556) used the following definition, including *Vibrio* as the first genus of *Spirillaceae*

Cells short bent rods, rigid, single or united into spirals Motile by means of a single (rarely two or three) polar flagellum, which is usually relatively short Many species liquefy gelatin and are active ammonifiers Aerobic and facultative No endospores Usually Gram-negative Water forms, a few parasites

The type species is *Vibrio cholerae* (Koch) Buchner

Buchanan (1918, p 178) followed the committee, using the following diagnosis

Short, bent rods, sometimes almost straight Motile by means of a single (rarely 2 or 3) polar flagellum Aërobic, and facultative Grow well on ordinary

media Frequently liquefy gelatin Not enlarged near center No spores
Usually Gram-negative

The type species is *Vibrio cholerae*

Castellani and Chalmers (1919, p 962) state

Spirillaceae motile, with short comma-like cells, possessing one, rarely more, flagellum at one, rarely at both ends

Type It is difficult to decide which is the type of this genus so defined

Remarks The important species is the *Vibrio comma* Koch, 1884, which is the cause of Asiatic cholera, but there are many others which cause paracholera

Bergey *et al* (1923, p 77) followed Castellani and Chalmers

It is apparent that the modern use of the term *Vibrio* is not consistent with the conception of *Vibrio lneola* as the type Apparently either the genus should be dropped as having a type species that cannot be identified, or there should be an agreement that *Vibrio cholerae*, a somewhat arbitrary choice, shall be selected as the type The latter alternative is the one here adopted

Vibron. A casual name for bacteria apparently first used by Pasteur (1876) in the phrase "vibron septique" Not a generic name

Vibronia. A family of protozoa used by Ehrenberg (1838) to include the genera *Bacterium*, *Spirillum*, *Vibrio*, *Spirochaeta* and *Spirodiscus*

Perty (1952, p 179) used this as the single family of his Sectio Lamprozoidia It contained two subfamilies *Spirillina* and *Bacterina*

It has also been used in a sense equivalent to "bacteria" (Quart Journ Mic Sci, 1855, p 205)

Vibronides. The sixth family of the first order *Gymnodes* of Boiy de St Vincent's (1826, p 533) group Microscopique

The order *Gymnodes* is described as without tests, perfectly smooth, without cilia or other vibratile organs The family is characterized as without appendages, linear or wormlike Five genera are included, *Spirillina*, *Melanella*, *Vibrio*, *Lacrumatoria* and *Pupella*

Vibronieae. A tribe of bacteria created by Trevisan (1879, p 137) with the following description

Bacteriaceae pluricellulares Individua e cellulis pluribus in somatia saepe filiformia dispositis confiata Sporae intra somatium matricale uniseriales Multiplicatio divisione somatorum vegetativa semper ad eandem directionem longitudinalem

It was divided into the subtribes *Euvibronieae*, *Myconostoceae* and *Cladothriceae*.

Vibroniens. The single family of the first order of the *Infusoria* named by Dujardin (1841, p 209) The diagnosis given is "Animaux filiformes extrêmement minces, sans organisation appreciable, sans organes locomoteurs visibles" Three genera are included *Bacterium*, *Vibrio* and *Spirillum*

Vibriothrix. A generic name proposed by Castellani (1917) Castellani and Chalmers (1919, p 1068) give the following description

The mycelial articles are motile, of very different shape bacillary, vibrio-like, spirillum-like, at times club-ended Globular or pear-shaped bodies of very variable size may be present Gram-negative, not acid-fast Cultivable on ordinary media

Type species *Vibriothrix zeylanica* Castellani, 1910

Synonyms *Spirillum zeylanicum* Castellani, 1910, *Vibrio zeylanicus* Castellani, 1913, *Bacillus zeylanicus* Castellani, 1913, *Vibriothrix zeylanica* Castellani, 1917, *Spirobacillus zeylanicus* Castellani, Spagnolo, and Russo, 1918

Remarks Very polymorphic organism vibrio-like, bacillus-like, and undulating forms being often found in the same preparation Very small, medium size, and occasionally large roundish bodies are at times observed, and club-like forms may also be present

The organism is motile, Gram-negative, not acid-fast Easily grown on ordinary media On potato the growth is often of a reddish colour In broth there is often a pellicle, preparations from the fluid medium generally show a predominance of vibrio-like or bacillary forms, while in the pellicle long undulating forms are often found On MacConkey's medium the colonies are white, and somewhat resemble those of the typhoid-dysentery group The organism does not ferment any of the usual laboratory carbohydrates or alcohols glucose, levulose, galactose, maltose, lactose, saccharose, mannitol, dulcitol, raffinose There is, in fact, frequently a production of alkalinity Milk is not clotted and is rendered alkaline, and certain strains after several weeks may induce a certain degree of peptonization The great majority of strains are non-pathogenic to rabbits and guinea-pigs

Weinbergillus. A generic name proposed by Heller (1922, p 31) with the following description

Highly proteolytic *Putrificoideae* that in meat medium cause the formation of balls of amino-acid crystals They digest the meat particles till their bulk is greatly reduced and form large amounts of ammonia, amino-nitrogen and fatty acids They digest the casein in milk rapidly Attack few or no sugars Produce little or no gas in agar media Gram-positive or Gram-negative rods with sub-terminal oval spores Colonies in deep agar small, delicate woolly structures May invade living tissue in company with other organisms, or at times alone, producing a rapid and complete digestion of muscular and connective tissue structures

Type species *W. histolyticus* (*Bacillus histolyticus* Weinberg and Séguin) as described by Henry (p 370) Weinberg and Séguin and the Committee allow con-

siderable variation in characters for the strains termed by them *Bacillus histolyticus*

Welchillus. A generic name proposed by Heller (1922, p 18) with the following description

Clostridiaceae that liquefy gelatin but do not produce hydrogen sulfid demonstrable by lead-acetate-paper test in blood-broth They produce much acid and gas on meat medium, but they do not digest it, nor do they digest casein, coagulated serum, or egg-white They produce stormy fermentation of milk and attack many sugars vigorously Their multiplication is extraordinarily rapid, they are killed by their own growth products in acid media, in which they fail to sporulate Deeply Gram-positive non-flagellate rods with square ends They produce oval subterminal or median spores in alkaline media, or in media free of fermentable carbohydrates, these spores do not bulge the sides of the bacilli When growing rapidly the rods are very short, and resemble closely no other type here listed except *Stoddardillus* When growing slowly the rods are less abundant and longer, and may be mistaken for *Martellillus* and similar organisms Typically intestinal saprophytes, ubiquitous Many strains produce toxin and invade tissue, forming gas, and causing the formation of oedema, and in many cases causing the disintegration of muscle and of connective tissue without the production of a foul odor This disintegration occurs only *in vivo*, and is probably due to the enzymes of the host tissue *Welchillus* are the most frequent anaerobic invaders noted on the hospital autopsy table The organisms are present in human intestines Causative agents of a probably greatly overestimated percentage of gas gangrene cases following war wounds Comparatively rare as animal invaders

Type species *W. aerogenes* (*Bacillus aerogenes-capsulatus* Welch and Nuttall) type I as defined by Simmonds (type IV of Esty) Ferments besides other carbohydrates, inulin and glycerol Usually pathogenic for guinea pigs

Wesenbergus A generic name proposed by Castellani and Chalmers (1919, p 934) for the eighth genus of the tribe *Ebertheae* with the following description

Ebertheae which ferment glucose completely and lactose partially producing acid, but no gas Milk not clotted

Type species *Wesenbergus wesenbergi* Castellani, 1913.

Winogradskya A generic name proposed by Trevisan (1889, p 12) De Toni and Trevisan (1889, p 1028) give the following description "Baculi cylindracei et filamenta aggregata in familias zooglaeicas repetite ramosas, capsula tenui gelatinosa inclusa Sporae ignotae"

One species is included *Winogradskya ramigera* (Itzig) Trev. This is the *Zoogloea ramigera* Itzigsohn

The genus is probably not tenable

Zoagalactina. A variant of the generic name *Zoagalactina* used by Sette (1824, p. 51) for an organism which in 1819 caused red spots to appear in various food stuffs such as bread, macaroni, rice, etc. Trevisan (1887, p. 278) says:

Vincenzo Sette delegato d'ufficio ad investigare le cause del fenomeno che il fanatismo del popoli attribuiva ad un potere sopranaturele, mise in chiara pel primo che la cagione unica n'era un frungo nuovo di genere e di specie, cui impose il binomio *Zoagalactina imetropa*. Che se ventiquattro annipiu tardi Ehrenberg erroneamente riferendo la *Zoagalactina* de Sette alle Monadi tragli Infusorj, l'appello *Monas prodigiosa*, non per questo, per la volgare legge id prioritato potrebbe essere posposto il piu antico nome specific; und eche come vel 1879 in quest 'aula medesima proposi che aversse a chiamarsi *Micrococcus imetrophus* d'ora umana dovrebbe dirsi *Bacillus imetrophus*.

The priority of *Zoagalactina*, or rather its definition was questioned by Schroeter (1872, p. 109). He says:

Sette beschrieb die rothen Schleimklumpschen schon 1824 als *Zoagalactina imetropa*, sah aber die einzelnen, sie constituirenden Organismen nicht, deshalb kann dieser Name keine Priorität für die Beziehung der Körperchen beanspruchen.

He further states that he has found in potatoes red slimy drops which were not bacteria, but yeasts. He contends that Sette's type is therefore not recognizable and should be abandoned.

The name *Serratia* (q.v.) evidently has priority.

Zoagalactina. Enlows (1920, p. 98) states that this is the original spelling for *Zoagalactina* (q.v.).

Zoogalactina. A variant spelling of *Zoagalactina* used by Schroeter (1872, p. 109) and others.

Zoogloea. A generic name proposed by Cohn (1854, p. 123). He gave the following description:

Cellulae minimae bacilliformes hyaline, gelatina hyalina in massas globosas, uvaeformes, mox membranaceae consociatae, dein singulae elapae, per aquam vacillantes. The species named was *Zoogloea termo*.

Itzigsohn (1867) named an organism *Zoogloea ramigera*. Bollinger (1870, p. 583) used the name *Zoogloea pulmonis equi*. Billroth (1874, p. 6) proposed that the terms *Coccoglia* or *Gliacoccus* should be substituted for *Zoogloea*. Eberth (1873) named an organism *Zoogloea Beigeliana*. Cohn definitely abandoned *Zoogloea* (1872b, p. 141) as a generic designation believing that this was simply a growth form.

Since that time the name has not been used except as a casual name Vuillemin (1913, p 526) designates this as a form genus to be preserved

The generic name is probably invalid because not based upon pure culture

Zopfiae. A name proposed by Winslow *et al* (Committee Soc. Am Bact, 1920, p 209) for the third tribe of the family *Bacteriaceae* with the following description "Gram-positive rods, growing freely on artificial media Not attacking carbohydrates"

The single genus is *Zopfius*

The tribe is included by Bergey *et al* (1923, p 193) as the fifth in the family *Bacteriaceae*

The correct spelling of this name probably is *Zopfiae*

Zopfiella. A genus proposed by Trevisan (1885, p 4) with the species *Zopfiella tumescens*, (*Bacillus tumescens*). The description given is as follows

Tre stadi di sviluppo vegetativo 1 Filamenti 2 Bacilli 3 Cocchi
Filamenti (stadio protoplastico tipico) cilindrici, articolati, micolori, dritti
Bacilli (stadio transitorio) cilindrici, inarticolati, filiformi, macrobacilli e microbacilli
Cocchi (stadio finale) derivati da microbacilli, d apprima in brevi monili, indo liberi Spore provenienti

The name was later abandoned (De Toni and Trevisan, 1889, p 972) becoming a synonym of *Bacillus*

Zopfius. A generic name proposed by Wenner and Rettger (1919, p 334) The generic description is summarized by Enlows as follows

Cells rod-shaped, usually about 0.8 by 3.5 μ in size, have somewhat rounded ends, and in young cultures occur in long evenly curved chains Gram-positive Motile by means of peritrichiate flagella No spores No capsules Facultative anaerobes No visible change in litmus milk Gelatin not liquefied and none of the carbohydrates are attacked A more or less characteristic spider-web growth on agar and gelatin plates, but inoculations in the condensation water of agar slants do not result in a spreading over the surface such as occurs in the genus *Proteus* The authors include here *Bacterium zopfii* Kurth, and *Proteus zenkeri* Hauser, which they regard as identical, after a study of a number of strains of each type finding few differentiating properties See *Proteus* (Hauser) em Wenner and Rettger The name *Zopfius* having been chosen as the name for new genus, the type species (by virtual tautonymy) would be *Zopfius zopfii* (Kurth) Wenner and Retter

Winslow *et al* (Committee Soc. Am Bact, 1920, p 210) include this as the single genus of the tribe *Zopfiae*, with the following characterization.

Long rods occurring in evenly curved chains Gram positive Motile Proteus-like growth on media Facultative anaerobes Carbohydrates and gelatin not attacked, hydrogen sulphide not formed

Type species, *Z. zopfii* (Kurth) Wenner and Rettger

Bergey *et al* (1923, p 193) include this genus in the tribe *Zopfiae* Apparently this name is an exact synonym of *Kurthia* which has priority Both are based on the same type species

Zygobacterium. According to De Toni and Trevisan (1889, p 1023) this generic name was used by Maggi (1887, p 318) for the species *Zygobacterium nitrosum* which is regarded by the former as a synonym of *Bacterium lineola* (Mueller) Cohn

Zymosotis. A generic name proposed by Salisbury (1868) Enlows (1920, p 99) gives the following summary

Z. regularis Spores very minute well defined in outline, and uniform in size and shape Multiply by duplicate segmentation, and develop into filaments with great rapidity Filaments are well defined, uniform in diameter, and have cross markings or interruptions in the inside tubular membrane at regular intervals, hence its name Found in human blood

Z. escularis The filaments in their early stages of development are mostly moniliform The more mature filaments have the outside tube continuous and uniform in diameter, while the inside membrane has not only interruptions at irregular intervals, but the interruptions are of variable length Where the inside membrane occurs it affords a double wall to the tube and communicates greater opacity than have the intervening spaces Found in the freshly drawn blood of horses affected by a fatal disease characterized by a remittent fever

BIBLIOGRAPHY

- AFANASSJEW, M J 1887 Die Aetiologie und klinische Bakteriologie des Keuch-
hustens St Petersburg Med Wochenschrift 4 323, 331, 339, 347
- AGARDE, C A 1821 Species Algarum Vol 1
1828 Icones Alg Eur Nos 21-23
- AGARDE, J G 1824 Syst Alg
1842 Algae Maris Mediterranei et Adriatici p 154
- ALESSAND, P E, e MAGGI, L 1887 Acque potabili
- ALTANA, GIUSEPPE 1909 Ueber einen vom Meerschweinchen isolierten Tetra-
genus Centralbl f Bakteriologie, Jena Abt 1 48 44
- AMBROŽ, ADOLF 1913 *Denitrobacterium thermophilum* spec nova, ein Beitrag
zur Biologie der thermophilen Bakterien Centralbl f Bakteriologie, Jena
Abt 2 37 3-16
- AMIRADŽIĆ, S S 1907 *Melococcus ostrjanini* Charifkov Med žurn 4
309-312
- ARKWRIGHT, J A, BACOT, A, AND DUNCAN, F M 1919 The association of
Rickettsia with French fever Jour Hyg, Cambridge 18 76-94
- ARLOING, S 1889 Sur l'étude bactériologique des lésions de la péripneumonie
contagieuse du boeuf Compt rend Acad d sc Paris 109 428
- ARNDT, RUDOLF 1880 Beobachtungen an *Spirochaete dentucola*, des Spirochaete
des Zahnschleimes Arch Path Anat u Physiol (Virchow) 79
76-86
- ALEXENFELD, THEODOR 1897 Ueber die chronische Diplobacillen conjunctivitis
Centralbl f Bakteriologie, Jena 21 1
- BABES, V 1883 Le rein et le foie dans la fièvre jaune Compt rend Acad
d sc Paris 97 682-685
1888 Sur l'hémoglobininurie bactérienne du boeuf Compt rend Acad
d sc Paris 107 692
1889 Die Aetiologie des seuchenhaften Haemoglobinurie des Rindes Arch
Path Anat u Physiol (Virchow) 115 81
1890 in Cornil, A V, and Babes, V Les Bacteries
- BAKER, C F 1902 Revision of the *Elephantopaceae* Trans Acad Sci St Louis
12 53
- BALBIANI 1886 Evolution des Micro-organismes animaux et végétaux Jour
Microg 10 535-544
- BARENDRECHT, H P 1901 Die Agglutination von Hefe Centralbl f Bak-
teriologie, Jena Abt 2 7 623-627
- BARTOSZEWICZ, ST, AND SCHWARZWASSER, JOSEF 1908 Eine neue Form von
Diplococcus Tetradiplodoccus filiformans lodzensis (Vorl Mitt) Cen-
tralbl f Bakteriologie, Jena Abt 2 21 614-616
- BAUMGARTEN, PAUL V 1890 Lehrbuch der pathologischen Mykologie Leipzig
- DE BEAUREPAIRE ARAGAS, H, AND VIENNA, G 1913 Pesquisas sobre o granu-
loma venereo Mem do Inst Oswaldo Cruz 5. 211-238

- BÉCHAMP, A 1867 Sur le corpuscle vibrant de la pébrine, considéré comme organisme producteur d'alcool Compt rend Acad d sc Par 64 231-232
- 1868 Sur la maladie à mycrozymas de vers a soie Compt rend Acad d. sc Par 56 1160
- BEHRENS 1899 Review Koch's Jahresbericht 10 244-246
- BELJERINCK, M W 1888 Die Bakterien der Papilionaceenknollchen Bot Ztschr 46 758
- 1889 Le *Photobacterium luminosum* (Bactérie lumineuse de la mer du Nord). Arch néerl d sc exactes 23 428-444
- 1893 Ueber die Butylalkoholgarung und das Butylferment Verhandl K Akad Wetensch Amsterdam Sect II Deel I No 10 p 51
- 1894 Ueber die Butylalkoholgarung und das Butylferment Abs Centralbl f Bakteriologie Abt 1 15 171
- 1896 Sur la Fermentation et le Ferment butyliques Arch néerl d sc exactes Series I 29 1
- 1898 Sur les diverses espèces de bacteries acétifiantes Arch néerl d sc exactes Ser II 2 180-189
- 1898 Ueber die Arten der Essigbakterien Centralbl f Bakteriologie Abt 2 4 209-216
- 1900 Schwefelwasserstoffbildung in den Stadtgraben und Aufstellung der Gattung *Aërobacter* Centralbl f Bakteriologie, Jena Abt 2 6 193-206
- 1901 Over oligonitrophile bacterien (On oligonitrophilous bacteria) Amsterdam Versl Wis Nat Afd K Akad Wet, 9 633-642 (Dutch), Amsterdam, Proc Sci K Akad Wet 3 586-595 (English)
- 1901 Sur la formation de l'hydrogen sulfuré dans les canaux et le genre nouveau *Aërobacter* Arch Néerl Sci Soc Holl Ser 2 4 1-18
- 1901 Sur les ferments lactiques de l'industrie Arch néerl d sc exactes Ser 2 6 212-243
- 1901 Sur les ferments lactiques de l'industrie Haarlem Arch néerl Sci Soc Holl Ser 2 6 212-243
- 1901 Über oligonitrophile Mikroben Centralbl f Bakteriologie Abt 2 7 561-582
- UND VAN DELDEN, A 1902 Ueber die Assimilation des freien Stickstoffs durch Bakterien Centralbl f Bakteriologie, Jena Abt 2 9 3-43
- 1903 Sur des microbes oligonitrophiles Arch néerl d sc exactes Sec 2 8 217
- UND VAN DELDEN, A 1903 Sur l'assimilation de l'azote libre par les bactéries Haarlem, Arch Néerl Sci Soc Holl Ser 2 8 319-373
- UND VAN DELDEN, A 1904 Sur les bacteries actives dans le rouissage du lin Haarlem, Arch Néerl Sci Soc Holl Ser 2 9 418-441
- 1904 Ueber die Bakterien, welche sich im Dunkeln mit Kohlensäure als Kohlenstoffe ernähren können Centralbl f Bakteriologie, Jena Abt 2 11 598
- 1906 Une sarcine de fermentation anaerobie obligaloire Haarlem, Arch Neerl Sci Soc Holl Ser 2 11 199-205
- 1908 Fermentation lactique dans le lait Arch néerl d sc exactes Ser 2. 13 359

- 1908 Die Erscheinung der Flockenbildung der Agglutination bei Alkoholfäulen Centralbl f Bakteriöl, Jena Abt 2 20 641
- 1911 Ueber Pigmentbildung bei Essigbakterien Centralbl f Bakteriöl, Jena Abt 2 29 169-176
- 1914 Ueber Schroter und Cohn's Lackmusmicrococcus Folia microbiol, Delft 2 185-200
- 1914 Ueber das Nitratterment und ueber Physiologische Artbildung Folia microbiol, Delft 3 91-113
- BENECKE, WILHELM 1912 Bau und Leben der Bakterien
- BERESTNEW, N 1899 Zur Frage der Klassifikation und Systematischen Stellung der Strahlenpilze Centralbl f Bakteriöl, Jena Abt 1 26 390
- BERGAMASCHI, GIUSEPPE 1822 Osservazioni micologiche ed enumerazione storica di tutti i funghi di Pavese Milano
- BERGEY, DAVID H *et al* 1923 Manual of Determinative Bacteriology Baltimore
- BERGONZINI, CURZIO 1881 Sopra un nuovo Bacterio Colorato Annuario della Societa dei naturalisti in Modena Ser II 14 149-158
- BERKELEY, M J 1857 Introduction to Cryptogamic Botany
- 1874 Notes on North American Fungi Grevillea 3 64
- BETTENCOURT, ANNIBAL, AND KOPKE, AYRES, RENZENDE, GOMES DE, AND MENDES, CORREIA 1904 Ueber die Aetiologie der Schlafkrankheit Centralbl f Bakteriöl, Jena Abt I 35 55
- BIEDERT, PH 1885 Beitrag zur Frage nach der Constanz der Spaltpilze (*Kokkobacillus zymogenus* and *Bacterium termo*) Virchow's Arch f path Anat, Berl 100 439
- BILLET, A 1888 Sur le cycle évolutif et les variations morphologiques d'une nouvelle Bacteriacee marine Compt rend Acad d sc Par 106 293
- 1888 Sur le cycle évolutif d'une nouvelle Bacteriacee chromogène et marine, *Bacterium Balbiani* Compt rend Acad d sc Par 107 423
- 1890 Contribution a l'étude de la Morphologie et du Developpement des Bacteriacees Extrait du Bulletin scientifique de la France et de la Belgique Paris
- BILLROTH (CHRISTIAN ALBERT) THEODORE 1874 Untersuchungen ueber die Vegetationsformen von *Coccobacteria septica*—Berlin
- BIZIO, B 1819 Gazzetta privilegiata Venice Aug 24
- 1823 Polenta porporp Biblioteca italiana o sia Giornale di lettera, scienza ed arti (Anno VIII) 30 275
- 1827 Opuscoli chimico-fisici
- 1844 Sur un altération du pain causee par le developpement d'un cryptogame observation faite, il y a tres longtemps d'un phenomene qui rentre dans la class de ceux sur lesquels Moser a appelé l' Compt rend Acad d sc Par I Sem 18 951
- BLACK, G V 1892 Micro-organisms of the oral cavity Transactions of the Illinois Dental Society, 1886, cited by Miller Mikroorganismus der Mundhöhle p 78
- BLANCHARD, RAPHAEL 1896 Parasites végétaux a l'exclusion des Bactéries Bouchards' Traité de Pathologie Generale 2 811-926

- 1906 Spirilles, spirochètes et autres microorganismes à corps spiralé La Semaine Medicale 26 1
- BOLLINGER, OTTO 1870 Mycosis der Lunge beim Pferde Virchow's Arch f path Anat, Berl 49 583
- 1888 Ueber Botryomykose beim Pferd Kleinere Mittheilungen XIII Deut Ztschr Thermed 13 177
- BONGERT 1901 *Corynebacterium pseudotuberculosis murum* ein neuer pathogener Bacillus für Mäuse Ztschr f Hyg u Infektionskrankh, Leipz 37 449-475
- BONOME, A 1888 Pleuro-pericarditis und cerebro-spinal Meningitis Sero-fibrinosa durch einen dem *Diplococcus pneumoniae* sehr ähnlichen Mikroorganismus erzeugt Centralbl f Bakteriologie, Jena 2 321
- BORREL, DUJARDIN-BEAUMETZ, JEANTET AND JOUAN 1910 Le Microbe de la Peripneumonie Ann de l'Inst Pasteur, Par 24 179
- BORY DE ST VINCENT, J B 1824 Encyclopédie méthodique 1826 Microscopiques Dictionnaire classique d'histoire naturelle 10 533-546
- BORZI, ANTONINO 1878 Note alla morfologia e biologia delle Alghe Ficocromaceae Nuovo Gior Bot Ital 10 236-288
- BOSANQUET, WM C 1911 Spirochaetes
- BREED, R S, CONN, H J, AND BAKER, J C 1918 Comments on the evolution and classification of bacteria Jour Bact, Balt 3 445-459
- AND CONN, H J 1919 The nomenclature of the Actinomycetaceae Jour Bact, Balt 4 585-602
- 1920 The nomenclature of the Actinomycetaceae (Addenda) Jour Bact, Balt 5 489-490
- BRUMPT, E 1910 Precis de Parasitologie Paris
- BRUSSOFF, A 1916 *Ferribacterium duplex*, eine stabchenförmige Eisenbakterie Centralbl f Bakteriologie, Jena Abt 2 45 546-554
- BUCHANAN, R E 1915 Nomenclature of the Coccaceae J Infect Dis, Chicago 17 528-541
- 1916 Studies in the nomenclature and classification of the bacteria I The problem of bacterial nomenclature Jour Bact 1 591-596
- 1917 Studies in the nomenclature and classification of the bacteria II The primary subdivisions of the Schizomycetes Jour Bact 2 155-162
- 1917 Studies in the nomenclature and classification of the bacteria III The families of the Eubacteriales Jour Bact 2 347-350
- 1917 Studies in the nomenclature and classification of the bacteria IV Subgroups and genera of the Coccaceae Jour Bact 2 603-617
- 1918 Studies in the nomenclature and classification of the bacteria V Subgroups and genera of the Bacteriaceae Jour Bact 3 27-61
- 1918 Studies in the nomenclature and classification of the bacteria VI Subdivisions and genera of the Spirillaceae and Nitrobacteriaceae Jour Bact 3 175-181
- 1918 Studies in the classification and nomenclature of the bacteria VII The subgroups and genera of the Chlamydo bacteriales Jour Bact 3 301-306

- 1918 Studies in the classification and nomenclature of the bacteria VIII
The subgroups and genera of the *Actinomycetales* Jour Bact 3:
403-406
- 1918 Studies in the classification and nomenclature of the bacteria IX
The subgroups and genera of the *Thiobacterales* Jour Bact 3
461-474
- 1918 Studies in the classification and nomenclature of the bacteria X
Subgroups and genera of the *Mycobacteriales* and *Spirochaetales* Jour
Bact 3 541-545
- BUDER, JOHANNES 1913 *Chloronium mirabile* Ber d deutsch Bot Ge-
sellsch, Berl 31 80-97
- BUMM, E 1885 Der Mikroorganismus der Gonorrhoeischen Schleimhaut Er-
krankungen, "Gonococcus-Neisser" Wiesbaden
- BURDON-SANDERSON, JOHN S 1871 Origin and Distribution of the Microzymes
in Water and in the Tissues and Liquids of the Organism Quart
Jour Micr Sc, Lond 11 323-352
- BURRILL, T J 1883 New species of *Micrococcus* (Bacteria) Amer Nat 17
319-320
- CASPARY, ROBT 1874 *Merismopedium Restenbachii*, n sp Schr Physikal
Okonom Gesell Konigsb 15 104
- CASTELLANI, ALDO, AND CHALMERS, ALBERT J 1919 Manual of Tropical
Medicine, 3rd ed
- 1920 Sur la classification de certain groupes de bacilles aerobies de l'intestin
humain Ann de l'Inst Pasteur, Par 34 600-621
- CHALMERS, ALBERT J, AND CHRISTOPHERSON, J B 1916 A Sudanese Actino-
mycosis Ann Trop Med and Parasitol, Liverp 10 223-283
- CHATTON, EDOUARD, AND PÉRARD, CHARLES 1913 Schizophytes du caecum du
cobaye I *Oscillospira guillermondi*, n g, n sp, II *Metabacterium*
polyspora n g, n sp Compt rend Soc de biol, Par 74 1159 and 1223
- CHAUTEMASSE, A, MATRUCHOT, L, AND GRUNBERG, A 1917 Une maladie
nouvelle, simulant le rhumatisme articulaire aigu, avec ses complica-
tions viscérales et causée par un microbe nouveau (*Mycobacillus syn-
novialis*) Bull Acad de méd, Par 3rd Ser 77 708-712
- 1917 Un microbe nouveau *Mycobacillus synovialis*, causant chez l'homme un
maladie évoluant comme le rhumatisme articulaire Compt rend Acad
d sc, Par 164 562-565
- CHESTER, F D 1897 A preliminary arrangement of the species of the genus
Bacterium 9th Ann Rept Del Coll Agr Exp Sta p 53-146
- 1897 Classification of the *Schizomycetes* 9th Ann Rept Del Coll Agr
Exp Sta p 62-63
- 1898 A preliminary arrangement of the species of the genus *Bacillus* 10th
Ann Rept Del Coll Agr Exp Sta p 100-137
- 1899 Description of certain species of bacteria isolated from cultivated soil
Ann Rept Del Coll Agr Exp Sta p 52-75
- 1899 Studies in systematic bacteriology Ann Rept Del Coll Agr Exp
Sta 11: 36

- 1901 Manual of Determinative Bacteriology
- 1903 Observations on an important group of soil bacteria 15th Ann Rept
Del Coll Agr Exp Sta p 42-96
- CHUDIakov, N 1897 Zur Lehre von der Anaerobiose Theil I (Russian)
Koch's Jahresbericht 8 44-46
- CIEŃKOWSKI 1878 Untersuchung über die Gallertbildungen des Zuckerruben-
saftes (Résumé allemand du mémoire russe Charkow) Arbeit d
Naturforsch Gesellsch a d Univ a Charkoff p 12
- CLEMENTS, FREDERICK E 1894 Family—*Bacteriaceae* in Part I Flora of
Nebraska p 25-28
- 1902 Greek and Latin in biological nomenclature University Studies
Pub by U of Nebraska 3 1-84
- 1909 The Genera of Fungi Minneapolis
- COATS, G, AND FORBES, J G 1910-1911 On the relation of the *Meningococcus*
intracellularis to Pseudoglioma Proc Roy Soc Med, Lond 4 Pathol
Section p 242-250
- COHN, FERDINAND 1854 Untersuchungen über die Entwicklungsgeschichte
der mikroskopischen Algen und Pilze Der Akademie übergeben den
1 Mai, 1853 Nov Act Acad Caes Leop-Carol Nat Cur 24 103-256
- 1872 Untersuch über Bakterien Beitr Biol Pflanz 1 127-224
- 1872 (a) Organismen in der Pockenlymphe Virchow's Arch f path Anat,
Berl 55 229-238
- 1870 Ueber den Brunnenfaden (*Crenothrix polyspora*) mit Bemerkungen
über die mikroskopische Analyse des Brunnenswassers Beitr Biol
Pflanz (Heft 1) 1 108-131
- 1875 Untersuchungen über Bakterien II (Heft 3) Beitr Biol Pflanz
1 141-208
- 1876 Untersuchungen über Bakterien IV Beiträge zur Biologie der
Bacillen (Heft 2) 2 249-277
- COLLIN, BERNARD 1913 Sur un ensemble de Protistes parasites des Batraciens
Arch zool exper 51 59-76
- CONN, H W 1909 Agricultural Bacteriology Ed 2
- CORDA, A C J 1839 Prachtflora europäischer Schimmelbildungen Leipzig
u Dresden p 27
- CORNIL AND BABES 1890 Les Bacteries
- CUNNINGHAM, ALLAN 1839 Flora Insularum Novae Zelandiae Precursor
Ann Nit Hist 2 47
- DALLINGER, W H 1878 On the measurement of the diameter of the flagella
of *Bacterium termo* a contribution to the question of the "Ultimate
Limit of Vision" with our present lenses Jour Roy Micr Soc, Long
1 159-175
- DANGEARD, A 1909 Memoire sur les Chytridiées Le Botaniste 1 55
- 1891 Contribution à l'étude des Bacteriacees vertes (*Eubacillus* nov gen)
Le Botaniste 2 151
- DA ROCHA, LIMA 1916 Zur Aetiologie des Fleckfiebers Berl klin Wehnschr
p 567-569

- DAVAINE, C 1863 Recherches sur les infusoires du sang dans la maladie connue sous le nom de sang de rate *Compt rend Acad d sc*, Par 57 220-223
- 1863 Nouvelle recherches sur la maladie du sang de rate considérée au point de vue de sa nature *Memoires lus a la Société de Biologie* Ser 3 5 193-202
- 1864 Nouvelles recherches sur la nature de la maladie charbonneuse connue sous le nom de sang de rate *Compt rend Acad d sc*, Par 59 393-396
- 1865 Sur la présence constante des Bactériidies dans les animaux affectés de la maladie charbonneuse *Compt Rend Acad d sc*, Par 61 334-335
- 1868 Dictionnaire Encyclopedique des Sciences Medicales Ser I. 8 21
- DE BARY, A 1884 *Vergl Morph u Biol d Pilze, Mycetozaa und Bacterien* Leipzig
- 1886 *Vorlesungen uber Bakterien*
- D'HERELLE, F 1914 Le Coccobacille des Sauterelles *Ann de l'Inst Pasteur*, Par 280 28-387
- DE KRUYFF, E 1906-8 Les Bacteries hydrolysant et oxydant les graisses *Bull* 1-15 Dept van Landbouw Dutch East Indies
- DE LANESSAU, J T 1880 *Revue Internationale des Sciences Biologiques* 5 245
- DESMAZIERES, J B 1826 Recherches microscopiques et physiologiques sur le genre *Mycoderma* *Ann Sci Nat Bot Ser 1* 10 42-67
- 1851 Sur les plantes cryptogames recement decouvertes en France *Ann Sci Nat Bot Ser 3* 16 319
- DE TONI, J B, AND TREVISAN, V 1889 *Schizomycetaceae* Naeg, in Saccardo, P A *Sylloge Fungorum* 8 923-1087
- DOBELL, C CLIFFORD 1911 *Paraspirillum vej dovskii* n g n sp, a new bacterial form *Arch f Protistenk*, Jena 24 97
- 1911 On *Cristispira veneris* n sp and the affinities and classification of spirochaets *Quart Jour Micr Sc*, Lond 56 507-540
- DOCHEZ, A R, AND AVERY, O T 1915 Varieties of *Pneumococcus* and their relation to lobar pneumonia *Jour Exper Med*, Balt 21 114
- DOFLEIN, F 1911 *Lehrbuch der Protozoenkunde*, Jena
- DOPTER, CH 1909 Étude de quelques germes isoles du Rhino-pharynx, voisin du Méningo-coque (para-meningococques) *Compt Rend Soc de biol*, Par 67 74-76
- DUBOSCO, O, AND LEBAILLY, CH 1912 *Spirella canis* n g, n sp, spirelle de l'estomac du chien *Compt rend Acad d Sc*, Par 154 835-837
- DUCLAUX, E 1882 Memoire sur le lait *Ann Inst Nat Agron* 4 23-138 (for years 1879-80) Chapter IV Ferments du lait pp 74-117
- 1895 Sur la Nutrition intra-cellulaire *Ann de l'Inst Pasteur*, Par 9 811-839
- DUJARDIN, FELIX 1841 *Hist nat des Zoophytes infusoires* Paris Mem sur une classification des Intusoires en rapport avec leur organisation *Compt rend Acad d sc* 11 281-286 1840 *Microsc Journ* 1 53-56 1841

- 1841 Histoire naturelle des Zoophytes Infusoires Paris
- DUNBAR, W 1893 Untersuchungen über choleraähnliche Wasserbakterien Deutsche med Wchnschr, Berl and Leipz 19 799
- DUNCKER, H C J 1887 Ueber *Actinomyces musculorum suis* Arch f wissensch u prakt Tierh, Berl 13 224
- DYER, W. T THISELTON 1875 Sexual reproduction of Thallophytes Quart Jour Micr Sc, Lond N S 15 295-326
- EDINGTON, ALEXANDER 1887 A method of prophylaxis and an investigation into the nature of the contagium of scarlet fever Brit Med Jour 1. 1265
- EDKINS, J S 1923 *Spirella regaudi* in the cat Parasitology 15 296-307
- EHRENBERG, C G 1828 Symbolae Physicae seu Icones et Descriptiones animalium evertibratorum sepositis insectis quae ex itinere per Africam Borealem et Asiam occidentalem IV Evertabrata Berlin
- 1830 Bertrage zur Kenntniss der Organisation der Infusorien und ihrer geographischen Verbreitung Abhandlungen d Berl Akad p 1-80
- 1832 Die geographische Verbreitung der Infusionsthierchen in Nord-Afrika und West Asien, etc Abhandl d K Akad d Wissensch zu Berlin Physik Klasse
- 1833 Dritter Bertrag zur Erkenntniss grosser Organisation in der Richtung des kleinsten Abhandl Berl Akad p 145-336
- 1836 Vorlauffige Mittheilungen über das wirkliche Vorkommen fossiler Infusionen und ihre grosse Verbreitung Ann Phys Ser 2 8 217
- 1838 Die Infusionsthierchen als vollkommende Organismen Leipzig
- 1840 Charakteristik vom 274 neuen Arten von Infusorien Berichte über die Verhandlungen der Königl Preuss Akademie der Wissenschaften zu Berlin 5 197-219
- 1848 Fortsetzung der Beobachtung des sogenannten Blutes im Brode als *Monas prodigiosa* Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königl Preuss Akademie der Wissenschaften zu Berlin 354-362
- EIDAM, E 1876 Ueber die Entwicklung des *Sphaerotilus natans* Jahresber Schles Gesell Vaterland Cult 54 133
- EISENBERG, JAMES 1886 Bakteriologische Diagnostik Hamburg and Liepzig (Voss)
- 1891 Bakteriologische Diagnostik Ed 3
- ELLIS, D 1897 On the discovery of a new genus of thread-bacteria (*Spirophyllum ferrugineum*) Proc Roy Soc, Edinb 27 part I (No 6) p 21
- 1907 A contribution to our knowledge of the thread bacteria Centralbl. f Bakteriol, Jena Abt 2 19 516
- 1909 Outlines of Bacteriology
- 1910 A contribution to our knowledge of the thread-bacteria (II). Centralbl f Bakteriol, Jena Abt 2 26
- ELSER, W J, AND HUNTOON, F M 1909 Studies on meningococcus Jour. Med Research, Bost 20 371-541

- ENGELMANN, T W 1883 *Bacterium photometricum* Ein Beitrag zur vergleichenden Physiologie des Lichts und Farbensinne Pflugers Arch f. Physiol 30 95-124
- ENGLER, A 1882 Ueber die im Kieler Hafen in dem sogenannten "toten Grund" vorkommenden Pilzformen Verhandl Bot Ver Brandenb 24: 17-20
- 1882 Ueber die Pilz-Vegetation des weissen oder todten Grundes in der Kieler Bucht Vierter Bericht der Commission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel fur 1877 bis 1881 Abt 1 187-193
- ENGLER, G 1912 Syllabus der Pflanzenfamilien Ed 7
- ENLWS, ELLA M A 1920 The generic names of bacteria Hygienic Laboratory—Bull No 121 Treasury Dept U S P H Service
- ESCHERICH, TH 1885 Die Darmbakterien des Neugeborenen und Sauglings. Fortschr d Med No 15 3 520
- 1886 Die Darmbakterien des Sauglings
- 1886 Beitrage zur Kenntniss der Darmbakterien Munchen med Wehnschr 33 759-763
- EVANS, ALICE C The nomenclature of the melitensis—abortus group of bacterial organisms Pub Health Rep U S Mar Hosp Serv, Wash 38-1943-1948
- FALKENHEIM, H 1885 Ueber Sarcine Arch f exper Path u Pharmacol, Leipz 19 339
- FAMINTZIN, A 1892 Arbeiten aus dem botanischen Laboratorium der Kaiserlichen Akademie der Wissenschaften zu St Petersburg No 2 Eine neue Bacterienform *Nevskia ramosa* Bul Acad Imp Sci St—Petersb 34 (N S 2) 481-486
- FAULL, J H 1915 A new myxobacterium (Proc Roy Soc Canada) Sci N S 43 469
- FEHLEISEN 1883 Die Aetiologie des Erysipels Berlin
- FICK, ADOLF E 1887 Ueber Mikroorganismen in Conjunctivalsack Wiesbaden
- FISCHER, ALFRED 1895 Untersuchungenuber Bakterium Jahrb f wissenschaft Bot, Berl 27 1-163
- 1897 Lectures on Bacteria Eng translation of Vorlesungen uber Bakterien
- 1903 Vorlesungen uber Bakterien 2nd ed
- FISCHER, BERNHARD 1894 Die Bakterien des Meeres Ergebnisse der Plankton—Expedition der Humboldt—Stiftung p 1-82
- FITZ, ALK 1878 Ueber Schizomyceten—Gahrungen III Ber Deut Chem Gesell 11 42-55
- FLUGGE, C 1886 Die Mikroorganismen, Leipzig
- 1908 Grundriss der Hygiene Ed 6
- FOÀ AND BORDONO-UFFREDUZZI 1888 Ztschr f Hyg u Infektionskrankh, Leipz 4 67
- FORTI, C 1896 Notizie complementari su alcuni studi di zimotecnica enologica eseguiti a tutto il 1896 Ann della soc chim di Milano Vol 3, fasci 1
- FORTINEAU, L 1905 *L'Erythrobacillus pyosepticus* Compt rend Acad d sc. Par et de la Soc de Biologie Paris 58 104

- FRANK, B 1889 Ueber die Pilzsymbiose der Leguminosen Ber d deutsch Bot Gesellsch, Berl 7 338
- FRENZEL, JOHANNES 1884 Ueber die Mikrozymas in der Leber und im Pankreas Biol Centralbl, Leipz 3 49-52
- 1897 Neue oder wenig bekannte Susswasserprotisten Biol Centralbl 17 801-808
- FREUDENREICH, EDUARD 1894 Les Microbes et leur rôle dans la Laiterie
- FROST, W D 1911 in Marshall's Microbiology
- FUCHS, C H 1841 Beitrage zur nahern Kenntniss der gesunden und fehlerhaften Milch der Hausthiere Magazin fur die gesammte Thuerheilkunde 7 133-198
- FUEHRMANN, F 1905 Morphologisch-biologische Untersuchungen uber ein neues Essigsauresbildendes Bakterium Beiheft zum bot Centralbl 19 1-31
- GAMELÉIA, N 1888 *Vibrio Metschnikovi* (n sp) et ses rapports avec le microbe du cholera asiatique Ann de l'Inst Pasteur, Par 2 482
- 1888 Zur Aetologie der Huhnercholera Centralbl f Bakteriologie, Jena 4 167
- GASPERINI, G 1889 Recherches morphologiques et biologiques sur un Microorganisme de l'Atmosphere, le *Streptothrix Foersteri* Cohn Ann Microg 10 449
- 1895 Discussioni Atti dell' XI congresso med internaz Roma 6 (igiene) 80-82
- GEDOELST, L 1902 Les Champignons Parasites de l'Homme et des Animaux Domestiques Paris
- GLASER, R W, AND CHAPMAN, J W 1912 Studies on the wilt disease or "flacherie" of the gypsy moth Science, N Y and Lancaster, n s 36 219-224
- GLEITSMANN 1913 Beitrag zur Entwicklungsgeschichte der Spirochäten (Borrelhen) Centralbl f Bakteriologie, Jena Abt I 68 31
- GONDER, RICHARD 1914 Spirohemaceae (Spirocheten) Prowazek's Handbuch der pathogenen Protozoen Lief 6 p 671
- GONNERMANN, M 1907 Ueber gallertbildende Bakterien Zeitschrift fur Zuckerindustrie und Landwirtschaft 36 877-888
- GOODSIR, JOHN 1842 History of a case in which a fluid periodically ejected from the stomach contained vegetable organisms of an undescribed form Edinburgh Med and Surg Jour 57 432
- GRIFFITH, J W 1853 On *Gallionella ferruginea* (Ehrenb) Ann and Mag Nat Hist Ser 2 12 438
- GROENEWEGE, J 1912 De rotting der tomatenvruchten, veroorzakt door *Phytobacter lycopersicum*, n sp meded van de Rijks Hoog Land- Tuin- en Boschbouwschool Deel v 5 Afl 5 217-239
- GROSS, J 1910 *Cristispira nov gen* Ein Beitrag zur Spirochätenfrage Mittheil zool Stat Neapel 20 41-93
- 1911 Ueber freilebende Spirohemaceen Mittheil Zool Stat Neapel 188-204
- 1912 Ueber Systematik, Struktur und Fortpflanzung der Spirohemaceae Centralbl f Bakteriologie, Jena Abt I 65 83

- GROVE, W B 1884 A Synopsis of the Bacteria and Yeast Fungi
- GRUBER, M 1891 *Micromyces Hoffmanni* Centralbl f Bakteriöl 10 648
- GRUBER, THEODOR 1895 Die Arten der Gattung *Sarcina* Arbeiten Bacteriologisches Institut der Technischen Hochschule zu Karlsruhe 1.239-290.
- GUERBET, M 1906 Notes sur la fermentation du yoghurt Compt rend Soc de biol, Par 60 495
- GUIGNARD, LEON 1890 Sur une nouvelle Bacteriacee marin, le *Streblotricha Bornetii* Compt rend Soc de biol, Par 42 124
- GUILLEBEAU, A 1890 Studien über Milchfehler und Euterentzündungen bei Rindern und Ziegen Landwirtsch Jahrb d Schweiz, Berne 4: 27-44
- 1891 Beiträge zur Lehre von den Ursachen der fadenziehenden Milch Landwirtsch Jahrb d Schweiz, Berne
- 1892 Studien u Milchfehlen und Euterentzündungen bei Rindern und Ziegeln I Ueber Ursachen der Euterentzündungen Landwirtsch Jahrb d Schweiz, Berne 4 27-44
- GUTTMANN, PAUL 1880 Zur Histologie des Blutes bei Febris recurrens Arch f path Anat u Phys Virchow 80 1-9
- HAASS, EVERHARD 1906 Beiträge zur Kenntniss der Aktinomyceten Centralbl f Bakteriöl, Jena Abt I 40 180-186
- HAASE, W 1887 Gesammelte Beobachtungen über die Influenza der Pferde Archiv für Tierheilkunde 13 347
- HAJEK, M 1888 Die Bakterien bei der akuten und chronischen Coryza, sowie bei der Ozaena, und deren Beziehungen zu den genannten Krankheiten Berliner klin Wochenschr 25 659
- HALLER, ERNST 1866 Bericht über mykologische Untersuchungen Bot Zeitung 24 383
- 1866 Die Pflanzlichen Parasiten des menschlichen Körpers für Aerzte, Botaniker und Studierende zugleich als Einleitung in das Studium der niederen Organismen Leipzig
- 1868 Ueber einen bei der Rotzkrankheit der Pferde auftretenden Parasiten, verglichen mit dem der Syphilis Bayerischen ärztlichen Intelligenzblatt p 327
- HANSEN, E 1879 De af mig i øl og ølurt jagttagne Organismer Meddel Carlsberg Lab Labor 2 222
- 1894 Gesammelte Theoretische Abhandlungen über Gärungsorganismen p 465
- HANSRIG, ANTON 1888 Neue Beiträge zur Kenntniss der halophilen der thermophilen und der Berg Algen-Flora, sowie der chemophilen Spaltpilz flora Böhmen 38 41, 87, 114, 149
- 1888 Beiträge zur Kenntniss der Kellerbakterien nebst Bemerkungen zur Systematik der Spaltpilze (Bacteria) Osterr Bot Ztschr 38 227-230, 263-267
- 1890 Ueber neue Süsswasser- und Meeres-Algen und Bacterien II Aero-phytische und im Meere lebende Bacterien Sitzungs Berichte der K böhm Gesellsch der Wissensch Math -Naturwiss Classe 1 1-34

- 1890 Ueber neue Susswasser Bakterien mit Bemerkungen zur Systematik dieser Phycophyten und uber den Einfluss des Lichtes auf die Ortsbewegungen des *Bacillus Pfefferi* Sitz-ber K bohm Gesellsch d Wissenschaften p 20-31
- 1891 Ueber die Bacteriaceen Gattung *Phragmidiothrix* Engler und einige *Leptothrix* arten Bot Ztg 49 313
- HARTLEB, R 1900 Die Morphologie und systematische Stellung der sogenannten Knollchenbakterien Chem Ztg 24 887
- HARTLEB, R AND STUTZER, A 1901 Bemerkungen zu der Mittheilung von Dr W Rullmann Centralbl f Bakteriöl, Jena Abt 2 3 621
- HARZ, C O 1877-78 *Actinomyces bovis*, ein neuer Schimmel in dem Gewebe des Rindes Jahresber d Muenchener Central-Thuerarzneischule p 125-140
- HAUSER, G 1885 Ueber die Entwicklungsgeschichte und pathogenen Eigenschaften einer faulniserregenden Bakterienart Sitz-ber d phys-mediz Sozietat zu Erlangen 156-171 After C Fischer Biol Centralbl, Leipz 5 36-38
- HAZEN, TRACY E 1896-99 Life history of *Sphaerella lacustris* Mem Torrey Bot Club 6 211-246
- HEIM, LUDWIG 1911 Lehrbuch der Bakteriologie
- HELLER, HILDA HEMPL 1921 Suggestions concerning a rational basis for the classification of the anaerobic bacteria Jour Bact, Balt 6 521-553
- 1922 Certain genera of the *Clostridiaceae* Jour Bact, Balt 7 1-36
- HENFREY, ARTHUR 1856 Notes on some fresh water confervoid algae new to Britain Transactions of the Microscopical Society of London New Series 4 49-54
- HENNEBERG, W 1909 Garungsbakteriologisches Praktikum, Betriebsuntersuchungen und Pilzkunde p 480
- HERTWIG 1886 Ueber den *Actinomyces musculorum* der Schweine Arch f wissensch u prakt Tierh, Berl 12 365-372
- HEWLETT, R I 1898 A Manual of Bacteriology p 18
- HINZE, G 1903 *Thiophysa volutans*, ein neues Schwefelbakterium Ber d deutsch Bot Gesellsch, Berl 21 309
- HOCHSTETTER, CH F 1842 Nova genera plantarum Africae australis tum tropicae borealis proponit et describit Flora 25 226
- HOELLING, A 1910 Die Kernverhaltnisse von *Fusiformis termatidis* Arch f Protistenk, Jena 19 240
- HOHL, J 1902 Ein neuer, aus Strohisolierter, "das Fadenziehen" der Milch verursachender Coccus (*Carphococcus pituitoparus*) Centralbl f Bakteriöl, Jena Abt 2 9 338-344 Also Molchztg 12 446-448
- HOYER, DIRK PIETER 1898 Bijdrage tot de kennis van de Azijnbacterien Proefschrift ter Verkriging van den Graad van Doctor in de Scheikunde van de Rijks-Universiteit te Leiden 115 pp Delft (Waltma jr) Abs Cent 2nd Abt 4 867-875
- HUEPPE, F 1886 Die Formen der Bakterien und ihre Beziehungen zu den Gattungen und Arten Wiesbaden
- HUSS, HARALD 1908 Eine fettspaltende Bakterie (*Bactradium hypolyticum* n sp) Centralbl f Bakteriöl, Jena Abt 2 20 474-484

- ITZIGSOHN, HERMANN 1858 Zur Naturgeschichte der *Sarcina ventriculi* Good-
sir. Arch f path Anat u Phys (Virchow) 13 541-549.
- JACQUÉ, LÉON, AND MASAY, FERNAND 1912 Le *Streptobacterium foetidum*,
agent pathogène nouveau de l'homme. Centralbl f Bakteriöl, Jena
Abt 1 Orig 62: 180-181
- JEGUNOW, M 1896 Bakterien-Gesellschaften Centralbl f Bakteriöl, Jena
Abt 2 2 11-21
- JENNINGS, A VAUGHN 1899 On a new genus of bacteria (*Astrobacter*) Proc
Roy Irish Acad Ser 3 5 315
- JOHNE, ALBERT 1881 Die Actinomykose oder Strahlenpilzerkrankung, eine
neue Infektionskrankheit Deut Ztschr Thiermed 7 141-192
- JORDAN, E O 1908 General Bacteriology
- KALCHBRENNER, REV CARL AND COOKE, M C 1880 Australian Fungi
Grevillea 9 17-34
- KARWACKI, LEON 1909 Casuistique bacteriologique *Coccobacillus fusiformis*
n sp Agent pathogene de la septicemie Gaz lek, Warszawa 29
679-681
- KENDALL, ARTHUR I 1902 A proposed classification and method of graphical
tabulation of the characters of bacteria Proc Amer Pub Health
Ann 28 484
- KERN, EDUARD 1882 *Dispora caucasica* nov g et nov sp, eine neue Bak-
terienform Biol Zent 2 135
1882 Ueber ein neues Milchferment aus dem Kaukasus Bul Soc Imp Nat.
Moscou 56 (part 2) 141-173
- KIRCHNER, O 1878 Algen Kryptogamen-flora von Schlesien
1895 Die Wurzelknöllchen der Sojabohne Beitr Biol Pflanz 7 213
- KITT, TH 1888 Der *Micrococcus ascoformans* und das Myofibrom des Pferdes
Centralbl f Bakteriöl, Jena 3 247
- KIEBS, EDWIN 1871 Die Ursachen der infectiosen Wundkrankheiten Cor -Bl
f schweiz Aerzte
1875 Beiträge zur Kenntniss der pathogenen Schistomyceten Arch f
exper Path u Pharmakol, Leipz 1873, I 31, 443, 1875, III 305, IV
107, 207, 409, 1876, V 350
1879 Das Contagium der Syphilis Arch f exper Path u Pharmakol,
Leipz 10 161-218
- KLECKI, V VON 1894 Ueber einige aus ranziger Butter kultivierte Mikro-
organismen Centralbl f Bakteriöl, Jena Abt I 15 354-362
- KLEIN, E 1885 Microorganisms and Disease
- KLIGLER, I J 1913 A systematic study of the *Coccaceae* in the collection of the
Museum of Natural History Jour Infect Dis, Chicago 12 432
1915 A Biochemical Study and Differentiation of Oral Bacteria, with Special
Reference to Dental Caries Diss Ph D Columbia University
- KLINGER, R 1912 Ueber einen neuen pathogenen Anaëroben aus mensch-
lichem Eiter (*Coccobacterium mucosum anaërobicum* n sp) Centralbl
f Bakteriöl, Jena Abt I Originale 62 186-191

- KLOCKER ALB 1903 Fermentation Organisms English Translation by Allan and Miller, p 332
- KOCH, ALFRED AND OELSNER, ALICE 1923 Ueber nucleoproteidespaltende Bakterien und ihre Bedeutung für die Erschliessung des Phosphorkapitals im Ackerboden Biochem Ztschr 134 76-96
- KOLLE, W, AND HETSCH, H 1911 Die experimentelle Bakteriologie und die Infektionskrankheiten 3rd ed
- KOPPANYI, EMERICH 1907 Ueber eine mit fibrinösen Pleuritis einhergehende Pyämie der Kaninchen Ztschr f Tiermed, Jena 11 429-454
- KOSSOWICZ, ALEX 1911 Einführung in die Mykologie der Nahrungsmittelgewerbe
1912 Einführung in die Agrikulturmykologie
- KROKIEWICZ, A 1904 Zur Pathogenese des Diplopneumococcus Frankel-Wechselbaum, ein Fall von Pneumococci sepsis metastatica Wien klin Wchnschr 17 563-565
- KRZYSZTAŁOWICZ AND SIEDLICKI 1908 Etude expérimentale de la Syphilis, morphologie de *Spirochaeta pallida* Bul Acad Sci Cracovie p 173-234
- KUTTNER, H 1895 Ueber einen neuen, beim Menschen gefundenen Eitererreger Ztschr f Hyg u Infektionskrankh, Leipz 19 263-290
- KUTZING, F T 1833 *Sphaerotilus natans*, eine neue Süßwasseralge Linnaea 8 385
1837 Mikroskopische Untersuchungen über die Hefe und Essigmutter Jour Prakt Chem 11 386
1843 Phycologia generalis
1849 Species Algarum
1904 Mikroskopische Untersuchungen über die Hefe und Essigmutter, nebst mehreren anderen dazu gehörigen vegetabilischen Gebilden Auszüge vorgetragen in der Versammlung des naturhistorischen Vereins der Harzes in Alexisbad am 26 Juli 1837 Reprinted in Delbrück, M and Schrohe, A Hefe, Gärung und Faulnis p 25-38
- KUNTH, KARL SIGISMUND 1850 Enumeratio plantarum, etc 5 35
- KURTH, H 1883 Ueber *Bacterium Zopfii*, eine neue Bakterienart Ber d deutsch Bot Gesellsch, Berl 1 97-100
1883 Ueber *Bacterium Zopfii*, eine neue Bakterienart Bot Ztg 41 369-386, 393-405, 409-420, 425-435
- LACHNER-SANDOVAL, V 1898 Ueber Strahlenpilze Inaugural Diss Strassburg
- LAGERHEIM, G 1901 Mykologische Studien III Beiträge zur Kenntniss der parasitischen Bakterien und der bakterioiden Pilze Bihang till K svenska Vetenskaps-Akademien Handlingar Band 26 No 4 1900 Bot Centralbl, Cassel 85 280-283
- LANKESTER, E RAY 1873 On a peach-coloured bacterium—*Bacterium rubescens* Quart Jour Micr Sc, Lond 13 408-410
1876 Further observations on a peach- or red-coloured bacterium—*Bacterium rubescens* Quart Jour Micr Sc, Lond N S 16 27-40
- LANZI, M. 1874 Sulla origine e natura dei batteri Arch di Med chir ed ig Roma 12 65, 129
1876 I Batteri Parassitici funghi Nuovo Gior Bot Ital 8 256-261

- LAUTERBORN, ROBERT 1898 Ueber *Modderula hartwigi* Frenzel Biol Centralbl, Cassel 18 95
 1906 Zur Kenntniss der sapropelischen Flora Allg bot Ztschr Karlsruhe 12 196-197
 1907 Eine neue Gattung der Schwefelbakterien (*Thioploca Schmidler* nov gen nov spec) Ber d deutsch Bot Gesellsch, Berl 25 238-242
 1913 Zur Kenntniss einiger sapropelischer Schizomyceten Allgemeine Bot Zeitschrift 19 99-100
- LEBERT, HERMANN 1856-57 Ueber die Krankheit des Insekts der Seide Verein z Beford d Seidenbaues d Prov Brandenb
- LEHMANN, K B, AND NEUMANN, R 1896 Atlas und Grundriss der Bakteriologie Munchen
 1901 Bacteriology English Translation of 2nd German edition by G H. Weaver 2 127
- LEIDY, JOSEPH 1849 Entophyta Proc Acad Nat Sc, Phila 4. 225
 1849 Entophyta Proc Acad Nat Sc, Phila 4 249
 1851 Descriptions of new entophyta growing within animals Proc Acad Nat Sc, Phila 5 35
 1852 A flora and fauna within living animals Smithsonian Contributions to knowledge 5 1-67
- LEUBE, W 1885 Ueber die ammoniakalische Harngahrung Archiv f path Anat (Virchow) 50 540-570
- LÉVEILLÉ, J H 1851 Organisation et Disposition methodique des especes qui composent le genre *Erysiphé* Annales science naturelles Botanique Series 3 16 109-179 and 381
- LEVY, E 1899 Ueber die Actinomyces gruppe (Aktinomyceten) Centralbl f Bakteriöl, Jena Abt I 26 1
- LEWIS, TIMOTHY RICHARDS 1879 The Microphytes which have been found in the blood and their relation to disease Quart J Micr Sc, Lond N S 19 356-404
- LIESKE, RUDOLF 1921 Morphologie und Biologie der Strahlenpilze
- LIGNIÈRES, M 1900 Contribution a l'étude et a la classification des Septicemies hemorrhagiques Rec de med vet, Par Ser 8 7 331-332
 1901 Contribution a l'étude et la classification des septicemies hemorrhagiques, les "Pasteurelloses" Ann de l'Inst Pasteur, Par 15. 734-736
- LIGNIÈRES, JOSE AND SPITZ, JORGE 1902 Actinobacilosis Boletin de Agricultura y Ganaderia Buenos Ayres 11 169-230
 1904 Contribution à l'étude, à la classification et à la nomenclature des affections connues sous le nom d'actinomycose Centralbl f Bakteriöl, Jena Abt I Originale 35 452-461
- LINDAU, GUSTAV 1898 Bibliography Schizomyceten 1894-1898 Sammelwerke, Lehrbucher, Monographien etc Gonococcus Neisseri p 100-103 Just's Bot Jahresber Abt 1 26 1-139
- LINDNER, P 1887 Die Sarcina-Organismen der Gahrungsgewerbe Diss Berlin 1888 Centralbl f Bakteriöl, Jena 2 340
 1909 Mikroskopische Betriebskontrolle in den Garungsgewerben Ed 5

- LINGELSHIM, W v 1912 Streptokokken Kolle and Wasserman's Handbuch der pathogenen Mikroorganismen Ed 2 p 494
- LINK, H 1795 Dissertationes botanicae, Sverini F 65
- LISTER, JOSEPH 1873 A further contribution to the natural history of bacteria and the germ theory of fermentation changes Quart Jour Micr. Sc, Long xiii 380-408
- LOEFFLER 1886 Die Aetiologie der Rotzkrankheit Arb a d k Gsndhtsamte, Berl 1 143
- LOHNIS, F Handbuch der landwirtschaftliche Bakteriologie
1913 Vorlesungen über landwirtschaftliche Bakteriologie
- LOHNIS, F, AND HANZAWA, J 1914 Die Stellung von *Azotobacter* im System Centralbl f Bakteriöl, Jena 2 Abt 42 1-8
- LOTSY, J P 1907 Vorträge über botanische Stammesgeschichte Vol I Algen und Pilze
- LOWI, EMIL 1920 Ueber den *Bacillus crassus* Lipschutz Wien klin Wchnschr 33 730-733
- LUDWIG, FRIEDRICH 1892 Lehrbuch der niederen kryptogamen
1898 Abstract of paper by Hoyer (1898) Centralbl f Bakteriöl, Jena 2 Abt 4 867-875
1899 Beobachtungen über Schleimflüsse der Baume im Jahr 1898 Ztschr f Pflanzenkr, Stuttgart 9 10-14
- LUDERS, JOH 1866 Ueber Abstammung und Entwicklung des *Bacterium Termo* Duj, *Vibrio lineola* Ehr Bot Ztg 24 33-39, 41-46
- LUERSSSEN, ARTUR, AND KUHN, M 1908 Yoghurt die bulgarische Sauermilch Centralbl f Bakteriöl, Jena Abt 2 20 241
- LUERSSSEN, CHR 1879 Botanik 1 22
- LUTZ, ADOLPH 1886 Zur Morphologie des Mikroorganismus der Lepra Separat Abdruck aus Dermatologische Studien Herausg von Dr P G Unna Heft 1 p 22
- LYALL, H W 1915 The types of pneumococci in tuberculous sputum Jour Exper M Balt 21 146
- LYNGBYE, H CHRISTIAN 1820 Hydrophyt Danica Account of a new arrangement of the Algae Edinb Phil Jour 2 282-287
- MAASSEN, ALBERT 1905 Über Gallertbildungen in den Saften der Zuckerfabriken Ein Beitrag zur kenntniss der gallertbildenden Bodenbakterien Arb a d k Gsndhtsamte, Berl Abt 5 1
- MACÉ, E 1897 Traité pratique de Bactériologie 3rd ed
1913 Traité pratique de Bactériologie 6th ed
- MACNAB, W R 1877 On the classification of the vegetable kingdom Jour Bot London 6 340
- MAGGI, LEOPOLDO 1886 Essai d'une classification protistologique des ferments vivants Jour Microg 10 80-85, 173-178, 327-333
1887 Acque potabili
- MAGNIN, A 1878 Les bactéries Paris
- MAIRE, RENÉ AND TISON, ADRIEN 1909 La Cytologie des Plasmodiophoracées et la classe des Phytomyxinae Ann Mycolog, Berl 7. 241

- MAKRINOV, I A 1916 Sur un nouveau microorganisme provoquant la fermentation de l'amidon, et des matières pectiques. Arch d Sci Biol de Petrograd 18 440-452
- MALERBA, P, AND SANNA-SALARIS, G 1888 Ricerche sul Gliscrobatterio Rendiconto della R. Accademia della Science fis e mat di Napoli Fasc 6 1888 according to Buchner Centralbl f Bakteriöl, Jena 4 486.
- MARPMANN, G 1889 Ueber die Erreger der Milchsäure-Gährung Ergänzungshefte zum Centralblatt für allgemeine Gesundheitspflege 2: 118-132
- MATZUSCHITA, T 1902 Bakteriologisches Diagnostik
- MAZE, N 1903 Sur la fermentation forménique et le ferment qui la produit Compt rend Acad d sc Par 137 887
- McNAB, A 1904 Ueber den "*Diplobacillus liquefaciens* (Petit)" und ueber sein Verhältnis zudem Morax-Axenfeldschen Diplobazillus der Blepharokonjunctivitis Klinische Monatsblätter für Augenheilkunde 42 54-70
- MENDEL, JOSEPH 1919 *Cladothrix* et infection d'origine dentaire (*Cladothrix matouchoti*) Compt rend Soc de biol, Par 82 583-686
- MERRILL, E D, AND WADE, H W 1919 The validity of the name *Discomyces* for the genus of fungi variously called *Actinomyces*, *Streptothrix* and *Nocardia* Philippine Jour Sc, Manila 14 55-69
- MESSEA, AL 1891 Contribuzione allo studio delle ciglia dei batterii e preposta di una classificazione Rivista d'Igiene e Sanità Pubblica Anno I No 14 Arch Sci Med (Turin), Torino e Palermo 15 233-236
- METCHNIKOFF, ELIE 1888 *Pasteuria ramosa*, un représentant des bacteries a divisione longitudinale Ann de l'Inst Pasteur, Par 2 166
- 1888 Ueber die phagocytaire Rolle der Tuberkelriesenzellen Virchow's Archiv 113 63-94
- 1889 Contributions a l'étude du pleomorphisme des bacteriens Ann de l'Inst Pasteur, Par 3 61
- METTENHEIMER, C 1858 Ueber *Leptothrix ochracea* Kutz, und ihre Beziehung zur *Gallionella ferruginea* Ehr Abhandl herausgeben von der Senckenburgischen naturforsch Ges Frankfurt a M 2 139
- MEYEN, J 1827 *Actinomyce* (Strahlenpilz), eine neue Pilzgattung Linnæa 2 433
- 1839 Jahresberichte über die Resultate der Arbeiten im Felde der physiologischen Botanik de Jahre 1838 Weigmann's Arch Naturgesch 2 1-152
- MEYER, ARTHUR 1897 Studien über die Morphologie und Entwicklungsgeschichte der Bakterien, ausgeführt an *Astasia asterospora* A M and *Bacillus tumescens* Zopf Flora 84 185
- 1898 Neues über die Morphologie der Batterienzelle und die Entwicklungsgeschichte der Batteriensporen Sitzungsberichte der Gesellschaft zur Beförderung der gesammten Naturwissenschaftler zu Marburg No 5 p 49
- 1912 Die Zelle der Bakterien, Jena
- MEYER, K F, AND SHAW, E B 1920 A comparison of the morphologic cultural and biochemical characteristics of *B abortus* and *B Melitensis* Studies on the genus *Brucella* nov gen I Jour Infect Dis, Chicago 27 173-184

- MIEHE, HUGO 1908 Beitrage zur Biologie, Morphologie und Systematik des Tuberkelbazillus Ztschr f Hyg u Infektionskrankh, Leipz 62 131
- MIGULA, W 1890 Bakterienkunde fur Landwirte Berlin
- 1894 Ueber ein neues System der Bakterien Arb Bact Inst Karlsruhe 1 235-238
- 1895 *Schizomycetes* Engler and Prantl Naturlichen Pflanzenfamilien p 20
- 1897 Ueber *Gallionella ferruginea* Ehrenberg Ber d deutsch Bot Gesellsch. Berl 15 321
- 1897-1900 System der Bakterien Vol 1 1897, Vol 2 1900
- 1904 Allgemeine Morphologie, Entwicklungsgeschichte, Anatomie und Systematik der Schizomyceten in Lafar Handbuch der technischen Mykologie 2nd ed p 1-149
- 1904 Botanisches Vademecum Wiesbaden
- MILLER, W D 1886 Einige gasbildende Spaltpilze des Verdauungstractus, ihr Schicksal im Magen und ihre Reaction auf verschiedene Speisen Deutsche med Wehnschr, Berl and Leipz 12 117-119
- 1888 Beitrage zur Kenntniss der Mundpilze Deutsche med Wehnschr, 14 612
- 1892 Mikroorganismen der Mundhohle Ed 2
- MIQUEL, M P 1879 Etudes sur la fermentation ammoniacale et les ferments de l'uree Ann Microg 1 414, 417, 506, 552
- 1880 Etudes sur la fermentation ammoniacale et les ferments de l'urée Ann Microg 2 13, 53, 122, 145, 367, 488
- 1881 Etudes sur la fermentation ammoniacale et les ferments de l'urée Ann Microg 3 275, 305
- MIQUEL, P, AND CAMBIER, R 1902 Traité de Bact Paris p 881
- MIYOSHI, MANABU 1897 Studien uber die Schwefelrasenbildung und die Schwefelbakterien der Thermen von Yumoto bei Nikko Jour Col Sci Imp Univ Tokyo 10 143-170
- MOLISCH, HANS 1906 Zwei neue Purpurbakterien mit Schwefelkorperchen Bot Ztg Abth I Orig 64 223-232
- 1907 Die Purpurbakterien Jena
- 1909 Festschrift fur Treub Extrait des Ann Jard Bot Buitenzorg Series 2 Supp 3 p 29-33
- 1910 Die Eisenbakterien Jena
- MORAX, V 1896 Note sur un diplobacille pathogene pour la conjonctivite humaine Ann de l'Inst Pasteur, Par 10 337
- MORENO, J MADRID 1901 Eine neue Art von *Ascobacillus*, entdeckt im Wasser des Lozayakanals bei Madrid Centralbl f Bakteriol Jena 30 111-114
- MORI, N 1913 Di un nuovo batterio patogene e di molti altri batteri nei quali quo provocarsi l'individuazione di un nucleo tipico Ann d Staz Sper p l malattie infettive del bestiame 1 1
- MORREN, CHARLES F A 1831 Mémoire sur les vibrions lamellinaires Mes-sager des sciences de Gand 9^e et 10^e liv 6^e vol 1830 Abstract Bull des sciences naturelles et de Geologie 27 203-205

- MUELLER, O F 1773 Vermium Terrestrium et fluviatiliu seu Animalium infusoriorum, helminthicorum et Testaceorum, non marinorum Vol 1
- MUELLER, O F 1786 Animalcula infusoria et marina, Hauniae
- MUHLHAUSER, F A 1884 Ueber Spirillen Arch f path Anat u Phys (Virchow) 47 84-106
- MUSGRAVE, W E, CLEGG, M T, AND POLK, MARY 1908 Streptothricosis with special reference to the etiology and classification of mycetoma Phillip Journ Sci, Ser B, Med Sci 3 447-544
- NADSON, G A 1906 Zur morphologie der niederer Algae I Über Veraenderungen bei *Stichococcus bacillaris* Nag in Abhangigkeit om den Bedingungen der Ernahrung II Über Endosporenbildung bei *Stichococcus bacillaris* Nag und *Chloroidium Krugeri* III *Chlorobium limicola* ein gruner chlorophyll-fuehrender Mikrobe Bull du Jard in Imp Bot de St Petersburg 6. 184-192
- NAEGELI, CARL WILHELM VON 1842 Botanische Beitrage Linnaea 16 279 Jahrb Wiss Bot (Pringsheim)
- 1857 Bericht uber die Verhandlungen der bot Section der 33 Versammlung deutscher Naturforscher und Arzter Bot Ztg p 760
- NICOLLE, M, AND CESARI, E 1914 Études sur les staphylocoques dorés Ann de l'Inst Pasteur, Par 28 219
- NIJDMAN, H W M 1907 *Aerobacter tartarivorum* (Dutch) Delft (vanMarken)
- NIKLEWSKI, BRONISLAW 1910 Ueber die Wasserstoffoxydation durch Mikroorganismen
- NISSEN, FRANZ 1889 Zur Kenntniss der bacterienvernichtenden Eigenschaft des Blutes Ztschr f Hyg u Infektionskrankh, Leipz 6 487-520
- NOGUCHI, HIDEYO 1917 *Spirochaeta icterohaemorrhagiae* in American wild rats and its relation to the Japanese and European strains Jour Exper Med, Balt 25 755-763
- NOGUCHI, HIDEYO, AND AKATSU, SEINAI 1917 Immunological studies on pure cultures of various spirochetes Jour Exper Med, Balt 25 765-788
- NOGUCHI, HIDEYO 1918 Morphological characteristics and nomenclature of *Leptospira (Spirochaeta) Icterohaemorrhagiae* (Inada and Ido) Jour Exper Med, Balt 27 575-592
- 1918 Further study on the Cultural conditions of *Leptospira (Spirochaeta) Icterohaemorrhagiae* Jour Exper Med, Balt 27 593-608, 609-625, 667-678
- OERSTED, A S 1840-1841 Beretning om en Excursion til Trindelen en Alluvial dannelsen i Odensefjord Naturhistorisk Tidsskrift 2 552-569
- 1844 De regionibus marinis p 44
- OGSTON, ALEX 1883 Micrococcus Poisoning Jour Anat et Physiol (Paris) 17 27
- OLSEN-SOPP, O J 1912 Taette, die urnordische Dauermilch und verwandte Milchsorten, sowie ihre Bedeutung fur die Volksernahrung Centralbl f Bakteriologie, Jena Abt 2 33 1-54
- ORLA-JENSEN, S 1909 Die Hauptlinien des natuerlichen Bakterien systems Centralbl f Bakteriologie, Jena Abt 2 22 305-346

- 1909 Vorschlag zu einer neuen bakteriologischen Nomenklatur Centralbl f Bakteriöl, Jena Abt 2 24 477
- 1919 The lactic acid bacteria Mém de l'Acad Roy d Sci et d Lettres de Danemark, Copenhagen, 1919 Sec d Sci 8me, sér 5. (no 2) 81-196
- 1921 The main lines of the natural bacterial system Jour Bact, Balt 6: 263-273
- OVERBECK, A 1891 Zur Kenntniss der Fettfarbstoff Production bei Spaltpilzen Nova Acta d k Leop Carol deutsch Akad d Naturf 55 399
- PASTEUR, LOUIS 1868 Etudes sur la vinaigre
- PERFILIEV, B V 1914 Sur une "bacterie verte" à chlorophylle, le *Pelodictyon clathratiforme* Lauterb Jour Microbiol 1 179-194
- PERRONCITO, EDUARD 1879 Ueber das epizootic Typhoid der Hühner Arch f wissensch u prakt Tierh, Berl 5 22-51
- 1879 Ueber den *Actinomyces bovis* und die Sarkome der Rinder Deut Ztschr Thiermed 5 31-40
- PERRONE 1905 Contribution a l'étude de la bacteriologie de l'appendicite Ann de l'Inst Pasteur, Par 19 371
- PERSOON 1822 Mycologia europaea Sectioprima
- PERTY, MAXMILIAN 1852 Zur Kenntniss kleinster Lebensformsn
- PETRUSCHKY 1913 Die pathogenen Trichomyceten und Trichobakterien *Streptothrix*, *Cladothrix*, *Leptothrix* Kolle and Wassermann Handbuch der pathogene Mikroorganismen 2nd ed 5 267-300
- PETSCHENKO, BORIS DE 1908 Sur la structure et le cycle évolutif de *Bacillopsis styloggae* n g et n sp Bull Acad Sci Cracovie pp 359-371
- 1911 *Drepanospira Mulleri* n g n sp parasite des paraméciums contribution a l'étude de la structure des bacteries Arch f Protistenk, Jena 22 248-298
- PFEIFFER. 1887 Synonymia botanicae
- PINOY, E, AND MORAX 1911 Sur les concrétions des voies lacrymales Etud mycologique Bull Soc Ophtalmologie de Paris No 3
- 1913 Actinomycoses et Mycetomes Bul Inst Pasteur 11 929
- POULSEN, V A 1880 Ueber einige mikroskopische Pflanzenorganismen Bot Ztg 38 509
- PRAZMOWSKI, ADAM 1879 Zur Entwicklungsgeschichte und Fermentwirkung einiger Bakterien-Arten Verlauffe Mitt Bot Ztg 37 409-424
- 1880 Untersuchungen über die Entwicklungsgeschichte und Fermentwirkung einiger Bakterien p 23
- PREISZ, HUGO 1894 Recherches comparatives sur les pseudotuberouloses bacillaires Ann de l'Inst Pasteur, Par 8 231
- PRILLIEUX 1879 Sur la nature et la cause de la formation des tubercules quinaissent sur les racines des Legumineuses Bul Soc Bot France 26 98
- RABENHORST, L 1864-1865 Flora Europaea Algarum
- RALFS, JOHN 1843 On the British Diatomaceae Ann and Mag Nat Hist Ser 1 12 351
- REICHENBACH, H L 1852 Gartenorchiden Bot Ztg 10 928

- REITZ, A 1906 Weitere bakteriologische Untersuchungen mit der Stuttgarter Markt- und Handelsbutter Centralbl f Bakteriöl Jena 2nd Abt 16 719
- RIST, E, AND KHOURY, J 1902 Etudes sur un lait fermenté comestible, le Leben d'Egypte Ann de l'Inst Pasteur Par 16 65-84
- RIVOLTA, S 1878 Sul così detto mal del rospo del Truta e Sull "*Actinomyces bovis*" di Harz Clin vet, Milano 1 169 and 201
- 1882 Ueber die Priorität der Beschreibung der Formen der Actinomykose und ihrer eigenthümlichen Elemente bei den Rindern (Virchow) Archiv f path Anat u Phys u f klinische Medecin 88 389-390.
- 1888 Pleurite infettiva nel gatto e nel cane Giornale di Anatomia Fisiologia e Patologia degli Animal 20 1-8
- ROBIN, CHARLES 1847 Des végétaux qui croissent sur les animaux vivants Paris
- 1853 Histoire naturelle des Végétaux Parasites qui croissant sur l'homme et sur les animaux vivants Paris
- ROSENBAACH, A J F 1884 Mikroorganismen bei den Wund Infektionskrankheiten
- 1909 Experimentelle, morphologische und klinische Studie über die krankheitserregenden Mikroorganismen des Schweinesrothlaufs, des Erysipeloides und der Mausesepsis Ztschr f Hyg u Infektionskrankh, Leipz 63 343-371
- ROTHEMANN, E A 1904 *Glioscrobacterium* als Ursache der schleimigen Garung des Menschurins Centralbl f Bakteriöl, Jena Abt 1 Originale 37 491
- ROUGENTZOFF, D 1914 La Flore intestinale des lapins nourris de carottes et des lapins soumise l'inanition Ann de l'Inst Pasteur, Par 28 639-661
- ROZE, E 1898 Sur un nouveau type générique des schizomycètes, le *Chatinella* Compt rend Acad d sc, Par 127 825 Bull de la soc mycol de France p 69
- RUATA, VIKTON 1909 Ueber einen Kokkobacillus der menschlichen Bindehaut Ein Beitrag zur Kenntniss der gram-negativen Diplokokken Centralbl f Bakteriöl, Jena Abt I Originale 51 630
- RULLMANN, W 1901 Ueber ein *Nitrosobacterium* mit neuen Wuchsformen Centralbl f Bakteriöl, Jena Abt 2 3 228
- SALISBURY, J H 1868 Microscopic examination of the blood, and vegetations found in variola, vaccinia and typhoid fever
- SAMBON, L 1907 In Manson's, Tropical Diseases
- SAMPIETRO, G 1908 Sulla vitalità e virulenza die batteri intestinali conservati nel terreno Ann d'Igiene Roma 18 119-135
- 1908 Sopra due casi di actinomiosi nell'uomo Ibid 391-461
- SAUVAGEAU, C, AND RADAIS, M 1892 Sur les genres *Cladothrix*, *Streptothrix*, *Actinomyces* et description de deux *Streptothrix* nouveaux Ann d l'Inst Pasteur, Par 6 242-273
- SCHAMBERG, F 1902 On the *Microbacillus seborrhoeae* (Sabourand) Zurn dermatol sifilidol Russ trans St Petersburg 2 293

- SCHATTENFROH, A, AND GRASSBERGER, R 1899 Ueber Buttersauregahrung
Centralbl f Bakteriöl, Jena 2nd Abt 5 697-702
- SCHAUDINN, F 1905 "Korrespondenzen" Deutsche med Wehnschr, Berl
and Leipz 31 1728
- SCHAUDINN UND HOFFMANN 1905 Vorläufiger Bericht über das Vorkommen
von Spirochäten in syphilitischen Krankheitsproduktion und bei Papillo-
men Arb K Gsndhtsamt 22 527
- SCHIEWIAKOFF, W 1893 Ueber einen neuen bacterienähnlichen Organismus des
Susswassers Heidelberg
- SCHINDLER, ANTON K 1905 Halorrhagidaceae Das Pflanzenreich Heft
23 82
- SCHLEGEL, M 1913 Morphologie des Aktinomycespilzes Kolle u Wasser-
man, Handbuch der pathogenen Mikroorganismen 2nd ed 5 303
- SCHMIDLECHNER, KARL 1905 Ein durch Vincentsche Bakterien verursachte
Puerperalerkrankung Zeitschr. für Geburtshilfe und Gynakologie
56, 291
- SCHMIDT AND WEIS 1902 Die Bakterien
- SCHNEIDER, A 1912 Pharmaceutical Bacteriology
- SCHORLER, B 1904 Beiträge zur Kenntniss der Eisenbakterien Centralbl
f Bakteriöl, Jena Abt 2 12 691
- SCHROETER, J 1872 Ueber einige durch Bakterien gebildete Pigmente Cohn's
Beiträge z Biologie d Pflanzen Heft 2 1 109-126
- 1886 *Schizomycetes* Cohn's Kryptogeme-Flora von Schlesien 3 136-174
- SCHUETZ 1887 Di Ursache der Brustseuche der Pferde Arch f wissensch
u prakt Tierh, Berl 13 27
- SCHWERS, HENRI 1908 Le fer dans les eaux souterraines Rev d'hyg, Par
30 11-43, 100-126, 185-221, 287-306
- SELIGMANN, E 1911 Bakteriologische Befunde bei Sauglingsgrippe Beiheft
Centralbl f Bakteriöl, Jena Abt 1 Referate 50 81*-82*
- SETTE, VINCENTO 1824 Sull'arrossimento straordinario di alcune sostanze
alimentose osservato nella provincia di Padova l'anno 1819 Venezia
1824
- SMITH, ALLAN J 1902 Laboratory Exercises in Bacteriology
- SMITH, ERWIN F 1905 Bacteria in Relation to Plant Diseases Vol 1 Car-
negie Institute Washington, D C
- SMITH, R GRIEG 1902 The gummosis of the sugarcane (*Boc vascularum*
Cobb) Sydney, N S W, Proc Linn Soc N S Whales 27 31-47
- 1902 An *Ascobacterium* from the sugar cane Ibid pp 137-145 Int
Cat Sci Lit 2 157-158
- 1902 Die Gummigärung von Zuckerrohrsaft Das Verderben der Krystalle
von rohem und raffiniertem Zucker Die Sauregärung des Rohrzuckers
Centralbl f Bakteriöl, Jena Abt 2 8 597-600
- SOLLIED, PETER RAV 1903 Studien über den Einfluss von Alkohol auf die an
verschiedenen Brauerei- und Brennereimaterialien sich vorfinden
Organismen, sowie Beschreibung einer gegen Alkohol sehr widerstands-
fähigen neuen *Pediococcus* art (*Pediococcus hennebergi*, n sp) Ztschr
Spiritusindus 26 481-482, 491-493

- STERNBERG, G M 1892 Manual of Bacteriology
- STILES, C W 1905 The international code of zoological nomenclature as applied to medicine Bull Hygienic Laboratory, Washington, D C 24
1905 The correct spelling of *Spirochaeta* Am Med 10 94
- STILES, C W, AND PFENDER, C A 1905 The generic name of the parasite of syphilis Am Med 10 936
- STOCKHAUSEN, FERD 1907 Okologie, "Anhaufungen" nach Beijerinck Beiträge zur natürlichen Reinzucht der Mikroorganismen Berlin
- STOKLASA, JULIUS 1908 Beitrag zur kenntniss der chemischen Vorgänge bei der Assimilation des elementaren Stickstoffs durch *Azotobacter* und *Radobacter* Centralbl f Bakteriöl, Jena Abt 2 21 282-509
- STREINZ, W M 1862 Nomenclator fungorum
- STUTZER, A, AND HARTLEB, R 1899 Untersuchungen über die beider Bildung von Salpeter beobachteten Mikroorganismen Mitt Landw Inst Breslau 1 75, 2 197
- SWELLENGREBEL, M. 1907 Sur la cytologie comparée des Spirochètes Ann d l'Inst Pasteur, Par 21 448 and 562
- TERNI, C 1894 *Actinomyces Gruberii* Atti dell'vi congresso med internaz Roma 6 (Igiene) 79
- THAXTER, ROLAND 1892 On the *Myzobacteriaceae*, a new order of *Schizomycetes* Bot Gaz, Chicago 17 394
- THESING, CURT 1906 *Spirochaete*, *Spironema* oder *Spirillum* Centralbl f Bakteriöl, Jena Abt I 40 351
- THIERCELIN, E 1899 Du diplocoque ou enterocoque son rôle dans la pathogenie de certaines affections digestives Bull Soc de Péd de Paris 1 208-223
1899 Zur un diplocoque saprophyte de l'intestin susceptible de devenir pathogénique Compt rend Soc de biol, Par 51 269-271
1902 Procèdes faciles pour isoler l'entérocoque des selles normale, filtration des selles, culture préalable en anaérobie Compt rend Soc de biol 54 1082
- THIN, GEORGE 1880 On *Bacterium foetidum*, an organism associated with profuse sweating from the soles of the feet Proc Roy Soc, Edinb 30 473-478
1881 *Bacterium decalvans*, an organism associated with the destruction of the hair in alopecia areata Proc Roy Soc, Edinb. 33 247-253
- THOMSON, ROBERT D 1852 Ueber die Natur und die chemischen Wirkungen der Essigmutter Annalen der Chemie und Pharmazie 83 89
- TIEGHEM, PH VAN Observations sur des bactériacées vertes, sur des phycochromacees, blanches, et sur les affinités de ces deux familles Bul. Soc Bot France 27 174-179
- TILDEN, JOSEPHINE 1910 Minnesota Algae 1 86
- TISSIER, HENRY 1900 Recherches sur la Flore Intestinale des Nourrissons (Etat normal et pathologique) Thèse pour le Doctorat en Médecine. Paris
1905 Répartition des microbes dans l'intestin du nourrisson Ann d l'Inst. Pasteur, Par 19 109-123

- TRÉCUL, A 1865 Matière amylacée et cryptogames amylières dans les vaisseaux du latex de plusieurs Apocynées Compt rend Acad d sc Par 61 156-160
- 1865 Production de plantules amylières dans les cellules végétales pendant la putréfaction Chlorophylle cristallisée Compt rend Acad d sc Par 61 432-436
- TRELEIGH, W 1885 Observations on several zoözoë and related forms Studies Biol Lab Johns Hopkins Univ 3 194-216
- TREVISAN, V 1842 Prospetto della Flora Euganea
- 1879 Prime linee d'introduzione allo studio dei Batteri italiani Rendiconti Reale Istituto Lombardo di Scienze e Lettere Ser II 12 13-21
- 1879 Prime linee d'introduzione allo studio dei Batteri italiani 2nd communication Rendiconti Reale Istituto Lombardo di Scienze e Lettere IV Ser II 12 13-21, 133-151
- 1879 Sulla causa dell'asfissia e dell'agglutianzione dei globuli del sangue nelle difterite Rendiconti Reale Istituto Lombardo di Scienze e Lettere Ser II 12 54-62
- 1882 Emilio Cornalia Atti della Accademia Fisio-Medico-Statistica di Milano p 100
- 1882 Processi Verbal Delle Adunanze Atti della Accademia Fisio-Medico-Statistica di Milano p 151
- 1883 Il Batterio del tifo abdominale (*Metallactar ileotyper*) e il Batterio della pellagra, (*Bacterium Maydis*) Atti della Accademia Fisio-Medico-statistica in Milano Ser 4 1 149-157
- 1884 A proposito del bacillo del cholera, Koch o Pacini? intorno al modo di agire del bacillo nel corpo umano Gazz med lomb, Milano 8 s 6 373, 547
- 1884 Il batterio del cholera è un vibrione, non un bacillo, è agente esogeno e causor occasionale della malattia, non agente endogeno e causa assoluta Gazz med lomb, Milano 8s 6 417-427
- 1885 Il fungo del cholera asiatico Questioni risolte Atti della Accademia Fisio-Medico-Statistica in Milano Ser 4 3 78-91
- 1885 Caratteri di alcuni nuovi generi di Batteriacee Atti della Accademia Fisio-Medico-Statistica in Milano Ser 4 3 92-107
- 1885 Quadro sinottico delle malattie dell'uomo e degli animali caratterizzate dalla presenza di batteri patogeni Atti della Accademia Fisio-Medico-Statistica in Milano Ser 4 3 107-119
- 1887 Sul Micrococco della rabbia e sulla possibilità di riconoscere durante il periodo d'incubazione, dall'esame del sangue della persona morsicata, se ha contratta l'infezione rabbica Rendiconti Reale Istituto Lombardo di Scienze e Lettere Ser II 20 88-105
- 1887 Se sia vero che il *Bacillo imetrofo* (*Micrococcus prodigiosus* di antichi autori) preceda sempre come agente specifico predisponente il calcino del baco da seta Rendiconti Reale Istituto Lombardo di Scienze e Lettere Ser II 20 797-804
- 1887 Caratteri di Alcuni Nuovi Generi di Batteriacee Reale Istituto Lombardo di Scienze e Lettere Rendiconti Pisto p 94
- 1889 *Klebsiella* Reale Istituto Lombardo di Scienze e Lettere. Rendiconti Ser II 30 777-790
- 1889 I generi e le specie delle Batteriacee Milan

- TROILI-PETERSSON, G 1903 Studien über die Mikroorganismen des schwedischen Gunterkases Centralbl f Bakteriöl, Jena Abt 2 11 120-143
- TSILINSKY, P 1889 Sur les Mucédinées Thermophiles Ann de l'Inst Pasteur, Par 13 500
- TURCZANINOW, NICOLAUS 1851 Synanthereae quaedam hucusque indscriptae Bull Soc natur Moscow 24 156
- UEKE, A 1898 Ein Beitrag zur kenntniss der Anaeroben Centralbl f Bakteriöl, Jena Abt I 23 996
- UNNA, P G 1892 Der *Streptobacillus* des weichen Schankers Monats f Prakt Derm 14 485-490
- UNNA, P G, AND TOMMASOLI, P Flora dermatologica (Fortsetzung) Bacillen, kokken und Hefeformen Monatshefte für praktische Dermatologie 9. 60
- VALENTIN, G G 1836 Repert für Anat und Phys 1 110
1837 *Hygrocybus intestinalis*, eine auf der lebendigen und ungestört functionirenden Schleimhaut des Darmcanales vegetirende Conferve Repert f Anat u Phys 1 110-114
- VAN LAER, H 1889 Contributions à l'histoire des ferments des hydrates de carbone Bacille des bières tournées Mem de l'Acad royale de Belgique 43
- VAN TIEGHEM, PH 1878 Sur la Gomme de Sucrierie (*Leuconostoc mesenteroides*) Ann Sci Nat Bot Ser 6 7 198
1879 Sur le ferment butyrique (*Bacillus amylobacter*) à l'époque de la houille Compt rend Acad d Sc, Par 89 1102-1104
1880 Sur quelques bacteries agregées Bul Soc Bot France 27 148-153
1884 Traité de Botanique Paris p 114
- VELEY, V H, AND VELEY, LILIAN J 1898 The microorganism of faulty rum London Frowde Abs Kochs J 9 154
- VINCENZI, L 1897 Di un nuovo tetragenopatogene (tetragenocitreo) Riforma Med 13 758-760
- VON BAER, K E 1827 Die Verwandtschafts-verhältnisse unter den Niedern Thierformen Nova Acta Phys-Med Acad Caes Leop Nat Cur 13 748
- VUILLEMIN, PAUL 1888 Les tubercules radicaux des légumineuses Ann de la Sci Agron France et Etrangère 4 121-212
1905 Sur la dénomination de l'agent de la syphilis Compt rend Acad d sc, Par 140 1567
1913 Genera Schizomycetum Ann Mycologie, Berl 11 546
- WAKSMAN, SELMAN, A 1919 Cultural studies of species of *Actinomyces* Soil Sc, New Brunswick, N J 8 71-207
- WAKSMAN, SELMAN A, AND JOFFE, J S 1922 Microorganisms concerned in the oxidation of sulfur in the soil II *Thiobacillus thiooxidans*, a new sulfur-oxidizing organism isolated from the soil Jour Bact, Balt, 7 239-256

- WALLROTH, FRED GUIL 1837 *Flora Cryptogamica German* 2 182
- WARMING, EUGENIUS 1875 *Om nogle ved Danmarks Kysterlevende Bakterien Videnskab Meddelelser fra den naturhist Forening i Kjobenhavn* 7 307-420
- 1876 *Observations sur quelques Bactéries qui se rencontrent sur les côtes du Danmark Résumé d'un mémoire publié dans les "Videnskabelige Meddelelser" de la Société d'histoire naturelle de Copenhague, Avec 4 Planches Den Naturhistoriske Forening Kjobenhaven Videnskabelige Meddelelser Appendix* 28 1-36
- WEICHELBAUM, A 1886 *Untersuchungen uber Pneumonie Bakt Theil Medizinische Jahrbucher* 82 Neue Folge 1 506
- 1887 *Zur Aetologie der acuten Endocarditis Centralbl f Bakteriologie, Jena* 2 209-217
- WENNER, J J, AND RETTGER, LEO F 1919 *A systematic study of the proteus group of bacteria Jour Bact, Balt* 4 331-353
- WEST, G S 1904 *The British Freshwater Algae*
- WEST, G S, AND GRIFFITHS, B M 1909 *Hillhousia mirabilis, a giant sulfur Bacterium Proc Roy Soc Long Biol Sci* 81 398-399
- WILLE, N. 1903 *Algologische Notizen x. Nyt Magazin f Naturvidenskaben* 41 89-185
- WINOGRADSKY, SERGIUS 1888 *Beitrage zu Morphologie und Physiologie der Bakterien Heft 1 Zur Morphologie und Physiologie der Schwefelbakterien Leipzig*
- 1892 *Contributions à la morphologie des organismes de la nutrition Arch Sci Biol (St Petersburg)* p 87
- WINSLOW, C E A, BROADHURST, JEAN, BUCHANAN, R E., KRUMWIEDE, CHARLES JR, ROGERS, L A, AND SMITH, G H (Committee of the Society of American Bacteriologists on Characterization and Classification of Bacterial types) 1917 *Preliminary report The families and genera of bacteria Jour Bact, Balt* 2 552
- 1920 *The families and genera of bacteria Final report of the Committee of the Society of American Bacteriologists on Characterization and Classification of Bacterial Types Jour Bact, Balt* 5 191-229
- WINSLOW, C E A, AND ROGERS, ANNE F 1905 *A revision of the Coccaceae Science, N S* 21 669-672
- 1906 *A statistical study of generic characters in the Coccaceae Jour Infect Dis, Chicago* 3 485-546
- WINSLOW, C E A, ROTHEBERG, W, AND PARSONS, E I 1920 *Notes on the classification of the white and orange Staphylococci Jour Bact, Balt* 5 145-167
- WINSLOW, C E A, AND WINSLOW, ANNE ROGERS 1908 *The Systematic Relationships of the Coccaceae*
- WINTER, GEORG 1880 *Die Pilze Rabenhorst's Kryptogamen flora* 2nd ed Vol 1
- WISLITCH, S M 1914 *Spirillum Kolhutzii, nov sp, und einige neue Schwefelbakterien Prof Molisch's Jour de Microbiologie (Russian)* 1 50.
- WOLFF, MAX 1907 *Pedioplane haechelii n g, n sp, und Planosarcina schaudinnii n sp Centralbl f Bakteriologie, Jena Abt* 2 18 9-26

- WOLLMAN, EUGENE 1912 Recherches sur les Microbes Amylolytiques de l'intestin Ann de l'Inst Pasteur, Par 26 610-634
- WRIGHT, JAMES HOMER 1904-1905 The biology of the microorganisms of Actinomycosis Jour Med Research New Series 8 349-350
- ZEIDLER, A 1896 Ueber eine Essigsäure bildende Thermobacterie Centralbl f Bakteriöl, Jena Abt II 2 739
- ZETTNOW, 1903 Beiträge zur Kenntnis von *Spirobacillus gigas* Festschrift z 60 Gebrütern v Robert Koch, Jena p 383-388
- ZOPF, W 1883 Die Spaltpilze Encyklop der Naturwiss Abth 1 Lief 32
1884 (3) Die Spaltpilze Auflage II
1885 Die Spaltpilze 3rd ed
1891 Ueber Ausscheidung von Fettfarbstoffen (Lipochromen) seitens gewisser Spaltpilze Ber d deutsch Bot Gesellsch, Berl 9 28
- ZUELZER, MARGARETE 1911 Ueber *Spirochaeta phacalis* Ehrbg und deren Verwandtschaftsbeziehungen Arch f Protistenk, Jena 24 51
- ZUKAL, HUGO 1885 Neue Pilze aus Nieder-Oesterreich Verhandl K K Zool Bot Gesell Wien p 333-342
1896 *Myxobotrys variabilis* Zuk als Repräsentant einer neuen Myxomyceten Ordnung Ber d deutsch Bot Gesellsch, Berl 14 346
1897 Ueber die Myxobakterien Ber d deutsch Bot Gesellsch 15 550

SUBJECT INDEX

- Acanthococcus*, 383
- Acetimonas*, 69, 152, 154, 434
- aceti*, 152
- schuetzenbachii*, 152, 153
- Acetobacter*, 96, 98, 103, 153, 389, 400, 519
- aceti*, 153, 154
- melanoginum*, 153
- pastorianus*, 153
- placatum*, 153
- Acetobacterium*, 153, 154
- xylinum*, 153, 154
- xylinum* var *Lagerheimii*, 154
- Achromatiaceae*, 84, 154, 156, 330, 502, 507, 510
- Achromatum*, 84, 154, 155, 156, 330, 381, 504
- Mulleri*, 155
- oxahferum*, 155, 156
- Achromobacter*, 105, 156
- liquefaciens*, 156
- Achromobactereae*, 104, 105, 156
- Acidobacteriaceae*, 69, 71, 156, 157, 225, 247, 287, 419, 431
- Acid fast group, 67
- Actinobacille*, 157
- Actinobacillosis*, 158
- Actinobacillus*, 72, 83, 96, 97, 106, 157, 158, 159, 161
- ligneresi*, 158
- paulotrophus*, 159
- Actinobacilo*, 157, 158, 159
- Actinobacillosis*, 158
- Actinobacter*, 26, 159, 160, 161, 289
- polymorphum*, 26, 160
- Actinobacterium*, 63, 159, 161
- israeli* var *Spitzeri*, 161
- lactis viscosum*, 161
- Actinocladothrix*, 161
- Actinococcus*, 161, 162
- cyaneus*, 161
- Actinomyce*, 162, 165, 166
- horkelii*, 162
- Actinomyces*, 24, 34, 45, 58, 67, 68, 69, 74, 83, 87, 89, 96, 106, 157, 159, 161, 162, 163, 164, 165, 166, 167, 168, 169, 171, 241, 246, 261, 264, 294, 295, 377, 384, 385, 402, 403, 404, 405, 406, 496, 497, 500, 516
- bovis*, 24, 161, 162, 163, 165, 166, 167, 294, 295, 403
- clinik I & II*, 161
- musculorum*, 163
- paulotrophus*, 159
- Actinomycetaceae*, 74, 83, 96, 106, 159, 167, 168, 169, 304, 353
- Actinomycetales*, 79, 96, 102, 106, 167, 385, 461
- Actinomycetes*, 58, 69, 168, 169, 248, 280, 333
- Actinomycosis*, 158
- Aerobacter*, 56, 82, 105, 169, 226, 312, 339, 452
- aerogenes*, 169, 170
- coli*, 169
- liquefaciens*, 169
- tartarivorum*, 169
- viscosum*, 169
- Aerogenesbacterium*, 99, 170, 277
- Aerogenesbacterium aerogenes*, 170
- Aethyl-Bacillus*, 170
- Agonium*, 170
- centrale*, 171
- Altinomyces*, 171.
- Albococcus*, 61, 65, 87, 90, 92, 96, 171, 172, 173, 193, 411, 488, 489, 492
- albus*, 172, 173
- canescens*, 171, 172
- candicans*, 171, 172
- candidus*, 171
- epidermidis*, 171, 173
- pyogenes*, 171, 172, 173
- rhenanus*, 171, 172
- tetragenus*, 171
- ureae*, 171
- Alcaligenes*, 93, 95, 105, 173, 298

- Alcaligenes faecalis*, 173
ivar, 173
Alkalibacteriaceae, 69, 71, 173, 202, 358, 419, 500, 519
Alkaligenes, 173, 243
faecalis, 243
Allococcaceae, 51, 60, 174
Amarantaceae, 260
 American Code, 145
Amoebobacter, 36, 42, 55, 70, 85, 174, 175, 176, 186, 508
bacillosus, 174, 175
granula, 174, 176
roseus, 174, 175
Amoebobacteriae, 176
Amoebobacteriaceae, 55, 63, 175, 176
Amoebobacteriae, 42
Amoebobacteriae, 84, 85, 174, 176, 186, 256, 506, 508, 512
Amoebobacterinae, 176
Amoebobacterioidae, 176
Amoebomonas, 70, 176
Amphitricha, 177
Amylobacter, 19, 177, 268, 269, 287, 520
butylicus, 177
ethylicus, 177
Amylobakter, 178, 268
Anabaena, 23
Anacardiaceae, 449
Anacystis, 37
 Anthrax group, 66
Aphaneroglia, 178, 433
aquae potabilis, 178
Aphanocapsa, 22, 37, 502, 503
Aphanothece, 22, 37, 511
clathratiformis, 417
luteola, 461
Aplanobacter, 61, 178, 179, 211, 225, 429, 430
Arlingnillus, 102, 179, 267, 370
Chauvoet, 102, 179
Arthrobacillus, 179
Arthrobacter, 48, 179
Arthrobacteriaceae, 179, 181
Arthrobacterium, 29, 180, 181, 199
aceti, 180
Arthrobactridium, 49, 180
Arthrobactrillum, 49, 180, 232
Arthrobactrinum, 49, 180, 233
Arthrobacteriaceae, 181
Arthrobacterium, 179, 181
Arthrobokkaceae, 181
Arthromitus, 18, 181, 182, 183
batrachorum, 183
cristatus, 181, 182, 332
nitidus, 181, 182
Arthrospirobacterium, 183
Arthro-Spirobacterium, 184
Arthrospirobacteriaceae, 184
Arthrostreptococcus, 184
Arthro-Streptokokkus, 181, 184
Ascobacillus, 184, 185, 186, 434
aquatilis, 185, 186
citreus, 184, 185
sacchari, 185
Ascobacteria, 30, 185
ulvina, 185, 186
Ascobacterium, 43, 185, 186
aquatile, 186
luteum, 185, 186
Ascococcae, 35, 41, 176, 186, 250, 274, 306, 315
Ascococcee, 274, 503
Ascococcus, 186, 187, 272
parvus, 187, 188
Ascococcus, 22, 23, 25, 27, 29, 31, 32, 33, 35, 37, 38, 41, 43, 45, 65, 91, 186, 187, 188, 189, 287, 306, 356, 375, 376, 389, 411, 458, 492
Billrothi, 23, 187, 188, 189, 190, 191
Billrothi var *thermophilus*, 189
buccalis, 189
cantabrigdensis, 190
cellaris, 189
equi, 190
mesenteroides, 24, 188, 190, 355, 356
vibrans, 188
Ascokokkus, 191
Askokokkus, 181, 191
Astasia, 53, 81, 191, 203
asterospora, 53, 191
Asterococcus, 72, 81, 191, 329.
mycondes, 192
Astrobacter, 54, 192, 412
Athorhodaceae, 63, 192, 444, 445, 446
Aurococcus, 61, 65, 92, 172, 193, 411, 456, 488, 492
aurantiacus, 193

- Aurococcus aureus*, 193
mollis, 193
 Authority, 111
Azotobacter, 58, 69, 74, 81, 87, 89, 96,
 98, 103, 193, 194, 195, 340, 400, 410,
 444
chroococcum, 194
Azotobacteriaceae, 96, 103, 195, 400, 444
Azotobacteriaceae, 195
Azotomonas, 194, 195, 434

Babesia, 42, 195, 492
erysipeloides, 195
xanthopyretica, 195
Bacillaceae, 51, 59, 60, 66, 73, 87, 88,
 91, 92, 95, 99, 103, 106, 195, 196,
 197, 209, 211, 230, 232, 243, 266,
 269, 297, 298, 299, 309, 322, 325,
 326, 367, 400, 414, 432
Bacillaceae, 195, 197
Bacilleae, 40, 51, 60, 80, 92, 94, 95,
 196, 197, 223, 233, 410, 415, 471
Bacilleae, 197, 200, 234, 267, 278, 360,
 409
Baciller, 48, 196, 197
Bacilococcus, 197
Bacillopsis, 198
stylopygae, 198
Bacillus, 21, 22, 26, 27, 28, 29, 30, 31,
 32, 33, 35, 37, 38, 40, 43, 44, 45, 46,
 47, 48, 50, 51, 53, 54, 57, 58, 59,
 60, 66, 67, 71, 73, 74, 75, 77, 78, 80,
 81, 90, 94, 99, 106, 153, 159, 163,
 173, 178, 191, 194, 196, 197, 198,
 199, 200, 201, 202, 203, 204, 205,
 206, 207, 218, 219, 221, 222, 233,
 225, 231, 234, 235, 257, 267, 275,
 288, 289, 290, 297, 300, 303, 306,
 308, 311, 312, 314, 322, 327, 339,
 347, 369, 370, 371, 382, 392, 420,
 429, 432, 434, 444, 447, 462, 463,
 464, 472, 475, 487, 490, 491, 500,
 517, 518, 519, 520, 525, 526, 531
aceti, 152, 518
acidophilus, 373
aerifoetidus, 463
aerogenes capsulatus, 101, 529
alvei, 278
amylobacter, 177, 267, 268, 269, 323
anthraxis, 21, 178, 198, 199, 201, 211,
 491
azotobacter, 194
asterosporus, 191
bifementans, 361
bifementans sporogenes, 361, 438
botulinus, 66, 302
bulgaricus, 296
büttchli, 366
camptospora, 183
carotarum, 201
caucasicus, 296
chauvoei, 244
cholerae-asaticae, 409
cholerae-gallinarum, 337
chroococcus, 194
citreus, 185
cochlearius, 101, 314
coli communis, 170
crassus, 429
dentrificans, 288
dentrificans II, 287
dentrificans agilis, 287
egens, 101, 490
enteritidis sporogenes, 179, 370.
erythromyxa, 446
erythrospora, 491
fallax, 101, 521
felis, 274
flexilis, 366
fusiiformis, 314
gigas, 491
hastilis, 360
histolyticus, 528, 529
hortulensis, 452
imetrophus, 464, 530
lactis aerogenes, 169
lacunatus, 291
lebens, 296
lodermos, 324
liquefaciens, 170
marcescens, 464
maximum buccalis, 366
megatherium, 452
multifermentans tenalbus, 101.
mycoides, 202
nitratator, 498

- Bacillus oedematis maligni*, 405, 438
oligocarbophilus, 159
paratuberculosis, 331
paraputrificus, 513
parasporogenes, 370
parvus, 518
perfringens, 361, 438
perlibratus, 290
peromella, 411
prodigiosus, 15, 358, 465
putrificus, 438, 439, 513
pyemiae cuniculi, 439
pyocyaneus, 229, 233
radicicola, 425, 443
radiobacter, 440
ramosus, 191
rhusiopathiae suis, 304
saccharo-butyrificus, 268
sanguineus, 257
schuetzenbachii, 152, 153
solmsii, 290
sphenoides, 100, 297
spirogyra, 366
sporogenes, 370
subtilis, 21, 53, 178, 191, 198, 201, 202, 203, 204, 267, 296, 306, 307, 369, 370, 491
subtilis var *cellaris*, 392
synxanthus, 229
tenuis, 517
terius, 101, 329
tetani, 398
tetanomorphus, 100, 359
tumescens, 531
typhosus, 369
ulna, 21, 198, 389
violaceus, 284, 358
vulgaris, 498
welchii, 370, 371
zeylanica, 528
Bacteriae, 96, 104, 105, 173, 205, 212, 306, 433
Bacteria, 99, 100, 205, 309, 310, 394
Bacteriaceae, 24, 29, 34, 35, 37, 43, 46, 47, 48, 49, 52, 54, 56, 57, 58, 72, 74, 79, 80, 87, 89, 90, 96, 99, 102, 103, 104, 156, 178, 197, 205, 206, 207, 208, 209, 212, 230, 234, 242, 243, 247, 260, 279, 299, 302, 303, 307, 308, 309, 328, 329, 338, 378, 384, 475, 531
Bacteriaceen, 206, 208
Bacteriacees, 207, 208
Bacteriaceae, 33, 207, 208
Bacteriales, 209, 460
Bacteridia, 209
Bacteridiè, 209
Bacteriidae, 92, 93, 209, 211, 310
Bacteridium, 20, 81, 93, 178, 179, 199, 200, 202, 203, 209, 210, 211, 217, 232, 369, 376
anthracis, 211
prodigiosum, 210, 211, 464
violaceum, 221, 436
Bacteriidae, 24, 80, 81, 205, 211, 298, 444, 458
Bacterien, 212
Bactéries, 212
Bacterinae, 81, 82, 212, 420
Bacterina, 18, 212, 216, 523, 527
Bacteriner, 196
Bacteriopsis, 31, 35, 200, 212, 213
aceti, 212
xanthina, 213
merismopedioides, 212
ovata, 213
pasteuriana, 212
rasmusseni, 212, 213
synxantha, 213
ureae, 213
Bacterium, 17, 18, 19, 20, 21, 22, 25, 26, 27, 28, 30, 31, 32, 33, 35, 37, 38, 41, 43, 44, 45, 46, 47, 49, 54, 57, 58, 59, 61, 66, 68, 71, 72, 74, 75, 77, 82, 87, 89, 99, 153, 156, 157, 159, 160, 169, 170, 178, 180, 181, 196, 198, 201, 206, 207, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 234, 242, 277, 298, 306, 308, 310, 360, 369, 372, 382, 386, 410, 419, 429, 434, 435, 452, 459, 498, 518, 519, 523, 527, 528
abortus, 226, 244
aceti, 153, 180, 220, 223, 224, 420, 491, 518

- Bacterium acetigenum*, 501
acidi-lactici, 221, 222
actinocladotrix, 161
aerogenes, 67, 169, 219, 226
aeruginosum, 218, 219, 220, 222, 227, 229
alcaligenes, 226
allantoides, 223
anthracis, 221, 223, 227
articulatum, 214, 360
aurantiacum, 222
balbiani, 223
bovis, 429
brunneum, 220, 229
candidum, 222
casei, 237
catenula, 215, 220, 369
caucasicum, 237
charwoei, 179
chlorinum, 222
cholerae-gallinarum, 223
cholerae suis, 452
coli, 169, 225, 226, 227
coli commune, 222, 439
decalvans, 221
diphtheriae, 67
duplex, 291
dysenteriae, 67
enchelys, 215
erythromyxa, 446
fluorescens non-liquefaciens, 288
foetidum, 221
fusiforme, 219, 360
griseum, 219
lanthinum, 213, 221
incarnatum, 222
influenzae, 67
lactis, 219, 242
lactis acidi, 242, 340
lactis aerogenes, 160, 170, 222
laminariae, 240, 337
leguminosarum, 425
lineola, 220, 288, 360, 532
litoreum, 219
lucens, 220
luteum, 222
malleri, 67
merismopoedioides, 180, 212
morbilli, 222
nitrobacter, 399
oxydans, 498
paratyphi, 66
pasteurianum, 153, 212, 223
pastorianum, 180
pediculatum, 397
pestis, 67
photometricum, 222
pneumoniae, 160
pneumoniae crouposae, 335, 336
punctum, 215, 220
pyocyaneum, 225, 227, 229
rhusiopathiae, 67
roseum, 360
rubescens, 219
septicaemiae, 223
subtile, 221
syncyanum, 218, 219, 220, 222, 229
synxanthum, 218, 220, 222
termo, 213, 214, 215, 216, 217, 218, 219, 220, 222, 223, 224, 225, 227, 228, 229, 498, 524
tortuosum, 222
tremulans, 215, 220
triloculare, 17, 213, 214, 215, 217, 226, 230, 360
tuberculosis, 67, 221
typha, 66, 298
typhosum, 221
ureae, 179, 272
violaceum, 222, 257, 284
viride, 221
xanthinum, 220, 229
xeroseos, 67
zylinum, 154
zopfi, 180, 222, 328, 337, 338, 531
Bacterius, 59, 230
Bacterioidae, 92, 94, 95, 104, 105, 230
Bacterioides, 94, 105, 230, 231, 315
fragilis, 230, 231
fusiformis, 230
Bactrella, 231, 369
bacillus, 231
filum, 231
urdula, 231
Bactridae, 49, 196, 231
Bactridium, 49, 51, 60, 197, 210, 211, 231.

- Bactrillum butyricum*, 232
coli, 232
hypolyticum, 232
megatherium, 232
radicicola, 425
subtilis, 232
typhi, 232
vulgaris, 232
Bactrillei, 49, 196, 232
Bactrillus, 58, 232
Bactrillum, 49, 51, 60, 197, 232
cyanogenum, 232
fluorescens, 232
pseudo-termo, 232
syncyaneum, 232
Bactrineli, 48, 232, 233
Bactrinum, 48, 51, 60, 197, 233
pyocyaneum, 51, 233
Bactrinus, 59, 233
Baculogenae, 34, 39, 40, 197, 223, 233, 234, 515
Baculogene, 234, 337, 459
Bakteriaceen, 234
Bakterium, 179, 181, 234
Balkanella, 93, 96, 234, 298
coagulans, 234
Bangia tenuis, 345
Bacteriacee, 206, 234, 274, 360
Beggiatoa, 18, 20, 22, 25, 26, 27, 28, 29, 31, 32, 34, 35, 36, 37, 40, 43, 44, 45, 46, 48, 50, 52, 53, 55, 59, 61, 64, 70, 74, 84, 234, 235, 236, 237, 258, 310, 312, 345, 351, 352, 353, 357, 381, 408, 461, 479, 515
alba, 234, 236, 310
arachnoideae var *uncinata*, 310
multiseptata, 423
punctata, 18, 234
leptomutiformis, 234
roseo-persicina, 258
Beggiatoaceae, 47, 48, 52, 53, 55, 56, 59, 74, 84, 209, 236, 502, 507, 512
Beggiatoeae, 37, 235, 237, 344
Betabacterium, 237, 238, 419
breve, 237
caucasicum, 237
longum, 237, 238
Betacoccus, 238, 239, 419
arabinosaceus, 239
bovis, 239
mesenteroides, 239
Bibliography, 532
Billetta, 240, 353
Blatta orientalis, 198, 332
Bollingeria, 41, 186, 240, 241, 295, 306
equi, 240, 295
vacchetiae, 240, 295
Borrelia, 64, 240, 241, 479, 481, 485
gallinarum, 240
Botanical code, International, 112
Botanical Society of America Code, 145
Botryococcus, 240, 241
Botryocoques, 241
Botryomyces, 38, 240, 241
ascoformans, 241
Botulobacillus, 71, 241, 439
botulinus, 242
Brachybacterium, 242
apiculatum, 242
Brachybacterien, 61, 242
Brucella, 242, 243, 244
abortus, 244
bronchisepticus, 244
melitensis, 243
Brussel's Code, 112
Bufo calamita, 183
Bursaria, 14, 15
Butyl-Bacillus, 244
Butyribacillus, 71, 244, 245, 247, 268, 438
pasteurianus, 245
Butyribacteriaceae, 69, 71, 244, 245, 247, 416, 419
Butyriclostridium, 245, 268, 439
Byolysis, 245
Calothrix, 23
lutescens, 345
Calymmatobacterium, 245, 335.
granulomatus, 245
Carboxydomonas, 69, 87, 89, 98, 103, 245, 246, 434
oligocarbophala, 246
Carpococcus, 246
pituitoparus, 246

- Carteria*, 246
Carteri, 246
Caryobacterium, 246, 247
 equi, 246
Caseobacterium, 71, 156, 157, 247
 casei, 247
Cassitha, 484
 Casual names, 111
Cellulobacillus, 71, 245, 247.
 methanigenes, 247
Cellulomonadeae, 104, 105, 247
Cellulomonas, 105, 247
 brazoteus, 248
Cenomesia, 41, 186, 248, 306
 albida, 248
 hlacina, 248
Cephalotrichinae, 68, 69, 152, 157, 169
 247, 248, 287, 342, 358, 359, 371,
 409, 419, 441, 466, 473, 501, 515
Chaetophora endivaeifolia, 297
Chaetophorae, 161
Chamaesiphon, 23, 37
Chara, 461
Chatinella, 249
 scissipara 249.
Chronyphe, 249
 carteri, 249
Chlamydatomus, 25, 41, 186 249, 250,
 316, 331, 458
 bergeli, 249, 250
 cellaris, 250
Chlamydobacteriaceae, 47, 48, 52, 53,
 54, 56, 57, 59 74, 83, 91, 106, 209,
 250, 263, 266, 283, 289, 290, 308,
 309, 351, 384, 424
Chlamydobacteriales, 79, 87, 88, 100,
 102, 106, 251, 461
Chlamydobacterinae, 251
Chlamydothrix, 55, 60, 74, 251, 252, 319,
 350
 epiphytica, 252
 ferruginea, 252, 318, 321, 486
 frutans, 252
 hyalina, 252
 ochracea, 72, 251, 252
 sideropous, 71, 252
Chlorobacterium, 252
 lactis, 252, 253.
Chlorobacteriaceae, 253
Chlorobakteriaceae, 253, 254, 461
Chlorobium, 253.
 limicola, 253
Chlorochromatium, 253, 254, 286, 417.
 aggregatum, 253, 254, 286
Chloronum, 254
 mirabile 254
Chlorophyllophyceae, 24
Chondromyces, 19, 62, 74, 86, 107, 254,
 255, 287, 393, 394, 395, 430, 431, 489
Chondromyces crocatus, 254, 255
Chorynebacterium, 255
Chromatraceae, 55, 63, 255, 256
Chromatraceae, 84, 85, 256, 442, 445, 450.
Chromatrideae, 84, 176, 256, 257, 445,
 503, 506
Chromatium, 33, 36, 56, 70, 75, 85,
 200, 207, 220, 255, 256, 257, 308,
 311, 410, 441, 442, 448
Chromatium minus, 256
 minutissimum, 256
 okenii, 256, 257
 vinosum, 256, 257
 violascens, 256
 weissii, 256
Chromatrideae, 257
Chromobacillus, 38, 200, 257, 371
Chromobacteriae, 96, 104, 257, 258, 313,
 436, 465
Chromobacteriaceae, 257
Chromobacterium, 82, 96, 99, 104, 212,
 257, 284
 violaceum, 221, 258, 436
Chromobeggiatoa, 37, 235, 258
Chromococcus, 258, 285
 violaceus, 258, 436
Chroococcaceae, 24
Chroococcus, 22, 37
Chthonoblastus, 23
Ciliata, 18
Cladochytrium, 258
 tuberculum, 258
Cladophyllum, 18, 181, 182, 258, 259
 comatum, 182, 259
 ramosissimum, 259
Cladothricaceae, 37
Cladothriceae, 26, 259, 260

- Cladothrichaceae*, 259, 260, 288
Cladothrichacei, 259, 260
Cladothricheae, 260
Cladothricheen, 259, 260, 261
Cladothrix, 23, 26, 27, 29, 30, 31, 32, 34, 35, 37, 39, 43, 44, 45, 46, 48, 50, 52, 53, 58, 59, 60, 67, 68, 70, 79, 250, 252, 259, 260, 261, 262, 263, 264, 265, 266, 335, 353, 402, 469, 496, 497, 515, 516
dichotoma, 23, 65, 72, 260, 262, 263, 264, 468
dichotoma var *leptochaetoformis*, 261
placodes, 263
Cladotricaceae, 39, 264, 515
Cladotricaceae, 264, 402
Cladotricheae, 264
Cladotricheen, 264
Clathrococcus, 264
roseo-persicinus, 264
Clathrocystis, 22, 32, 37, 264, 265, 276, 277, 342
aeruginosa, 264, 277
roseo-persicina, 19, 219, 264
Clerodendron, 484
Clitridium, 265
Cloaca, 93, 265, 432
cloacae, 265
Clonothrix, 61, 74, 83, 106, 251, 265, 266
fusca, 65, 72, 265, 266
gracillima, 266
Clostridiaceae, 100, 266, 302, 439, 463
Clostridiaceae, 51, 60, 196, 266
Clostridioidae, 100, 179, 266, 267, 268, 297, 313, 329, 359, 371, 382, 405, 406, 451, 489, 521, 529
Clostridium, 19, 27, 28, 31, 33, 35, 40, 44, 46, 49, 51, 53, 59, 60, 76, 80, 87, 90, 99, 100, 106, 177, 178, 179, 197, 199, 202, 206, 207, 234, 242, 245, 266, 267, 306, 308, 324, 411, 415, 419, 428, 438, 439, 520
amylobacter, 268, 269
butyricum, 100, 267, 268, 269, 520
feseri, 267
pasteurianum, 244
polymyxa, 267
Clostrillum, 49, 232, 269
Clostrinum, 49, 233, 269
Clostridium, 269
Coccaceae, 46, 47, 49, 51, 52, 54, 56, 58, 59, 60, 65, 66, 72, 73, 74, 78, 79, 87, 88, 90, 91, 96, 97, 103, 174, 209, 269, 270, 171, 274, 292, 308, 309, 326, 331, 337, 368, 373, 384, 395, 396, 411, 466, 467, 492
Coccaceae, 269
Coccacei, 32, 271, 273
Coccobacillus, 38, 200, 271, 272, 337
acridiorum, 272
avicidus, 38, 272
conjunctivae, 272
fusiformis, 272
perfoetans, 272
zymogenes, 271, 272
Cocco-Bacillus, 272
Cocco-Bacillus anaerobius perfoetans, 272
Coccobacteria, 21, 32, 270, 272, 273, 337, 361
septica, 21, 187, 272, 274, 275, 276, 291, 320, 328, 347, 361, 366, 372, 373, 374, 382, 420, 457, 466, 491, 492, 495
Coccobacteriaceae, 273, 471, 476
Coccobacterium, 273, 337
fehls, 274
mucosum anaerobium, 274
prodigiosum, 274
Cocco Bacterium, 274
Coccochloris, 22
Coccogenae, 35, 39, 41, 186, 248, 250, 274, 373
Coccogene, 274, 458, 459, 492, 515
Coccoglia, 274, 530
Coccomonas, 274, 275, 276
Coccus, 275, 276
Coccothrichaceae, 31, 275
Coccothrix, 31, 200, 275, 337, 386, 387, 389, 463
leprae, 275
tuberculosis, 275
Coccotrichum, 275, 387
Coccus, 275, 276, 337, 419, 493
albus non-liquefaciens, 276
aquatilis, 276
candicans, 276

- Coccus cumulus minor*, 276
Coccus salinarum septicus, 276
 Code, Lehmann and Neumann, 109
 Codes of nomenclature, 108
 Code of Nomenclature, Zoological, 136
Coelosphaerium, 22, 248
Cohnia, 27, 37, 265, 276, 277, 341, 376
 roseo-persicina, 276
Cohnella, 276
Cohnastreptothrix, 73, 95, 164, 167, 276,
 404, 405
 foersteri, 164
 israeli, 164, 277
Coleothrix, 277
 methystes, 277
Colibacterium, 99, 170, 277
 Colon bacillus group, 66
Commelinaceae, 484
 Committee on Classification, 119
Conserva moniliformis, 316
 nummuloides, 316
 ochracea, 317
Conidiopora, 94
Conidiothrix, 278
Coniothecium, 278
 syphiliticum, 278
Coniomyces, 163
Cornelia, 40, 197, 278, 306, 310, 428
 alvei, 278, 279, 310
 pasteuri, 278
Cornelia sphinx, 428
 tremula, 428
Corynebacterium, 50, 52, 53, 57, 82, 87,
 89, 94, 98, 106, 161, 168, 212, 255,
 279, 280, 333, 378, 384, 385, 386, 387
 diphtheriae, 279, 378
Corynemonas, 69, 169, 280
 diphtheriae, 280
Corynethrix, 58, 280
Corynoladus, 18, 280
 radiatus, 280
Crenothrichaceae, 280, 288, 390, 391
Crenothrix, 20, 23, 25, 27, 29, 30, 31,
 32, 34, 35, 37, 39, 43, 44, 45, 46,
 48, 50, 52, 53, 55, 58, 59, 60, 70, 74,
 83, 106, 250, 251, 265, 281, 282, 283,
 310, 312, 340, 345, 351, 352, 234, 424,
 515,
 foetida, 282, 423
 kuhmana, 282, 283, 310, 423
 marina, 282, 423
 mucor, 423
 multiseptata, 423
 polyspora, 20, 65, 72, 281, 282, 283, 363
Crenothrichaceae, 283
Crenothricaceae, 39, 283, 515
Cristispira, 72, 78, 86, 88, 107, 283, 453,
 479, 481, 482, 484
 balbiana, 78, 284
 interrogationis, 283
 veneris, 283, 284
Cromobacterium, 28, 257, 284
 violaceum, 28, 284
Cromococcus, 258, 285
 violaceus, 285
Crypta, 245, 285
 syphilitica, 285
Cryptococcus, 20, 286, 518
 xanthogenicus, 286
Cryptodesma, 18, 286
 tenuis, 286
Cyanophyceae, 24
Cychnidium, 15
Cylindrogloea, 254, 286
 bachifera, 286
Cystobacter, 34, 45, 207, 255, 286, 287,
 308, 430
 erectus, 255, 286
 fuscus, 286
Cystococcaceae, 189, 287, 331, 356
Cytobacter, 287

Daphnia pulex, 415
Deazotomitrarazobacterium, 287
 thermophilum, 287
Deazotomitrarazobacterium, 287
Deazotomitrarazobacterium, 287
Denitrobacillus, 287
Denitrobacterium, 71, 156, 157, 287
 agile, 287
 thermophilum, 287
Denitromonas, 70, 288, 314, 358, 359,
 466
 fluorescens, 288
Desmobacteria, 21, 35, 37, 260, 280, 288,
 344

- Desmobacteriaceae*, 74, 288, 289
Desmobacteries, 200, 289
Desmobacterinae, 74, 288, 289, 326
Desmobacterium, 289
Detoniella, 39, 289, 332, 338, 348
ochracea, 289
Dialister, 106, 289.
pneumosintes, 289
Diastizae, 289, 411
Dicoccia, 41, 289, 290, 310, 337
glossophila, 290
Didymohelz, 19, 83, 106, 251 290, 317,
 319, 485
ferruginea, 290, 321
ochracea, 319
Diffusionsbacillus, 290
Diphtheria group, 67
Diplectridium, 49, 290
Diplobacillus, 290, 291
brevis endocarditidis, 290
liquefaciens, 291
Morax, 291
Morax-Azenfeld, 291
Diplobacteria, 291.
Diplobacterium, 291
Diplococcus, 291
Diplococcus, 31, 44, 45, 65, 66, 80, 92
 96, 104, 291, 293, 336, 376 396, 411,
 429, 434, 492
albicans tardissimus, 292
coryzae, 292
lanceolatus capsulatus, 66
luteus, 292
pneumoniae, 31, 292, 293, 336, 434
roseus 292
subflavus, 292
Diplokokken, 293
Diplokokkus, 43, 293
Diplopneumococcus, 293
Diplostreptococcus, 293, 294
intestinalis, 294
pleuropneumoniae, 293, 294
Diplovibrio, 294
Discomaces, 294
Discomyces, 24, 158, 163, 164, 165, 166,
 240, 294, 402
boris, 24, 295
equi, 163, 240, 241, 294, 295
pleuriticus, 163, 294
pleuriticus canis famaharis, 240, 294.
somaliensis, 333
Discomycetaceae, 295
Discomycetes, 164, 295
Dispora, 28, 289, 296, 297, 409
caucasica, 28, 296
Douglasillus, 100, 266, 297
sphenoides, 100, 297
Drepanospira, 297
mulleri, 297
Drepanospira, 297
Dysenteroides, 93, 96, 297, 298
metadysentericus, 297

Ebertheae, 92, 95, 173, 234, 297, 298, 301,
 306, 343, 452, 465, 529
Eberthella, 82, 105, 226, 298
typhi, 298
Eberthus, 93, 96, 298
typhosus, 298
Eccrina, 18, 298
longa, 298, 299
moniliforma, 298
Enterbacterium, 299
Enterococcus, 299
Elephantopus, 477
Encapsulateae, 92, 95, 104, 105, 299, 200
Encapsulatus, 92, 95, 105, 299, 331, 336.
pneumoniae, 299
Enchelis, 14, 15
Endobacterium, 300
Endostreptococcus, 300
Endostreptokokkus, 300
Enterobrus, 300
attenuatus, 300
elegans, 300
spiralis, 300
Enterobryus, 18, 298, 301
elegans, 182
Enterococcus, 301
saccharomyces, 301
Enterocoque, 301
Enteroides, 93, 298, 301
entericus, 301
Entophyta, 301
Erebonema, 301, 354
divaricatum, 301

- Erebonema hercynicum*, 301, 302, 355
Ermengillus, 302, 439
 botulinus, 302
Eruvinae, 96, 104, 302, 303
Eruvina, 87, 90, 99, 105, 303
 amylovora, 303
Eruvinae, 105, 302, 303, 424
Eruvinae, 302, 303
Erysipelococcus, 303
Erysipelothrix, 71, 82, 96, 97, 106, 212, 303, 304
 erysipeloides, 303
 murisepticus, 303
 porci, 303, 304
 rhusiopathiae, 304
Erythrobacillus, 62, 96, 99, 257, 304, 465
 prodigiosus, 304
 pyosepticus, 304
Erythroconis, 305, 343
 littoralis, 305
Escherichia, 93, 105, 298, 306
 coli, 306
Eusacococceae, 41, 186, 248, 306
Eubacilleae, 40, 197, 306
Eubacillus, 38, 44, 75, 200, 306, 307
 multisporus, 307
Eu-Bacillus, 81, 203, 307
Eubacteria, 37, 56, 61, 74, 75, 206, 297
 307, 308, 371, 390, 391
Eubacteriaceae, 308
Eubacteriales, 78, 79, 87, 88, 89, 91, 100, 102, 103, 196, 208, 251, 266, 308, 309, 334, 385, 399, 400, 319, 434, 459, 460, 461, 470, 471
Eubacteriae, 24, 100, 212, 309, 310, 374
Eu-Bacterium, 82, 226
Eubeggiatoa, 37, 235, 310
Eucoccaceae, 310, 375, 387
Eucoccus, 310
Eu-Cornelia, 310
Eucornelia, 278, 310
Eucrenothrix, 282, 310, 423
Euglenaceae, 24
Euklebsielleae, 41, 289, 310, 337
Eumantegazzaea, 310, 360
Eumacrospira, 75, 310
Eumonas, 311
Eumyriophyllum, 485
Eu-Pacina, 311, 409
Eupacina, 311, 409
Euplanococcus, 311
Euplanosarcina, 311
Eupseudomonas, 311
Eusarcina, 311
Euspirillum, 75, 312
Eu-Spirillum, 312
Euspirosoma, 54, 312, 385
Euvibrioniae, 25, 235, 281, 312, 477, 482
Fenobacter, 312
Ferribacterium, 312
 duplex, 312
Fizococcus, 76, 313
Fizosarcina, 76, 313
Fizostreptococcus, 76, 313
Flavobacterium, 104, 185, 313
 aquatilis, 313
Flemingillus, 101, 266, 313
 cochlearius, 101, 314
Flexner group, 93
Flexnerella, 93, 314
Fluorescent group, 66
Fluormonas, 99, 314, 358, 434
Fungi-Imperfecti, 94
Fusiforminae, 81, 212, 314
Fusiformis, 72, 81, 87, 89, 97, 106, 168, 231, 314, 315, 360, 385
 dentium, 314
 muris, 314
 termitidis, 314, 315
Gaffkya, 31, 35, 41, 186, 315, 316
 tetragena, 315, 365
Gaffkya, 41, 186, 250, 315
Gaillonella, 17, 316, 317, 318, 319, 320
 ferruginea, 17, 251, 316, 317
 lineata, 316
Galactobacterium, 319
Galactococcus, 320
 albus, 320
 fulvus, 320
 versicolor, 320
Galeria melanella, 363
Galkhonella, 17, 74, 252, 316, 317, 318, 320, 350, 485, 486
 ferruginea, 19, 64, 72, 283, 290, 317, 318, 321, 332, 349, 363, 485
 ferruginea var *lata*, 319, 348

- Gammarus locusta*, 423
 Genera conservanda, 118, 119
Glaeogenes, 320
 Glanders group, 67
Glaucothrix, 37.
Gleobacter, 320
Gliabacteria, 320
Gliacoccus, 272, 274, 320
Gliacoccus, 320, 530
 aceti, 320
Glia-kokkus, 320
Glschrobacterium, 320, 321
Glschrobacterium, 320
Gloeocapsa, 22, 37
Gloegenae, 265, 454
Gloeogenes, 22, 188, 355
Gloeosphaera, 318, 321
Gloeosphaera ferruginea, 318, 321
Gloeotheca, 22, 37
Gloeothila, 290, 317, 321
Gloeothila ferruginea, 217, 321
 ochracea, 321
Glucobacter, 321
Glycobacter, 321, 322
 peptolyticus, 322
 proteolyticus, 321
Gomphosphaeria, 22
Gonium, 15, 322
Gonium hyalinum, 322
Gonococcus, 292, 322, 376
 gonorrhoeae, 322
 neisseri, 322
Gracilondeae, 92, 93, 95, 322, 323
Gracilondes, 93, 322, 323
 albafaciens, 323
Granulobacillus, 323
 saccharobutyricus immobilis lique-
 faciens, 323
 saccharobutyricus mobilis non-lique-
 faciens, 268
Granulobacter, 50, 59, 177, 178, 323, 324,
 415
 butylicum, 323
Granulobacter immobilis, 411
 lactobutyricum, 323
 pectinovorum, 416
 polymyza, 323
 saccharobutyricum, 323
Granulobakter, 324
 butylicum, 324
 pectinovorum, 324
 saccharo-butyricum, 268
 urocephalum, 324
Grippestreptokokkus, 324
Gummibacillus, 324
Gymnobacteria, 324, 382, 419
Gymnodes, 16, 527
Gyrococcus, 324
 flaccidifex, 324
Haematococcus, 325, 521
 navalis, 325
Hamatokokkus, 325
Haemophyleae, 92, 325
Haemophilus, 325, 329
 influenzae, 325
Halbacterium, 47, 325
 aurantiacum, 325
 liquefaciens, 325
 pellucidum, 325, 326
 polymorphum, 325
 purpureum, 325
 roseum, 325
 rubrofusum, 325
Halorrhagidaceae, 485
Haplobacteriaceae, 326, 327
Haplobacterinae, 51, 59, 73, 326
Haplobakterien, 327
Hay bacillus group, 66
Helicobacterium, 327, 328
 aerogenes, 327
 klebsii, 327
Helicomonaden, 327, 328
Helicomonas, 327
 sphyiliticum, 327
Helikobakterium, 328
 zopfii, 328
Helicomonas, 328
Helminthica, 14
Helobacteria, 328
Hemisporales, 94
Hemophilaeae, 328
Hemophyleae, 95, 96, 104, 106, 289, 328,
 329
Hemophilinae, 81, 192, 212, 328
Hemophilus, 81, 87, 89, 99, 106, 325,
 328, 329
 influenzae, 329

SUBJECT INDEX

- Henricillus*, 101, 266, **329**
tertius, 101
Hiblerillus, 101, 267, **329**
megalosporus, 330
rectus, 330
septimus, 330
sextus, 101, 330
Hillhousia, 71, 84, 154, **330**, 331
mirabilis, 330
Homococcaceae, 51, 60, **331**
Homonyms, 124, 127, 148
Hormosiphon, 23
Hyalococcus, 33, 34, 37, 38, 45, 287 **331**,
336, 376
bergii, 250, 331, 428
cellaris, 250
pneumoniae, 331, 336
Hydrogenomonas, 69, 87, 89, 98, 103, **331**,
332, 434
flava, 331, 332
pantotropha, 331, 332
vitrea, 331, 332
Hydrotremellinae, 162
Hygrococcus, 20, 259, **332**, 343, 369, 459
intestinalis, 183, 332
ochracea, 316, 345
olivacea, 345
Hyphales, 94, 377, 378
Hyphothrix, 23, 37, **332**, 345
callitrichae, 462
Hyphomycetes, 50, 94, 163, 166, **333**, 385
Hyphomicrobium, **333**, 401
vulgare, 333
Hyphomycetes, 489, 495, 516
Hypnococcus, 61, **333**
Hyponym, 148

Indrella, **333**
somaliensis, 333
Indrellopsis, **333**
somaliensis, 333
Indolococcus, **333**, 334, 418
Influenza group, 67
Infusoria, 14, 16
Iodococcus, **334**
Iron bacteria, 64

Jodococcus, **334**

magnus, 334, 335.
parvus, 334
vaginatus, 334
Julus, 181, 258, 332
marginatus, 182, 259, 300

Kalymmabacterium, **335**
Karphococcus, 335
Karphokokkus, **335**
Keratophyton, **335**
Kladotrix, **335**
Klebsiella, 31, 35, 41, 77, 289, 300, 310,
320, **335**, 337, 434
aerobia, 335
cellaris, 392
crouposa, 335, 336
edingtoni, 184
enteritidis, 336
friedlanderi, 336
leydeni, 434
pneumoniae, 335, 336
rhinoscleromatis, 335, 336
sahvaris, 335, 336
Klebsiellae 35, 41 233, 289, 299, 310,
336, 337
Klebsiellae, 234, **337**
Kohhaccean, **337**
Kokken, **337**
Kokkobacillus. **337**
aucidus, 337
zymogenus, 337
Kokkobacteria, 337
Kokkobacterium, 337
lyssae, 337
Kokkothrix, 337
Kokkus, 43, 276, 337
Kolpoda, 15
Kurthia, 31, 34, 39, 240, **337**, 338, 353,
532
zopfii, 337, 338
Kurthiae, 39, 289, **338**, 515

Lacrimatoria, 16, 527
Lactobacillaceae, 87, 89, 96, **338**, 339
Lactobacillae, **338**, 340, 501
Lactobacilleae, 96, 104, 105, **338**
Lactobacillus, 82, 87, 90, 99, 105, 153,
212, 242, 247, **338**, 339, 340, 452

- Lactobacillus caucasicus*, 339
delbrückii, 339
fermentatum, 237
fermentum, 339
longus, 339
Lactobacter, 340
Lactobacteriaceae, 340
Lactococcus, 339, 340, 341
agglutinans, 341, 356
hollandiae, 341
lactis, 340, 341
Lactosarcina, 341
Lampozonidea, 18, 341
Lamprocystaceae, 55, 63, 341
Lamprocystae, 84, 85, 256, 341, 342
Lamprocystis, 27, 33, 34, 36, 37, 41, 55,
 70, 74, 85, 186, 248, 264, 265, 277,
 306, 311, 341, 449, 504, 508
gelatinosa, 509
roseo-persicina, 342, 377, 428, 433, 510
violacea, 510
Lampropedia, 33, 34, 37, 42, 85, 305,
 342, 376, 458, 506
hyalina, 322, 343
reitenbachii, 364
violacea, 436
Lampropediaceae, 506
Lankoides, 93, 96, 298, 343
pyogenes, 343
Lauraceae, 484
Leptomitus, 259, 343
divergens, 344
Leptonema, 344, 352
nivea, 344
Leptosira, 107, 344
icterohaemorrhagiae, 344
icteroides, 344
Leptothricaceae, 37, 237, 288
Leptothrichaceae, 344
Leptothricaceae, 37
Leptothricheae, 20, 344, 348, 518
Leptothricheen, 345
Leptothrix, 18, 20, 22, 26, 27, 28, 29, 30,
 31, 32, 34, 35, 37, 43, 45, 46, 50,
 53, 57, 67, 68, 70, 83, 106, 206, 221,
 251, 252, 260, 261, 278, 289, 317,
 319, 345, 348, 347, 348, 349, 350,
 351, 352, 389, 405, 440, 496, 497, 515,
 516
aeruginea, 345
buccalis, 18, 346, 347, 348, 350, 352
cellaris, 348
fontana, 345
insectorum, 346
kuhmana, 283
lutescens, 345
maxima, 351
ochracea, 18, 64, 251, 278, 283, 318,
 345, 346, 348, 349, 350, 351, 362.
parasitica, 347, 348
thurethana, 348, 408
Leptotrichaceae, 344, 351, 391
Leptotrichacei, 34, 351
Leptotrichaeae, 39, 345, 351, 515
Leptotrichheen, 351
Leptotrichia, 25, 34, 39, 83, 87, 89, 97,
 106, 168, 213, 312, 347, 348, 351, 352,
 353, 357, 385, 440
amphibola, 353
buccalis, 347, 353, 353, 440
mucor, 357
nivea, 344
ochracea, 348
thurethana, 408
rigidula, 344
Leptotrichieae, 34, 353
Leptotrichiae, 240, 337, 352, 353
Leptotrichiella, 353
Leucocystis, 33, 34, 37, 38, 41, 186, 287,
 302, 306, 353, 354, 355, 376
cellaris, 353, 391
cellaris var *cavernarum*, 355
pneumoniae, 354
Leuconostoc, 24, 28, 29, 31, 33, 35, 37,
 38, 42, 44, 45, 51, 74, 80, 96, 97,
 104, 181, 188, 190, 206, 240, 269,
 287, 341, 355, 356, 357, 395, 467, 494,
 492
agglutinans, 356
lagerheimii, 154, 356
lagerheimii var *subterraneum*, 356
mesenteroides, 24, 355, 357
Leucothrix, 353, 357
Liliaceae, 276
Lamnochlide, 23
Laneola, 16, 357
Lapobacter, 63, 357
Liquidobacterium, 71, 173, 357, 358, 432

- Liquidobacterium miquelii*, 520
prodigiosum, 358, 519
vulgare, 358
Liquidococcus, 70, 358, 359
Liquidomonas, 70, 288, 314, 358, 359
fluorescens, 358
Liquidovibrio, 70, 359
comma, 359
Loefflerella, 420
Lophotricha, 359
Lummbacteriaceae, 68, 70, 157, 248,
 288, 358, 359, 466
Lyngbya, 37
ochracea, 317, 345, 349
Lyngbyeae, 234
Lyngbyoideae, 234

Macintoshillus, 100, 266, 359
tetanomorphus, 100, 359
Macrococcus, 359
Makrokokkus, 359
Malleomyces, 360
equestris, 360
Mantegazzaea, 25, 40, 78, 197, 306, 310,
 312, 315, 360, 441, 442
articulata, 360
cienkowski, 360
hashis, 360
Martellillus, 361, 370, 439, 440, 441,
 529
bifermentans, 361
Mastichomonas, 361
Mastigophora, 471
Megabacteria, 361
Megabacterien, 361.
Megabacterium, 361
Megacoccus, 361
Megacoccus, 359, 362
Megalohrix, 362
discophora, 362
Melanella, 16, 363, 527
atoma, 363
erythraea, 363
flexuosa, 363,
monadina, 363
spirillum, 363
Melococcus, 363
ostrjanni, 363

Meloseria, 316, 363
ochracea, 290, 363
Meloseria, 317, 318, 319, 363
ochracea, 317, 363
Meningococcus, 292, 363, 364
intracellularis, 364
Meningokokkus, 364
intracellularis, 364
Merismopedia, 22, 30, 43, 44, 45, 269,
 305, 343, 364, 365, 375, 376, 454, 455,
 511
gonorrhoeae, 364
littoralis, 364
Merismopedium, 37, 365
reitenbachii, 364, 365
Merista, 30, 43, 77, 181, 343, 364, 365,
 376, 417, 426
laevigata, 365
tetragena, 365
ureae, 365
Merizomyria, 23, 365
aponina var *fasciculata*, 366
Merulus lacrymans, 433
Mesobacteria, 366
Mesobacterien, 366
Mesobacterium, 366
Mesococcus, 366
Mesococcus, 366
Metabacterium, 77, 81, 197, 366, 367, 368.
polyspora, 366, 367
Metacoccaceae, 61, 65, 270, 368
Metacoccus, 368
Metallacter, 18, 25, 200, 212, 216, 231,
 312, 368, 369, 370, 523
amylobacter, 369
anthracis, 369
bacillus, 368, 369
ileotyphi, 369
tremulus, 369
ulna, 369
Metchnikovillus, 370
sporogenes, 370, 439
Methanomonas, 69, 87, 89, 98, 103, 370,
 371, 434
methanica, 371
Metonym, 148
Meyerillus, 102, 267, 371
sadowa, 102, 371

- Microloa teres*, 216
Microbacillus, 371
 seborrhoeae, 371
Microbacteria, 21, 35, 38, 200, 207, 217, 223, 371
Microbacterien, 371, 372, 381
Microbacteries, 372
Microbacterium, 99, 372, 419
 flavum, 372
 lacticum, 372, 373
 mesentericum, 372
Micrococca, 211
Micrococceae, 42, 79, 80, 87, 91, 92, 103, 104, 271, 274, 368, 373, 377, 398, 447, 457
Micrococceae, 274, 396, 488
Micrococcoglia, 373
Micrococcus, 373, 374.
Micrococcus, 21, 22, 24, 26, 27, 28, 29, 30, 32, 33, 35, 37, 42, 43, 44, 45, 46, 47, 49, 51, 53, 54, 56, 58, 59, 60, 65, 66, 67, 71, 72, 74, 76, 77, 80, 87, 90, 92, 104, 156, 157, 171, 172, 174, 188, 189, 190, 191, 211, 246, 269, 270, 276, 310, 313, 339, 355, 358, 364, 365, 368, 373, 374, 375, 376, 377, 381, 416, 417, 422, 425, 426, 434, 447, 466, 467, 488, 493, 494, 526
 aceti, 320, 420, 440
 ascoformans, 240, 241
 aureus, 172
 bengelii, 428
 billrothii, 190
 bombycis, 410
 botryogenes, 240, 241
 butyricus, 362
 candicans, 375, 376
 casei liquefaciens, 499
 catarrhalis, 66
 cellaris, 355
 cholerae-gallinarum, 414
 citrinus, 376
 erythromyxa, 446
 fickii, 276
 flavus, 376
 gallicidus, 414
 gonorrhoeae, 66, 322
 metrophus, 530
 intracellularis meningitidis, 66
 lactis, 374
 luteus, 373, 374, 376, 377
 mehliensis, 243
 mirificus, 410
 nitrosus, 401
 ovalis, 301
 phosphorescens, 422
 prodigosus, 373, 377, 410, 464
 pyogenes, 171, 172
 rhodochrous, 446
 scarlatinus, 419
 septicus, 374
 tetragenus, 66, 315, 365, 416
 tetragenus mobilis ventriculi, 426
 ureae, 213, 272, 365, 521
 violaceus, 436
Microhaloa, 376, 377
 aeruginosa, 377
 botryoides, 377
 rupestris, 377
Micromyces, 44, 163, 377, 403
 hoffmanni, 377
 zygogoni, 377
Microphyta, 377
Microsiphonales, 94, 377, 278, 385, 405
Microsiphonés, 279, 378, 404, 415, 460, 462
Microsphaera, 21, 374, 378
 vaccinae, 21, 378
Microspira, 33, 34, 48, 54, 57, 59, 61, 74, 75, 207, 297, 308, 310, 359, 378, 379, 409, 410, 422, 435, 437, 449, 470, 473, 524, 526
 buccalis, 379
 comma, 379
 finkleri, 379
Microspironema, 62, 379, 380, 478, 483, 513
 pallidum, 380
Microspora, 380
 tyrosinatica, 380
Microsporon, 20, 68, 374, 380
 septicum, 20, 380
Microsporum, 380
Microzoa, 380
Microzoma, 374, 380, 381
 bombycis, 380

- Microzymba cretae*, 380
 Microzyme, 380
 Mikrobakterien, 381
Mikrococcus, 44, 381
Mikrokokkus, 43, 181, 381
Mikromyces, 381
Mikrospironeuma, 381
 Mikrozymas, 381
Modderula, 155, 381
 hartwegi, 155, 381
Monadina, 17
Monas, 14, 15, 17, 32, 311, 361, 376, 382
 capusculum, 17, 311
 lens, 15, 361
 mica, 382
 mülleri, 155
 oteni, 219, 256, 361, 442
 prodigiosa, 382, 464, 530
 pulmonale, 382
 punctum, 15, 361
 terno, 15, 213, 214, 215, 218, 361
Monobacillus, 382
 Monobacteria, 382
 Monococcus, 382
Monococcus, 382
Monotricha, 382
Monovibrio, 382
 Morax-Axenfeld bacillus, 291
Mortierella, 249
Mucor, 249
Multifermentans, 101, 267, 382
 tenalbus, 101, 382
Mycacanthococcus, 42, 383
 cellaris, 383
Mycobacillus, 383, 384
 synovialis, 383
Mycobacteriaceae, 52, 53, 56, 57, 73, 87,
 88, 89, 91, 94, 96, 106, 167, 168, 279,
 309, 326, 378, 384, 385, 387
Mycobacterineae, 80, 83, 385
Mycobacterium, 50, 52, 53, 57, 58, 83,
 87, 89, 94, 97, 106, 159, 161, 168,
 275, 333, 384, 385, 386, 387, 389, 462
 leprae, 387
 rhusiopathiae, 304
 tuberculosis, 385, 387
Mycococcaceae, 38, 270, 287, 310, 387,
 391, 467
Mycoderma, 16, 81, 87, 89, 96, 153, 154,
 259, 387, 388, 389, 444, 519
 aceti, 212, 388, 518
 cerevisiae, 388
 mesentericum, 16, 387
 ollare, 387, 388
 pasteurianum, 388
Mycomonas, 69, 169, 389
Myconostoc, 23, 26, 27, 30, 32, 33, 34,
 37, 38, 41, 45, 54, 207, 308, 312,
 337, 389, 390, 485, 486
 gregarium, 23, 389, 485
Myconostocaceae, 38, 308, 389, 390, 391
Myconostocaceae, 26, 34, 41, 337, 390, 527
Mycophyceae, 283, 391
Mycotetraedron, 42, 391
 cellare, 391
Mycothamnion, 391
 fodinatum, 391
Mycothece, 37, 38, 287, 391
 cellaris, 391, 392
Mycothrax, 392
Myriophyllum, 485
Myrsine urillei, 365
Myxobacillus, 64, 392
 betae, 392
Myxobacter, 46, 392, 393, 430
 aureus, 392, 430
Myxobacteria, 61, 393, 431
Myxobacteriaceae, 15, 75, 86, 107, 255,
 326, 392, 393, 394, 395, 430, 489
Myxobacterales, 79, 87, 88, 91, 102, 107,
 309, 393, 394, 459, 461
Myxobacterineae, 100, 310, 394
Myxobacterium, 394
Myxobacterales, 394, 460
Myxobacillus, 392, 394
 variabilis, 255, 394
Myxobotryaceae, 394
Myzococcus, 46, 72, 74, 86, 107, 255,
 393, 395, 430
 currosus, 395
 clavatus, 395
 corallinoides, 395
 cruentus, 395
 disciformis, 395
 digitatus, 395

- Myxococcus macrosporus*, 395
pyriformis, 395
rubescens, 395
striptatus, 395
virescens, 395
Myxokokkus, 64, 395
betae, 395
Myxomycetales, 102
Myxomycetes, 425, 443

Neisseriae, 103, 395, 396
Neisseria, 42, 77, 80, 87, 90, 91, 96, 97,
103, 292, 293, 322, 364, 373, 395, 396,
397, 492
gonorrhoeae, 396, 397
Neisseriae, 396
Nematogenae, 19, 22, 188, 234, 260, 346,
355, 389, 397, 493
Nematogenes, 198, 347, 397, 472, 477,
496, 524
Nevskia, 46, 397
ramosa, 397
Newsia, 397
Nicolaerillus, 397, 398, 427, 439
tetani, 398
Nigrococcus, 92, 373, 398
nigrescens, 398
Nitribacillus, 398
Nitrobacter, 46, 83, 87, 89, 98, 103, 398,
399, 401
ohgotrophum, 399
polytrophum, 399
winnogradskyi, 399
Nitrobacteriae, 92, 95, 96, 103, 154, 246,
332, 399, 400, 402, 501
Nitrobacteriaceae, 79, 83, 87, 88, 89, 96,
98, 103, 195, 308, 399, 400
Nitrobacteriae, 399, 400
Nitrobacterium, 400
nitrobacter, 400
Nitromicrobium, 400, 401
germinans, 400
Nitromikrobium, 401
Nitromonas, 69, 401, 402, 434
Nitrosobacterium, 401
novae formae, 401
Nitrosobakterium, 401
Nitrosococcus, 46, 83, 399, 401, 402
americanus, 401
nitrosus, 401
Nitrosomonas, 46, 69, 83, 87, 89, 98,
103, 399, 400, 401, 402, 434, 498
europoea, 402, 426
javanensis, 402
Nocardia, 39, 67, 83, 87, 89, 94, 96, 163,
165, 166, 167, 246, 264, 277, 294, 295,
333, 378, 402, 403, 404, 405, 415,
496, 516
actinomycetes, 161, 163, 402
bovis, 404
farcinica, 402
foersteri, 378
somaliensis, 333
Nocardiaceae, 94, 277, 378, 404, 405, 516
Nodofolium, 72, 405
ferrugineum, 319, 405
Nomenclature, 108
Nomina hybrida, 124
Nosema, 19, 405, 519
bombycis, 405, 410, 459
Nostoc, 23, 37, 356, 490
Nostocaceae, 24, 29, 355
Novillus, 102, 267, 405
maligni, 102, 405

Octopsis, 405
cholerae-gallinarum, 405
equorum, 405
lyssae, 405
xanthogenum, 405
Oenobacillus, 50, 406
abbae, 50, 406
Omelianskillus, 100, 266, 406
hydrogenicus, 100, 406
methanicus, 406
Oospora, 50, 53, 57, 58, 163, 164, 166,
167, 333, 403, 406, 496
Ophidobacteries, 406, 524
Ophidomonas, 38, 407, 408, 472, 511
jenensis, 407, 408
sanguinea, 407
Ophryothrix, 352, 353, 408
thurethana, 408
Orchadaceae, 276
Oscillaria, 23, 37, 234, 235, 236
Oscillariaceae, 19

- Oscillariaceae*, 20, 355
Oscillatoria, 18, 234, 461
 ochracea, 318
Oscillatoriaceae, 24
Oscillospora, 77, 332, 367, 408
 guillemondii, 408
Ourococcus, 521
Oxydobacteriaceae, 68, 69, 152, 194, 245,
 248, 331, 370, 401, 402, 409
Pacima, 31, 35, 40, 197, 296, 306, 311,
 379, 409, 410, 437, 475, 524
 caucasica, 296, 409
 cecu, 409
 cholerae-asiatricae, 311, 409, 437
 finkleri, 409
 lewisii, 409
 loeffleri, 409
 putrificans, 409
Paciniae, 410
Palmella, 302, 410, 464
 infusum, 216
 maritima, 410
 prodigiosa, 410
Palmellaceae, 24
Pankistophyton, 410
 ovatum, 213, 410
Parachromatium, 410
Paracloster, 48, 60, 266, 267, 410
 butyraceus, 411
Paracoccaceae, 61, 65, 171, 270, 411
Parameningococcus, 411
Paramoecium, 15
Paraplectrum, 48, 60, 411
 foetidum, 411, 438
 peroniella, 411
Paraschizae, 289, 412
Paraspirillum, 73, 83, 412, 471
 vejvodskii, 412
Passalus cornutus, 286, 300
Pasteurella, 35, 40, 82, 87, 89, 93, 99, 105,
 197, 212, 272, 290, 291, 306, 405,
 412, 413, 414, 415
 cholerae-gallinarum, 412, 414
 davarnae, 412
 gallicida, 414
 suilla, 412
 weichselbaumii, 291
Pasteurellaeae, 92, 93, 95, 96, 104, 105,
 414
Pasteuria, 35, 40, 197, 378, 412, 415
 ramosa, 35, 378, 415
Pasteuriaceae, 415
Pasteuriaceae, 40, 197, 415
Pectinobacter, 415
 amylophilum, 415
Pectobacillus, 71 245, 416
Pediococcus, 29, 42, 51, 59, 60, 331, 416,
 417, 426, 458
 acidus-lactici, 416
 albus, 416
 cerevisiae, 29, 416, 417
 hennebergii, 417
 violaceus, 436
Pediococcus, 416, 417
Pedioplana, 64, 417
 haeckeli, 417
Pelochromatium, 253, 417
 roseum, 417
Pelodictyon, 253, 417
 clathratiforme, 417
Pelagoloea, 253, 417
 chlorina, 417
Peloploca, 418
 undulata, 418
 taeniata, 417
Pelosigna, 418
 cohnii, 418
Pelosphaera, 418
 rotans, 418
Pentapteris, 455
Peptonococcus, 157, 334, 418
Peritricha, 419
Peritrichinales, 419
Peritrichinae, 69, 71, 156, 173, 287, 358,
 419, 431, 439, 500
Perroncitoa, 42, 419, 492
 scarlatinosa, 419
Petalobacillus, 420
Petalobacteria, 272, 420
Petalobacterium, 420
Petalococcus, 21, 273, 420
Petalococcus, 420
 aceti var *Duclaux*, 420
Petaloglypha, 420
Peziza, 487

- Pezizales*, 164
Pfeifferella, 82, 96, 98, 106, 212, 360, 420, 421
 malleri, 420
Photobacillus, 421
Photobacter, 169, 339, 421
Photobacter fischeri, 421
 indicum, 421
 luminosum, 421
 phosphorescens, 421
 splendidum, 421
 splendor, 421
Photobacterium, 38, 421
 annulare, 422
 carabicum, 422
 coronatum, 422
 degenerans, 422
 delgadense, 422
 fischeri, 421
 glutinosum, 422
 indicum, 421, 422
 luminosum, 421
 papillare, 422
 phosphorescens, 421, 422
 tuberosum, 422
Photobakterium, 422
Photomonas, 99, 422, 434
Photospirillum, 422
 dunbari, 422
Phragmidiothrix, 29, 31, 34, 40, 43, 46, 48, 53, 55, 58, 59, 74, 250, 282, 310, 345, 351, 353, 423, 424
 multiseptata, 423, 424
Phycobacteriaceae, 74, 424
Phycochromaceae, 24
Phytobacter, 424
 lycopersicum, 424
Phytomonas, 105, 424, 425
 campestris, 425
Phytomyza, 425, 443, 444
 leguminosarum, 425
 lupini, 425
Phytomyzini, 425, 443
Phytozoidia, 18
 Plague group, 67
Planococcus, 47, 51, 54, 56, 58, 60, 74, 76, 77, 174, 270, 311, 313, 425, 426
 citreus, 426
 europaeus, 426
Planomerista, 77, 426
 ventriculi, 426
Planosarcina, 47, 51, 54, 56, 58, 60, 76, 270, 311, 331, 426, 427, 455
 agilis, 426, 427
 mobilis, 426
Planostreptococcus, 76, 427
Plasmodiophora, 425
Plectonema, 37
Plectrideae, 51
Plectridae, 60, 196, 427
Plectridium, 49, 51, 60, 80, 197, 242, 268, 398, 411, 427, 428, 438
 paludosum, 427
 putrificum, 427
 tetani, 398, 427
Plectrillum, 49, 232, 428
Plectrinum, 49, 233, 428
Plennobacterium, 64, 428
Plennobakterium, 428
Pleurumices, 163
Pleurococcus, 428
 bergeli, 250, 428
 roseo-persicinus, 265, 428
Pleurospora, 278, 310, 428
Plocamobacterium, 429
 crassum, 429
Pneumobacillus, 331, 429
 liquefaciens, *bovis*, 429
Pneumococcus, 292, 336, 429
 flavescens, 429
 gutta-cerei, 429
 lanceolatus, 429
 lichenoides, 429
 mucosus, 429
 Point of departure, 117, 148
Pollendera, 178, 179, 200, 429, 430
 anthracis, 429
Pollendra, 430
Polyangium, 15, 62, 74, 86, 107, 392, 393, 395, 430
 compositum, 430
 fusum, 430
 primgenum, 430
 septatum, 430
 simplex, 430
 sorediatum, 430

- Polyangium vitellinum*, 15, 255, 430.
Polybacteria, 30, 185, 431
 catenata, 431
 sulphurea, 431
Polycephalum, 255, 431
 aurantiacum, 431
Polycoccus, 22, 37
Polycystis, 22, 37
Polydesmus granulatus, 298
 virginensis, 182, 298, 299
Polyedrium, 391
Polygastrica, 17
 Polynomial nomenclature, 110
 Priority, 116, 141
Propionibacterium, 71, 156, 157, 419, 431
Propionococcus, 99, 431
Proteae, 92, 93, 95, 265, 432
Proteobacter, 432
 pseudopulcher, 432
 septicum, 432
 shatol, 432
Proteus, 15, 31, 43, 82, 93, 96, 98, 105, 179, 181, 200, 202, 212, 232, 357, 358, 359, 432, 433, 434, 486, 491, 519, 531
 mirabilis, 432
 vulgaris, 432, 433
 zenkeri, 432, 531
Protobacteries, 178, 433
Protococcus, 265, 433
 dentalis, 433
 roseo-persicinus, 433
Protomycetum, 434, 486
 recurrentis, 434, 486
Protophyta, 24
Pseudobacterium, 434
Pseudodiplococcus, 434
 pneumonicus, 434
Pseudodysentery group, 93
Pseudomeningococcus, 434
Pseudomonadaceae, 87, 88, 89, 99, 434
Pseudomonadales, 419, 434
Pseudomonas, 47, 54, 57, 61, 68, 74, 75, 82, 87, 89, 99, 104, 196, 207, 212, 225, 229, 232, 288, 311, 339, 358, 359, 402, 422, 434, 435, 436, 437
 aeruginosa, 229, 434, 436
 pyocyanea, 229, 434
 radicicola, 425
 violacea, 435, 436
Pseudorhizobium, 56, 437
 ramosum, 437
Pseudosarcina, 437
Pseudosarcine, 437
Pseudospira, 311, 409, 437, 481
Pseudospirillum, 312
Punctula, 29, 437
 cubica, 438
 glomerata, 438
 rosea, 437
Punctum, 438
 saltans, 438
Pupella, 16, 527
Putribacillus, 71, 438, 439
 putrificus, 438
 vulgaris, 438
Putribacteriaceae, 69, 71, 241, 419, 438, 439
Putriclostridium, 99, 245, 268, 439.
Putrificoideae, 100, 266, 302, 361, 370, 397, 398, 439, 440, 441, 451, 463, 513, 528
Putrificus, 438, 439.
 brenstocki, 439
Pyobacillus, 439
 capsulatus cuniculi, 439
Pyobacterium, 439, 440
 fischeri, 299, 439
Pyococcus, 440

Radiobacter, 440
Radiooccus, 440
Rapanea, 365
Rasmussenia, 39, 213, 338, 348, 353, 440
 anceps, 213, 440
 buccalis, 440
 gigantea, 440
 maxima, 440
 variabilis, 440
Recordillus, 439, 440
 fragilis, 441
Reducibacteriaceae, 68, 70, 248, 441, 466, 473
Regillus, 439, 441
 progrediens, 441

- Rhabdochromatium*, 56, 70, 255, 360, **441**, **442**
fusiforme, 441
minus, 441
roseum, 441
Rhabdomonas, 32, 36, 70, 85, 220, 256, 360, 441, **442**
rosea, 441, **442**
Rhizobacterium, **442**
japonicum, 442
Rhizobinae, 81, 212, **444**
Rhizobium, 38, 81, 87, 89, 96, 98, 103, 195, 258, 400, **442**, **443**, **444**
leguminosarum, 425, **443**
Rhizobius, 442
Rhizomonas, 69, 169, 434, **443**, **444**
Rhodobacillus, 64, 75, 86, 192, **444**, **448**
palustris, **444**
Rhodobacteria, 63, 192, **444**, **445**
Rhodobacteriaceae, 55, 68, 70, 73, 74, 84, 176, 248, 256, 326, 342, **445**, **447**, **448**, **449**, **450**, **502**
Rhodobacterioidae, 84, 85, **444**, **445**, **449**, **450**
Rhodobacteroidae, **445**
Rhodobacterium, 64, 86, 192, **445**
capsulatum, **445**
Rhodobacteriaceae, 417
Rhodocapsa, 63, 70, 85, 192, 256, **445**, **502**
suspensa, **445**
Rhodocapsaceae, 63
Rhodococcus, 44, 61, 63, 65, 70, 76, 80, 87, 90, 92, 97, 104, 192, 313, 368, 373, **446**, **447**, **449**, **506**
capsulatus, **447**
erythromyxa, **446**
fulvus, **447**
rhodochrous, **446**
roseus, **447**
Rhodocystis, 63, 70, 85, 192, **445**, **448**, **504**
gelatinosa, **448**
Rhododictyon, 70, **448**, **506**
Rhodomonas, 70, 257, **448**
Rhodonostoc, 63, 87, 192, **445**, **448**
capsulatum, **448**
Rhodopolycoccus, 70, **448**, **508**
Rhodosarcina, 70, 76, **448**, **449**, **509**
Rhodospira, 86, **445**, **449**
capsulata, **449**
Rhodospira, 75, 310, **449**
Rhodospirillum, 64, 70, 75, 86, 192, **445**, **449**, **511**
giganteum, **449**
photometricum, **449**
rubrum, **449**
Rhodotheca, 63, 70, 85, 192, 256, **449**, **450**, **512**
pendens, 350
Rhodovibrio, 64, 86, 192, **445**, **450**
parvus, **450**
Rickettsia, **450**, **451**
pediculi, **451**
prowazeki, **451**
quantana, **451**
Rivulillus, 101, 266, 267, **451**
vibrio, 101, **451**
Rivularia, 23, **490**
Rivulariaceae, 24
Robertsoniella, 439, **451**
primus, **451**
 Rules of botanical nomenclature, 112
Saccharobacillus, 59, **451**, **452**
pastorianus, **451**
Saccharobacter, **452**
Saccharomycetaceae, 24
Saccharomycetes, 24
Salmonella, 82, 93, 96, 105, 226, 298, **452**, **453**
paratyphi, **452**
schottmulleri, **453**
Saprospira, 72, 86, 88, 107, **453**, **481**, **484**
grandis, **453**
nana, **453**
Sarcina, 18, 19, 22, 25, 26, 27, 28, 30, 32, 33, 35, 37, 42, 43, 44, 45, 46, 47, 49, 51, 53, 54, 56, 58, 59, 60, 65, 66, 67, 71, 73, 74, 76, 77, 80, 87, 90, 92, 97, 104, 156, 157, 181, 269, 270, 276, 311, 313, 331, 341, 365, 368, 373, 375, 416, 427, 449, **453**, **454**, **455**, **456**, **457**, **458**, **459**, **509**, **519**, **521**
aurantiaca, **456**

- Sarcina citrea*, 456
flava, 456
littoralis, 305
lutea, 456
rosea, 455, 509
urinae, 454
ventriculi, 18, 322, 364, 453, 454, 455, 456, 457
Sarcinacoccus, 457
Sarcinaglobulus, 457, 458
punctum, 457
Sarcinastrum, 56, 457
urospora, 457, 458
Sarcinaceae, 25, 42, 212, 249, 274, 454, 458
Sarcinee, 274, 416, 458, 502
Sarcinoglobulus, 458
Schinzia, 425, 443, 458, 459
leguminosarum, 425, 443, 459
Schizofiti, 206
Schizomycetacea, 91, 394, 459, 460
Schizomycetaceae, 39, 233, 459, 460, 515
Schizomyceten, 460
Schizomycetes, 19, 24, 47, 49, 54, 74, 78, 87, 88, 102, 205, 206, 251, 289, 303, 309, 394, 412, 419, 459, 460, 461, 481, 516, 517
Schizophyceae, 251, 391, 501
Schizophyita, 22, 24, 397
Schizosiphon, 23
Schmidlea, 253, 461
luteola, 461
Schuetzia, 42, 461, 492
disser, 461
equi, 462
lagerhermia, 462
laughlini, 461
poelsii, 461
Sclerothrix, 38, 200, 275, 378, 385, 386, 462, 463
calitrichae, 462
kochii, 386, 463
tuberculosis, 378, 462
beigelianum, 250
Scytonema, 260
Scytonemaceae, 24
Seguinillus, 439, 463
aerofetidus, 463
Semiclostridium, 62, 463
citreum, 463.
commune, 463
flavum, 463
rubrum, 463
Septicemia group, 67.
Serratia, 15, 77, 82, 104, 203, 212, 274, 304, 369, 374, 463, 464, 465, 530
marcescens, 15, 464, 465
marescens, 464
subtilis, 203, 464
Shigella, 93, 96, 298, 314, 465
dysenteriae, 465
flexneri, 314
Siderocapsa, 71, 80, 465
major, 72, 466
treubii, 72, 465, 466
Siderocapseeae, 80, 466
Siphonomyxa, 466
nosocomii viennensis, 466
Solidococcus, 70, 359, 466
Solidovibrio, 70, 441, 466
Spermotira, 23
Sphaerella, 325
Sphaerobacteria, 21, 35, 38, 273, 387, 467, 469
Sphaerobacteries, 375, 467
Sphaerococcaceae, 467
Sphaerococcus, 467
conferioides, 467
lactis acidii, 467
Sphaerokolhus, 467
Sphaerotilus, 17, 20, 27, 32, 34, 39, 44, 45, 52, 55, 74, 83, 106, 251, 261, 262, 264, 467, 468, 469
glaucus, 468
lacteus, 467
natans, 17, 467
ochraceus, 468
thermalis, 366, 467
Spherobacteria, 469
Sprella, 469, 470
canis, 469
regaudi, 470
Spirikna, 16, 470, 527
Spirillaceae, 47, 50, 51, 52, 57, 59, 60, 64, 67, 73, 74, 79, 83, 87, 88, 89, 91, 94, 97, 103, 104, 209, 297, 308, 309, 326, 379, 384, 412, 470, 474, 479, 480, 526, 527.

- Spirilleae*, 34, 40, 197, 471
Spirillum, 18, 216, 471, 523, 527
Spirillineae, 19, 471
Spirillobacteriaceae, 273, 471, 476
Spirilloflagellata, 471
Spirillum, 17, 18, 19, 20, 21, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37, 40, 43, 44, 45, 46, 48, 50, 52, 53, 54, 57, 59, 60, 62, 64, 67, 68, 70, 73, 74, 75, 77, 83, 87, 90, 94, 97, 104, 197, 199, 206, 207, 215, 217, 306, 308, 312, 318, 379, 407, 434, 441, 449, 459, 466, 470, 471, 472, 473, 474, 475, 476, 483, 486, 487, 511, 519, 522, 524, 525, 527, 528
desulfuricans, 432
giganteum, 470
obermeieri, 434, 438.
phcatlis, 471
proeclarum, 183
rubrum, 52, 449
rufum, 20, 472
sanguineum, 407
serpens, 312
spirillum, 474
tenuis, 17, 20, 312, 434, 472
undula, 17, 20, 52, 312, 471, 472, 473, 474, 523
violaceum, 436
volutans, 20, 389, 407, 471, 472, 474, 475
zeylanicum, 528
Spirobacillus, 38, 62, 474, 475, 476
cienkowski, 474
gigas, 475
zeylanicus, 528
Spirobacteria, 21, 35, 37, 43, 62, 207, 470, 472, 475
Spirobacteriaceae, 273, 471, 476
Spirobacterium, 476
Spirobacterium, 476.
Spirobacillus, 75, 476
Spirochaeta, 17, 18, 20, 27, 28, 32, 35, 43, 57, 59, 62, 64, 70, 72, 73, 78, 86, 88, 107, 184, 215, 241, 312, 379, 470, 472, 475, 476, 477, 478, 479, 480, 481, 482, 484, 485, 513, 515, 522, 527.
balbianii, 64.
biflexa, 343
buccalis, 64
gallinarum, 64, 240
pallida, 483
phcatlis, 17, 20, 64, 78, 476, 477, 479, 480, 486
recurrentis, 485
Spirochaetacea, 480, 481
Spirochaetaceae, 64, 86, 107, 240, 284, 344, 453, 479, 480, 481, 482, 514
Spirochaetales, 102, 107, 461, 481
Spirochaete, 21, 23, 25, 26, 30, 33, 34, 37, 43, 44, 45, 56, 50, 52, 54, 59, 60, 73, 74, 77, 207, 308, 379, 470, 471, 477, 478, 481, 485.
cohnii, 477
ferruginea, 318, 477.
obermeieri, 52, 477, 479
pallidum, 513
phcatlis, 477, 478
schroeteri, 477
Spirochaetidae, 480, 481, 485
Spirochaetodea, 72, 481
Spirochaetaceae, 482
Spirochaete, 476, 482
obermeieri, 434
Spirochaetus, 17, 215, 482, 527
fulvus, 17, 482
Spiromonas, 25, 27, 32, 40, 197, 306, 312, 418, 434, 482, 483
cohnii, 418, 471, 483
volutilis, 482, 483
Spiromonas, 62, 78, 107, 379, 478, 483, 484, 513
recurrentis, 78
Spiromonaceae, 283, 453, 484
Spirophillum, 53, 64, 70, 318, 350, 484, 485, 515
ferrugineum, 319, 484, 485
Spiroschaudinna, 240, 241, 481 485
duttoni, 485
marcouxi, 485
recurrentis, 485
Spirosoma, 47, 54, 57, 59, 62, 74, 75, 78, 94, 297, 312, 390, 470, 471, 473, 475, 485, 486
Spirothrix, 434, 486
Spirulina, 23, 37, 43, 179, 181, 318, 319, 432, 486, 487
ferruginea, 318, 486

- Spirulina plicatilis*, 486
Sporonema, 18, 216, 487
 gracile, 368, 487
Sporosarcina, 157, 487
Sporospirillum, 487
Staphylococcus, 30, 35, 42, 43, 44, 65, 66,
 73, 76, 80, 87, 90, 96, 97, 104, 171,
 172, 173, 193, 241, 320, 358, 373, 375,
 376, 440, 487, 488, 489, 492
 albus, 172, 488
 aureus, 171, 172, 193, 488, 489
 epidermidis, 173
 epidermidis albus, 173
 pyogenes albus, 171, 193, 488
 pyogenes aureus, 66, 171, 193, 488
Staphylokokken, 489
Staphylokokkus, 489
Stigmatella, 19, 254, 489
 aurantaca, 489
Stoddardillus, 101, 267, 489, 490, 529
 egens, 101, 490
Streptothrichia, 44, 490
 bornetii, 44, 490
Streptothrichia, 490
Streptobacillus, 490, 491
 anaerobicus-rectus, 330
 lebens, 490
 pseudotuberculosis rodentium, 490
 taette, 491
 terrae, 490, 491
Streptobacter, 200, 491
Streptobacteria, 200, 273, 369, 491
 gigas-pericardii, 491
Streptobacterium, 419, 491
 foetidum, 491, 492
Streptobazillus ulceris mollis, 490
Streptococceae, 42, 65, 79, 80, 87, 91,
 103, 104, 271, 274, 293, 357, 396, 397,
 411, 488, 492, 494, 495
Streptococcee, 195, 274, 461, 492
Streptococcus, 273, 492
Streptococcus, 23, 30, 33, 35, 37, 42, 43,
 44, 45, 46, 47, 49, 51, 53, 54, 56,
 58, 59, 60, 66, 67, 71, 74, 76, 77, 80,
 87, 90, 92, 97, 104, 156, 157, 180, 184
 195, 242, 269, 270, 313, 331, 341, 356,
 357, 363, 365, 374, 375, 411, 419, 427,
 447, 488, 492, 493, 494, 495
 bombycis, 380
 equi, 462
 erysipelatis, 303, 493, 494
 erysipelatos, 493, 495
 lacticus, 431
 lactis, 219
 mesenteroides, 66, 239, 494
 pyogenes, 380, 493, 494
 rubiginosus, 419
 violaceus, 258, 436
 xanthopyreticus, 195
Streptokokken, 495
Streptokokkus, 43, 495
Streptomesobacterio, 353
Streptomicrococcus, 495
Streptospirillum, 495
Streptothrichaceae, 73
Streptothrix, 23, 26, 32, 37, 43, 48, 53
 57, 59, 68, 164, 165, 166, 167, 246,
 250, 251, 252, 259, 261, 262, 402, 403,
 406, 495, 496, 497, 516.
 foersteri, 23, 73, 163, 164, 404, 496
 fusca, 495
 paulotrophus, 159
Streptovibrio, 497
Stylopyga orientalis, 198
Sulfomonas, 69, 497, 498, 502
 denitrificans, 498
Sulphur bacteria, 36, 63, 176
Synechococcus, 22, 37
Swine erysipelas group, 67
Synonyms, 124, 127, 148

Termobacterium, 59, 498, 500
 aceti, 498
 album, 498
 erythrinum, 498
 fuscescens, 498
 iridescens, 498
 lutescens, 498
Termobakterien, 498
Testacea, 14
Tetrabacterium, 498
Tetracoccus, 419, 499, 500
 butyri, 499
 casei, 499
 liquefaciens, 499
 mastitidis, 499

- Tetracoccus mycodermatus*, 499
 pyogenes albus, 499
 pyogenes aureus, 499
Tetradiplococcus, 499
 filiformans lodzensis, 499
Tetraedron, 391
Tetragenus, 500
 citreus, 500
 tardissimus, 500
Tetrakokklus, 500
 variola, 500
Thallophyta, 24
Thallosporales, 94
Thermoactinomyces, 500
 lanuginosus, 500
 vulgaris, 500
Thermobacillus, 500
Thermobacterium, 419, 498, 500, 501
 aceti, 500
 bulgaricum, 501
 cereale, 501
 lactis, 501
 zeidlerii, 501
Thiobacillus, 61, 103, 501
 denitrificans, 501
 thio-oxidans, 501
 thioparus, 498, 501
Thiobacter, 501
Thiobacteria, 54, 55, 61, 74, 501
Thiobacteriaceae, 68, 69, 248, 498, 501,
 503, 506
Thiobacteriales, 79, 84, 87, 88, 91, 100,
 102, 107, 154, 445, 459, 461, 502
Thiobacterium, 501
Thiocapsa, 36, 42, 55, 70, 85, 445, 449,
 458, 502, 503
 roseo-persicina, 502, 503
Thiocapsaceae, 55, 63, 503
Thiocapseeae, 84, 85, 256, 503, 504, 509,
 510
Thiococcus, 70, 502, 503
Thiocysts, 36, 41, 55, 70, 85, 186, 306,
 311, 342, 447, 449, 503, 504
 rufa, 503.
 violacea, 436, 503, 504.
Thioderma, 53, 85, 504, 506.
Thioderma roseum, 504
 rubrum, 504
Thiodictyeae, 40, 197
Thiodictyon, 36, 40, 55, 70, 85, 175, 176,
 197, 417, 448, 505
 elegans, 505, 506
Thiomonas, 69, 257, 502, 506
Thiopedia, 36, 55, 70, 311, 343, 447, 506
 rosea, 305, 506
Thiopediaceae, 55, 63, 506
Thiopediaceae, 84, 85, 256, 343, 504, 506
Thiophysa, 61, 84, 154, 506, 507, 509
 volutans, 507
Thioploca, 64, 84, 237, 507
 schmidlei, 508
Thioplococcus, 36, 42, 55, 70, 85, 175,
 176, 189, 310, 448, 458, 508
 ruber, 508, 509
Thiorhodaceae, 63, 256, 444
Thiosarcina, 36, 55, 70, 75, 85, 311,
 448, 503, 509
 rosea, 509
Thiosphaera, 53, 85, 503, 509, 510
 gelatinosa, 509, 510.
Thiosphaerella, 509.
 amykifera, 509
Thiosphaerion, 53, 85, 503, 510
 violaceum, 510.
Thiospira, 510
 bipunctata, 510
 vinogradskii, 510
Thiospirillum, 36, 56, 70, 75, 85, 255,
 256, 312, 407, 408, 449, 472, 502, 511,
 512
 jenensis, 511
 sanguineum, 407, 511
Thiothece, 36, 41, 55, 70, 85, 175, 176
 186, 306, 355, 450, 511, 512
 gelatinosa, 511, 512
Thiothrix, 36, 48, 50, 52, 53, 55, 58,
 61, 70, 74, 84, 236, 237, 250, 352,
 353, 512, 513, 515
 nuvea, 510, 513
Thiotrix, 513
Tillatna, 513
 diphtherica, 513
 scarlatinosa, 513.
Tisserillus, 439, 513
 paraputrificus, 513.
Torula, 221

- Torula aceti*, 518.
Treponema, 62, 63, 64, 72, 78, 86, 88,
 107, 241, 380, 478, 479, 481, 482,
 483, 513
pallida, 64, 78, 513, 514
Treponemadae, 480, 514
Trichobacteria, 177, 324, 514, 516
Trichobacteriaceae, 52, 60, 69, 70, 248,
 289, 349, 484, 514, 515
Trichobacteriacei, 326
Trichobacterinae, 52, 60, 515
Trichobakterien, 515
Trichogenae, 34, 39, 283, 459, 515, 516
Trichomycetes, 67, 497, 515, 516
Trichomyzeten, 516
Tricogene, 264, 353, 516
Trichosporum, 250
beigelii, 250
Trochascia, 383
Trypanoplasma, 63
Trypanosoma, 63, 478
Trypanosomidae, 62
 Type, 142
 Type basis code, 145
 Typonym, 148
Tythrrix, 26, 202, 406, 517, 518
catenula, 517
claviforme, 517
distortus, 517
filiformis, 517
geniculatus, 517
scaber, 517
tenuis, 517, 518
turgidus, 517
urocephalum, 517
virgula, 517
Ulva, 185
Ulmia, 20, 153, 154, 388, 518, 519
aceti, 388, 518, 519
myxophila, 518
rubi-idae, 518
sambuci, 518
Umbina, 19, 153, 154, 519
aceti, 459, 519
Urobacillus, 24, 71, 173, 519, 521
duclauxii, 520
freundenreichii, 520
leubii, 520
maddoxii, 520
maqueli, 520
pasteurii, 519, 520
schützenbergii, 520
Urobacter, 520
Urocephalum, 19, 177, 520.
Urococcus, 24, 58, 521
ureae, 521
van tieghemi, 521
Urosarcina, 24, 521.
Urospora mirabilis, 457
Vallorillus, 101, 266, 521.
fallax, 101, 521
Verbenaceae, 484
Vermes, 14
Vibrio, 14, 15, 16, 17, 18, 19, 20, 21, 23,
 25, 26, 28, 30, 31, 32, 34, 35, 37, 40,
 43, 44, 46, 50, 51, 59, 60, 61, 62, 64,
 67, 73, 77, 83, 87, 89, 94, 97, 104,
 197, 199, 206, 212, 215, 216, 217, 218,
 231, 288, 294, 306, 312, 359, 363, 379,
 382, 406, 410, 422, 437, 459, 466, 470,
 471, 472, 475, 476, 497, 519, 521,
 522, 523, 524, 525, 526, 527, 528
amphibolus, 525
bacillus, 15, 16, 17, 20, 231, 368, 369
 522, 523
cholerae, 409, 524, 525, 526, 527
cholerae-asianicae, 51, 67, 526
comma, 526
cyanogenus, 218, 229, 523
lineola, 15, 16, 17, 20, 216, 363, 522,
 523, 524, 527
prolixer, 17, 20, 522
proteus, 525
rugula, 15, 16, 17, 20, 363, 522, 523
 524, 525
serpens, 15, 20, 522, 523, 524
spirillum, 15, 16, 471, 474, 522
subtilis, 17, 20, 369, 522
syncyanus, 17, 218, 219, 229, 522, 523.
synxanthus, 17, 213, 218, 522, 523
tremulans, 17, 20, 522
typhosus, 525
undula, 15, 231, 522
urnae, 525

- Vibrio xanthogenus*, 218, 523
zeylanicus, 527
 Vibrios, 526, **527**
 septique, 101, 279, 527
Vibriones, 217
Vibriomna, 17, 215, 523, **527**
Vibrionida, 18, 216, 523
Vibrionidsa, 16, 363, 470, 522, **527**
Vibrionieae, 259, 260, 312, 390, **527**.
Vibrioniens, 528
Vibriothrix, **528**
 zeylanica, 528
 Vienna code, 112
 Volvox, 14, 15

Weinbergillus, 439, **528**
 histolyticus, 528
Welchillus, 101, 267, 489, **529**
 aerogenes, 101, 529
Wesenbergus, 93, 96, 298, **529**
 wesenbergi, 529
Winogradskyia, 41, 197, 310, 337, **529**.
 ramagera, 529

Zaogalactina, 15, 304, **530**
 imetropha, 15, 464, 530
Zoagalactina, **530**
Zoogalactina, 374, **530**
Zoogloea, 19, 216, 218, 220, 274, 320,
 530
 bergeliana, 250
 pulmonis equi, 240, 241, 530
 ramagera, 217, 529, 530
 termo, 216, 217, 530
 Zoological Code of Nomenclature, 136
Zopfiae, 96, 104, 105, **531**, 532
Zopfiae, 531
Zopfiella, 31, 34, **531**
 tumescens, 531
Zopfielleae, 34
Zopfius, 96, 98, 105, 338, **531**
 zopfi, 531, 532
Zygobacterium 35, **532**
Zymosotis, 532
 escularis, 532
 regularis, 532

AUTHOR INDEX

- Abel, 300
 Adamson, 371
 Affanasieff, 161
 Agardh, 316, 325, 332, 343, 363, 433, 467
 Altana, 500
 Amato, 470
 Ambroz, 287
 Amiradzibi, 363
 Arkwright, 450
 Arlong, 429
 Avery, 429
 Avenfeld, 291

 Babes, 43, 186, 189, 195, 200, 203, 208,
 271, 272, 296, 325, 375, 432, 455,
 488, 494
 Bacot, 450
 Baker, 226, 227, 230, 477
 Balbiani, 168, 459
 Balcke, 29, 416
 Barendrecht, 341, 356
 Bartoszewicz, 499
 Baumgarten, 42, 43, 201, 204, 223, 227,
 260, 276, 293, 337, 353, 376, 456, 473,
 494, 495
 Beaurefaire-Arago, 245, 335
 Bechamp, 380
 Behrens, 401
 Beijerinck, 38, 50, 56, 58, 61, 153, 159,
 161, 169, 177, 193, 194, 290, 312,
 323, 324, 338, 339, 340, 341, 356, 357,
 380, 398, 399, 421, 425, 432, 443, 452,
 457, 498, 501, 519, 521
 Benecke, 73, 196, 270, 278, 288, 289, 326,
 384, 393, 445, 470, 501
 Berestnew, 163, 406, 497
 Bergamaschi, 464
 Bergey, 102, 154, 155, 156, 158, 165, 167,
 168, 170, 173, 176, 194, 195, 196, 203,
 209, 212, 226, 227, 230, 236, 237, 246,
 247, 251, 255, 256, 257, 258, 266, 269,
 272, 279, 283, 284, 289, 293, 298, 299,
 300, 302, 303, 304, 306, 313, 315, 328,
 329, 330, 332, 338, 340, 342, 344, 351,
 353, 357, 371, 373, 377, 385, 387, 393,
 394, 395, 396, 397, 399, 400, 402, 414,
 420, 424, 425, 428, 430, 433, 436, 442,
 444, 445, 446, 447, 448, 449, 450, 453,
 457, 461, 465, 469, 471, 474, 480, 484,
 492, 495, 501, 502, 503, 504, 505, 506,
 507, 508, 509, 510, 511, 512, 513, 514,
 527, 531, 532
 Bergonzini, 28, 221, 257, 258, 284, 285,
 286
 Berkeley, 19, 254, 489
 Bettencourt, 61, 294, 333
 Biedert, 337
 Bienstock, 439, 513
 Billet, 186, 189, 200, 223, 240, 265, 276,
 291, 294, 337, 348, 364, 365, 375, 390,
 455, 473, 477, 481, 491, 494, 495, 499,
 525
 Billroth, 21, 185, 186, 187, 217, 272, 273,
 275, 276, 291, 320, 328, 347, 361, 366,
 371, 372, 373, 374, 382, 420, 457, 466,
 491, 492, 493, 495
 Bizio, 15, 463, 464
 Bizzozero, 469
 Blanchard, 62, 164, 295, 379, 403, 404,
 470, 473, 475, 478, 481, 483, 485, 513,
 526
 Bollinger, 23, 38, 162, 241, 295, 530
 Bongert, 58, 280
 Bonome, 434
 Bonordon, 430
 Bordono-Uffreduzzi, 363
 Borrel, 72, 191
 Borsa, 330, 381
 Bory de St Vincent, 16, 316, 363, 470,
 522, 527
 Borzi, 408
 Bosanquet, 73, 478, 479
 Bostroem, 167
 Bredemann, 244

- Breed, 96, 102, 165, 226, 227, 230, 464
 Broadhurst, 86
 Brown, C H, 13
 Brown, Margaret, 13
 Brumpton, 72, 157, 158, 161, 295, 333
 Brussoff, 312.
 Buchanan, R E, 78, 86, 102, 107, 153,
 154, 156, 158, 165, 167, 169, 172, 175,
 176, 191, 192, 193, 194, 197, 203, 208,
 211, 212, 225, 227, 236, 237, 242, 251,
 255, 256, 257, 258, 263, 266, 268, 271,
 279, 283, 284, 290, 293, 298, 303, 307,
 308, 314, 315, 328, 329, 330, 339, 341,
 342, 343, 351, 353, 367, 368, 373, 376,
 385, 387, 388, 393, 395, 396, 397, 399,
 401, 402, 412, 414, 420, 425, 427, 430,
 433, 436, 442, 443, 444, 445, 446, 447,
 448, 449, 450, 452, 453, 457, 460, 465,
 466, 469, 471, 474, 480, 481, 488, 492,
 494, 502, 503, 504, 506, 507, 508, 509,
 510, 511, 512, 513, 526
 Buchner, 524, 526
 Buder, 254
 Bumm, 291
 Burdon-Sanderson, 380
 Burnill, 414
 Busk, 322

 Cambier, 421, 422
 Carter, 249
 Caruel, 206
 Caspary, 364, 365
 Castellani, 91, 95, 102, 165, 173, 196,
 197, 209, 211, 226, 227, 230, 234, 243,
 249, 251, 264, 265, 277, 293, 297, 298,
 299, 301, 306, 308, 314, 315, 322, 323,
 325, 329, 333, 343, 351, 373, 378, 385,
 387, 394, 397, 398, 399, 400, 404, 405,
 414, 432, 433, 447, 452, 453, 457, 459,
 465, 471, 480, 485, 492, 495, 502, 514,
 516, 527, 528, 529.
 Cattani, 330.
 Certes, 475
 Cesari, 241
 Chalmers, 91, 95, 102, 162, 165, 173,
 196, 197, 209, 211, 226, 227, 230, 234,
 243, 249, 251, 264, 265, 277, 293, 295,
 297, 298, 299, 301, 306, 308, 314, 315,
 322, 323, 325, 329, 333, 343, 351, 373,
 377, 385, 387, 394, 397, 398, 399, 400,
 404, 405, 414, 432, 433, 447, 452, 453,
 457, 459, 465, 471, 480, 481, 485, 492,
 495, 502, 514, 516, 527, 528, 529
 Chantemasse, 383
 Chapman, 324
 Chatton, 77, 183, 332, 366, 408
 Chester, 52, 53, 56, 202, 203, 204, 208,
 224, 227, 236, 237, 250, 262, 270, 279,
 282, 348, 376, 379, 384, 385, 386, 406,
 424, 426, 427, 435, 456, 470, 473, 478,
 485, 494, 497, 512
 Choukevitch, 330, 370
 Christopherson, 102, 295, 333, 404
 Chudiakov, 232
 Cienkowski, 188, 190, 261, 355, 356
 Clairmonte, 300
 Clark, Clarissa, 13
 Clark, W M, 170, 278
 Clegg, 162, 246, 406, 497
 Clements, 12, 123, 201, 207, 209, 224,
 236, 237, 250, 375, 393, 394, 440, 460
 Coats, 363
 Cohn, Ferdinand, 19, 20, 21, 22, 110,
 163, 171, 188, 198, 199, 203, 204, 205,
 210, 212, 216, 217, 218, 219, 220, 222,
 227, 228, 229, 235, 252, 260, 263, 264,
 265, 277, 281, 288, 301, 302, 320, 328,
 347, 355, 365, 368, 369, 370, 371, 372,
 373, 374, 378, 379, 389, 390, 392, 397,
 406, 407, 442, 454, 459, 467, 468, 470,
 472, 475, 477, 481, 486, 493, 496, 520,
 524, 530
 Collin, 183
 Confare, Ruth, 13
 Conn, 67, 165, 202, 204, 225, 226, 227,
 230, 368, 380, 435
 Cooke, 431
 Coppen-Jones, 192, 195, 326
 Corda, 110, 163, 164, 495, 496, 497
 Cornil, 43, 189, 200, 203, 208, 271, 272,
 296, 375, 432, 455, 488, 494
 Corsini, 237
 Cunningham, 365
 Curtiss, 254, 489

 Dallinger, 220, 227.

- Dangeard, 44, 306, 377.
 Davane, 20, 178, 209, 210, 212, 217, 227,
 472, 523
 Davis, 170, 278
 DeBary, 29, 179, 180, 199, 203, 204, 355,
 472
 DeCandolle, 264
 deKruyff, 63, 357
 DeLanessan, 27, 28
 Delbruck, 498
 D'Herelle, 272
 Dennstatt, 458
 Desmazieres, 388, 487
 DeToni, 38, 39, 161, 170, 171, 174, 177,
 180, 184, 189, 195, 197, 200, 203, 213,
 223, 227, 233, 236, 240, 241, 248, 249,
 257, 258, 261, 264, 267, 271, 272, 274,
 275, 276, 278, 282, 283, 287, 289, 291,
 294, 296, 302, 305, 306, 310, 311, 312,
 315, 318, 327, 331, 332, 336, 337, 338,
 342, 343, 344, 346, 348, 353, 355, 356,
 357, 360, 364, 366, 369, 373, 375, 379,
 380, 390, 391, 392, 396, 402, 404, 407,
 409, 415, 417, 419, 423, 428, 429, 440,
 441, 455, 458, 459, 461, 462, 468, 471,
 472, 475, 477, 481, 483, 487, 488, 491,
 492, 494, 496, 502, 503, 505, 506, 508,
 509, 511, 515, 517, 518, 521, 525, 529,
 531, 532
 Diesing, 216, 361
 Dobell, 72, 73, 283, 314, 367, 412, 471,
 479, 483, 513
 Dochez, 429
 Doflein, 62, 72, 283, 325, 479, 513
 Donaldson, 370
 Dopter, 411
 Drysdale, 227
 Dubosco, 469
 Duclaux, 26, 115, 159, 160, 177, 287, 517
 Dujardin, 17, 191, 215, 216, 217, 218,
 223, 227, 471, 476, 482, 523, 524, 528
 Dunbar, 422
 Duncan, 450
 Duncker, 163
 Eberth, 530
 Edington, 184, 419
 Edkms, 470
 Ehrenberg, 16, 17, 191, 213, 214, 215,
 218, 226, 229, 290, 316, 317, 319, 320,
 322, 363, 368, 382, 407, 471, 474, 476,
 477, 481, 482, 486, 522, 524, 526, 527.
 Eidam, 468
 Eisenberg, 276, 325, 337, 416, 417, 467.
 Ellis, 53, 64, 72, 202, 204, 208, 225, 227,
 251, 252, 262, 265, 270, 278, 282, 318,
 319, 349, 363, 376, 379, 405, 424, 426,
 427, 435, 456, 470, 473, 478, 481, 484,
 485, 494
 Elser, 434
 Engelmann, 222
 Engler, 74, 118, 423, 424, 449.
 Enlows, 95, 153, 158, 169, 175, 180, 183,
 184, 185, 241, 245, 247, 252, 253, 254,
 269, 285, 286, 287, 288, 290, 297, 303,
 304, 320, 323, 327, 328, 333, 335, 343,
 361, 362, 363, 366, 367, 371, 380, 388,
 415, 421, 422, 424, 436, 438, 450, 454,
 457, 458, 490, 500, 509, 510, 511, 519
 530, 531, 532
 Erichsen, 434
 Escherich, 222, 227, 301, 328
 Esmarch, 449
 Evans, 173, 242, 243
 Falkenheim, 455
 Famintzin, 46, 397
 Fantham, 480
 Faull, 394
 Fehleisen, 493
 Fick, 276
 Firtsch, 525
 Fischer, Alfred, 48, 50, 59, 174, 179
 180, 195, 197, 201, 204, 224, 227, 231,
 232, 236, 251, 261, 262, 266, 267, 269,
 270, 282, 290, 320, 325, 326, 327, 331,
 375, 376, 379, 410, 411, 413, 416, 425,
 426, 427, 428, 456, 470, 473, 477, 478,
 481, 494, 512, 514, 515, 526
 Fischer, Bernhard, 47, 422
 Fitz, 170, 244, 362
 Fitzgerald, 300
 Fleischner, 243
 Flugge, 31, 32, 65, 66, 189, 196, 200,
 203, 222, 225, 227, 235, 265, 270, 282,
 290, 291, 347, 375, 381, 422, 423, 455,
 456, 468, 470, 472, 473, 477, 488, 494,
 524, 526.

- Foa, 363
 Forbes, 364
 Forti, 50, 406
 Fortineau, 62, 304, 465
 Foulerton, 165
 Francke, 417
 Frank, 38, 425, 443, 458
 Frankland, 197
 Frenzel, 155, 381
 Freudenreich, 46, 203, 223, 227
 Fricke, 300
 Friedberger, 392
 Friedlander, 354, 429
 Friere, 286
 Frost, 72, 175, 176, 202, 204, 208, 225,
 236, 237, 251, 252, 255, 257, 270, 282,
 303, 341, 342, 376, 379, 424, 426, 427,
 435, 456, 470, 473, 478, 485, 494, 501,
 502, 503, 504, 505, 506, 509, 511, 512,
 515
 Fuchs, 28, 229, 523
 Fuhrmann, 470, 500

 Gamaleia, 38, 272, 525
 Gasperini, 163, 496
 Gedoelst, 164
 Glaser, 324
 Gleitsmann, 241, 479, 484, 485
 Gonder, 78, 283, 381, 479, 484, 513
 Gonnermann, 64, 392, 394, 395, 428
 Goodsir, 18, 453
 Grassberger, 178, 244, 323, 324, 411
 Griffith, 19, 290, 317
 Griffiths, 71, 330
 Grumberg, 383
 Groenewege, 424
 Gross, 72, 283, 453, 479, 483, 484
 Grove, 29, 188, 199, 203, 222, 227, 235,
 257, 261, 277, 281, 347, 355, 374,
 454, 459, 468, 472, 477
 Gruber, 44, 323, 377, 381, 455
 Guerbet, 491
 Guignard, 44, 490
 Guillebeau, 319, 320
 Guttman, 486

 Haase, 291
 Haass, 63, 161
 Haller, 20, 217, 278, 360, 373, 392, 513

 Hammer, 102
 Hankin, 190
 Hansen, 388, 416
 Hansgirt, 36, 37, 42, 189, 200, 205, 207,
 223, 235, 237, 257, 258, 260, 261, 270,
 282, 283, 287, 306, 307, 308, 310, 318,
 319, 331, 344, 345, 348, 351, 355, 356,
 371, 375, 383, 387, 390, 391, 407, 408,
 423, 460, 467, 472, 475, 477, 524
 Hanzawa, 194
 Harrison, 102
 Hartleb, 56, 333, 400, 401, 437.
 Harz, 23, 166, 167, 295, 406
 Hassall, 521
 Hauser, 31, 432
 Hazen, 325
 Heim, 72, 73, 203, 225, 227, 236, 262,
 282, 350, 456, 479, 481, 489, 494, 497,
 512, 516
 Heller, Hilda Hempl, 99, 100, 179, 205,
 266, 268, 297, 302, 309, 310, 313, 329,
 359, 361, 370, 371, 382, 394, 397, 405,
 406, 427, 439, 440, 441, 451, 463, 489,
 502, 513, 521, 528, 529
 Hempl, 451
 Henfrey, 19, 264
 Henneberg, 451, 498
 Henry, 370, 528
 Hertwig, 163
 Hetsch, 73
 Hewlett, 53, 202, 204, 262, 376, 456,
 473, 494, 497
 v Hible, 330, 370
 Hiltner, 443
 Hinze, 61, 506
 Hochstetter, 484
 Hoelling, 72, 314
 Hoffman, 513
 Hohl, 246
 Hollig, 301
 Hoyer, 153, 154
 Huebner, 364
 Huntton, 102, 434
 Huppe, 111, 179, 180, 181, 184, 191, 200,
 203, 204, 207, 222, 227, 228, 234, 236,
 260, 261, 267, 270, 282, 300, 337, 345,
 348, 352, 355, 365, 375, 381, 423, 427,
 455, 472, 476, 477, 486, 487, 488, 494,
 495, 524

- Huss, 232
 Israel, 161, 167
 Itzigsohn, 217, 392, 530
 Jacqu , 491
 Jeantet, 191
 Jegunow, 506
 Jennings, 54, 192, 289
 Joffe, 501
 Johne, 171
 Jordan, 67, 164, 262, 349, 404, 497, 515
 Jouan, 191
 Kalchbrenner, 431
 Kallwacks, 300
 Karwacki, 272
 Kat, 421
 Kendall, 58, 202, 204, 208, 224, 227, 230, 232, 233, 236, 237, 250, 262, 270, 282, 376, 379, 386, 424, 426, 427, 456, 470, 473, 478, 481, 485, 494, 497
 Kern, 28, 296
 Khoury, 491
 Kirchner, 318, 442, 486
 Kitt, 241, 413
 Klebahn, 118
 Klebs, 20, 184, 320, 327, 337, 369, 380, 382, 500
 Klecki, 499
 Klein, 289, 370, 372, 411, 476
 Kligler, 171, 263, 350, 456
 Kline, 307
 Klinger, 274
 Klocker, 59, 208, 224, 270, 376, 417, 456
 Kolle, 73
 Kopke, 61, 333
 Koppanyi, 439
 Kossowicz, 341, 406, 428
 Krokiewicz, 293
 Krumwiede, 86
 Kruse, 177, 324, 382, 500
 Krzysztalawicz, 471
 Kuetzing, 18, 153, 154, 161, 234, 241, 252, 265, 290, 332, 365, 377, 518, 521
 Kuhn, 294
 Kunth, 276
 Kunze, 231
 Kurth, 222, 337.
 Kuttner, 299, 439
 Kutzing, 17, 38, 286, 301, 302, 317, 318, 321, 343, 344, 345, 347, 351, 364, 386, 388, 391, 462, 467
 Lachner-Sandoval, 163, 164, 168
 Lafar, 388
 Lagerheim, 56, 457, 458
 Lankester, 219, 227, 459
 Lanzi, 206, 220, 222, 234, 518
 Lauterborn, 64, 155, 253, 286, 381, 417, 418, 461, 507
 Lebailly, 469
 Lebert, 410, 434
 Lehmann, 49, 58, 109, 164, 168, 190, 202, 203, 207, 224, 227, 270, 279, 291, 300, 307, 333, 356, 364, 365, 375, 379, 385, 386, 406, 415, 416, 432, 456, 470, 473, 478, 481, 486, 488, 494, 497, 518, 524, 525, 526
 Leichmann, 242, 340
 Leidy, 18, 181, 258, 259, 280, 286, 298, 300, 301, 332, 523
 Leube, 271, 272
 Leveille, 378
 Levine, Max, 13
 Levy, 163
 Lewis, 434, 486
 Lieske, 165, 351
 Lign res, 157, 158, 159, 161, 413, 452, 453
 Lima, 450
 Lindau, 163, 322
 Lindley, 484
 Lindner, 29, 324, 416, 498
 Lingelsheim, 293, 294
 Link, 15, 430, 486
 Linnaeus, 110
 Lister, 219, 227, 242
 Loeffler, 278, 526
 Lohnis, 76, 160, 194, 203, 225, 227, 242, 318, 376, 437, 456, 473, 478, 494, 526.
 Lotsy, 63, 204, 425.
 Luders, 217, 227
 Ludwig, 44, 153, 154, 163, 201, 203, 223, 227, 236, 261, 282, 287, 331, 348, 355, 375, 390, 440, 443, 455, 468, 473, 477, 481, 494

- Luerssen, 26, 199, 203, 220, 222, 227,
 235, 294, 347, 374, 454, 472, 477, 481,
 524
 Lutz, 31, 275
 Lyall, 429
 Lyngbye, 410

 McIntosh, 314, 359, 513
 McNab, 24, 206, 291

 Maassen, 62, 463
 Macé, 185, 186, 189, 190, 202, 204, 207,
 210, 224, 263, 270, 348, 356, 376, 432,
 437, 517, 521, 525
 Maggi, 35, 160, 178, 180, 200, 222, 289,
 290, 294, 296, 320, 353, 355, 362, 372,
 375, 382, 406, 420, 433, 440, 467, 490,
 491, 497, 498, 517, 524, 532
 Magnin, 24, 188, 199, 203, 220, 227, 235,
 260, 347, 374, 390, 454, 469, 472, 496,
 524
 Magnus, 118
 Maire, 425
 Makrinov, 415
 Malerba, 320
 Mantegazza, 360
 Martelly, 361
 Marpmann, 467
 Masay, 491
 Matruchot, 383
 Matzuschita, 59, 202, 204, 224, 227, 376,
 446, 456, 473, 478, 481, 494, 526
 Meek, 379
 Mendes, 61
 Meneghini, 428
 Merrill, 158, 162, 165, 246, 295, 377, 404
 Messea, 177, 324, 359, 382, 419, 514
 Metschnikoff, 35, 38, 370, 385, 386, 415,
 462, 474
 Mettenheimer, 317.
 Meyen, 162, 364
 Meyer, 53, 75, 191, 242, 243, 307, 308,
 310, 312, 313, 370, 376, 379, 426, 427,
 435, 444, 447, 448, 449, 475, 476, 485
 Micellone, 294
 Miehe, 384, 385, 386
 Migula, 44, 47, 53, 54, 61, 155, 156, 172,
 175, 176, 178, 185, 189, 191, 201, 202,
 204, 207, 208, 223, 224, 225, 227, 228,
 236, 237, 250, 251, 255, 256, 257, 261,
 262, 264, 269, 270, 282, 283, 294, 305,
 307, 308, 311, 312, 318, 319, 322, 324,
 334, 341, 342, 343, 355, 356, 369, 375,
 376, 379, 386, 390, 397, 399, 401, 402,
 407, 415, 422, 424, 425, 426, 427, 435,
 436, 441, 442, 445, 446, 455, 468, 470,
 473, 477, 478, 481, 482, 483, 485, 488,
 490, 493, 494, 496, 501, 502, 503, 504,
 505, 506, 508, 509, 510, 511, 512, 517,
 518, 519, 520, 526
 Miller, 189, 327, 334, 359, 433
 Miquel, 24, 382, 421, 422, 499, 519, 520,
 521
 Miller, 189, 327, 334, 359, 433
 Miyoshi, 53, 504, 509, 510
 Molisch, 63, 64, 71, 192, 262, 265, 282,
 317, 318, 350, 405, 444, 445, 446, 448,
 449, 450, 465, 469, 485, 503
 Moore, 164
 Morax, 73, 291
 Moreno, 185
 Mori, 246
 Morren, 231, 369
 Mueller, O F, 14, 15, 16, 213, 214, 215,
 322, 368, 382, 471, 521, 522, 524, 526
 Muhlhauser, 439
 Muur, 291
 Musgrave, 162, 246, 406, 497

 Nadson, 253, 509
 Naegeli, 19, 153, 154, 405, 458, 459, 518,
 519
 Neisser, 322
 Neumann, 49, 58, 109, 164, 168, 190,
 202, 203, 207, 224, 227, 270, 279, 291,
 300, 307, 333, 356, 364, 365, 375, 379,
 385, 386, 406, 415, 416, 432, 456, 470,
 473, 478, 481, 486, 488, 494, 497, 518
 524, 525, 526
 Nicolle, 241
 Nijdam, 169
 Niklewski, 331, 332
 Nissen, 276
 Noguchi, 344
 Novy, 405
 Nowakowski, 258

- Obermeier, 434
 Oersted, 170, 305, 357
 Ogston, 193, 374, 487, 488, 493
 Olson-Sopp, 491
 Omeliansky, 406, 511
 Orla-Jensen, 68, 71, 99, 134, 152, 154,
 156, 157, 164, 169, 170, 173, 176, 194,
 195, 202, 203, 225, 227, 236, 237, 238,
 241, 244, 245, 247, 248, 252, 257, 262,
 268, 273, 276, 277, 280, 282, 287, 288,
 314, 318, 331, 333, 340, 342, 349, 357,
 358, 359, 370, 372, 376, 389, 401, 402,
 409, 416, 418, 419, 422, 431, 432, 434,
 438, 439, 441, 442, 443, 444, 445, 447,
 448, 449, 450, 456, 466, 468, 471, 473,
 476, 478, 483, 484, 487, 494, 497, 498,
 499, 500, 501, 502, 503, 504, 506, 508,
 509, 511, 512, 515, 519
 Overbeck, 446

 Pacini, 524, 525
 Parsons, 173, 193
 Pasteur, 278, 388, 416, 518, 527
 Perard, 77, 183, 332, 366, 408
 Perfiliev, 286
 Perkins, 300
 Perroncito, 162, 163, 272
 Perrone, 294
 Persoon, 16, 153, 154, 387
 Perty, 18, 191, 212, 216, 227, 256, 332,
 341, 368, 369, 370, 471, 477, 482, 487,
 523, 527
 Petit, 291
 Petruschky, 164, 349, 497, 515, 516
 Petschenko, 198, 297
 Pfeiffer, 161, 164
 Pfender, 62, 379, 478, 483, 513
 Pinoy, 73, 164, 165, 167, 277, 404
 Polk, 162, 246
 Potron, 166
 Pouchet, 525
 Poulsen, 457
 Prazmowski, 27, 177, 267, 269, 365
 Preisz, 490
 Prilleaux, 443
 Pringsheim, 244
 Prowazek, 450

 Quehl, 395, 430

 Rabenhorst, 19, 21, 216, 234, 241, 265,
 283, 305, 318, 321, 343, 344, 346, 364,
 365, 368, 471, 472, 477, 486, 518, 524
 Radais, 163, 261, 403, 406, 496
 Rafinesque, 484
 Ralfs, 290, 317, 363
 Reichenbach, 276
 Rertz, 161
 Renzende, 61, 333
 Rettger, 338, 531
 Rist, 490
 Ritchie, 291
 Rivolta, 24, 163, 164, 273, 274, 294, 295,
 337
 Robertson, 451
 Robin, 18, 182, 259, 322, 345, 346, 364,
 454
 Rogers, Ann, 61, 78, 171, 190, 193, 292,
 356, 411
 Rogers, 170, 191, 278, 363, 446
 Romer, 301
 Rosenbach, 30, 71, 171, 172, 193, 299,
 303, 335, 375, 487, 493, 494
 Rothberg, 173, 193
 Rothmann, 321
 Rougentzoff, 301
 Roze, 249
 Ruata, 272
 Rullmann, 401

 Sabouraud, 371
 Saccardo, 255, 395, 430, 489, 495, 518
 Salisbury, 245, 285, 532
 Sambon, 241, 485
 Sampietro, 161
 Sanna-Salaris, 320
 Sauvageau, 163, 261, 402, 406, 496
 Schambert, 371
 Schattenfroh, 178, 244, 323, 324, 411
 Schaudinn, 62, 367, 379, 380, 478, 483,
 513, 514
 Schewiakoff, 155, 156, 381
 Schindler, 485
 Schlegel, 161, 164
 Schmidlechner, 429
 Schmidt, 196, 202, 224, 236, 237, 250,
 251, 264, 270, 274, 282, 435, 456, 470,
 513, 515

- Schneider, 202, 204, 208, 225, 227, 270,
376, 379, 384, 435, 456, 470, 473, 481,
482, 494
- Schorler, 61, 265
- Schneider, 73, 237, 443
- Schroeter, 27, 32, 163, 189, 200, 203, 206,
208, 210, 218, 221, 222, 227, 229, 236,
255, 256, 259, 260, 261, 264, 265, 267,
270, 273, 282, 286, 288, 302, 307, 322,
331, 341, 342, 346, 348, 351, 353, 355,
356, 375, 378, 379, 380, 390, 425, 433,
443, 455, 458, 468, 472, 477, 481, 491,
494, 509, 518, 524, 526, 530
- Schutz, 293
- Schwarzwasser, 499
- Schwers, 319, 349, 362
- Seguin, 330, 361, 463, 528
- Seligmann, 324
- Sette, 15, 530
- Shaw, 242, 243
- Siedlicki, 471
- Silberschmidt, 161
- Summonds, 529
- Smith, Allan, J., 59, 202, 204, 208, 224,
227, 250, 262, 270, 282, 305, 348, 376,
379, 384, 386, 424, 426, 427, 435, 456,
470, 473, 478, 485, 494, 497
- Smith, Erwin F., 61, 160, 169, 170, 175,
176, 177, 178, 179, 180, 181, 192, 194,
195, 202, 204, 208, 210, 213, 224, 225,
227, 228, 229, 232, 233, 236, 237, 240,
241, 248, 250, 251, 255, 256, 257, 268,
270, 272, 275, 276, 277, 279, 282, 290,
291, 295, 297, 302, 304, 305, 308, 312,
315, 320, 321, 324, 326, 328, 331, 334,
336, 338, 340, 341, 342, 343, 349, 355,
356, 359, 361, 362, 363, 364, 366, 369,
372, 376, 377, 379, 380, 382, 390, 392,
393, 396, 397, 399, 401, 402, 403, 406,
408, 410, 411, 413, 415, 417, 419, 420,
421, 424, 425, 426, 427, 429, 430, 432,
435, 442, 443, 445, 452, 456, 462, 467,
468, 470, 473, 475, 478, 483, 485, 490,
491, 494, 497, 499, 500, 501, 502, 504,
505, 506, 508, 509, 510, 511, 512, 518,
519, 520, 521, 526
- Smith, 86
- Sollhed, 417.
- Spitz, 157, 158, 159, 161
- Sternberg, 45, 189, 201, 204, 220, 223,
227, 264, 267, 292, 351, 356, 364, 375,
455, 473, 477, 481, 487, 494, 525
- Stiles, 62, 113, 114, 121, 131, 379, 382,
474, 478, 481, 482, 483, 513
- Stockhausen, 194, 323, 324, 341
- Stoddard, 382, 390
- Stoklasa, 440
- Stormer, 443
- Streinz, 162
- Strickland, 113
- Stutzer, 333, 400, 401
- Swellengrebel, 64, 240, 470, 479, 480,
513
- Tavel, 294
- Terni, 163
- Thaxter, R., 15, 19, 46, 255, 287, 392,
393, 395, 430, 431, 489
- Thesing, 479, 481
- Thienmann, 249
- Thin, 221
- Thiercelin, 301
- Thuselton-Dyer, 459
- Thomson, 388
- Tilden, 486
- Tison, 425
- Tissier, 272, 301, 361
- Tizzoni, 330
- Tommasoli, 184, 185
- Trecul, 19, 27, 177, 269, 520
- Trelease, 222
- Trevisan, 18, 24, 31, 34, 38, 39, 161, 163,
171, 174, 176, 177, 180, 184, 186, 188,
189, 195, 197, 199, 200, 203, 206, 211,
212, 213, 220, 223, 227, 231, 233, 234,
235, 236, 240, 241, 248, 249, 257, 258,
260, 261, 264, 267, 271, 272, 273, 274,
275, 276, 278, 281, 282, 283, 287, 289,
291, 294, 296, 302, 305, 306, 309, 310,
311, 312, 315, 318, 322, 327, 331, 332,
335, 336, 337, 338, 342, 343, 344, 346,
347, 348, 351, 352, 353, 355, 356, 357,
360, 363, 364, 366, 368, 369, 370, 373,
374, 375, 379, 380, 389, 390, 391, 392,
396, 402, 404, 405, 407, 408, 409, 412,
414, 415, 416, 417, 419, 423, 428, 429,

- 434, 437, 440, 441, 442, 454, 455, 458,
459, 461, 462, 468, 471, 472, 475, 477,
481, 482, 483, 487, 488, 491, 492, 494,
496, 502, 503, 505, 506, 509, 511, 513,
515, 517, 518, 521, 524, 525, 527, 529
530, 531, 532
- Troili-Petersson, 61, 242
- Tsilinsky, 500
- Tulasne, 513
- Turczaninow, 476
- Turpin, 486
- Ucke, 490
- Unna, 184, 185, 337, 381, 490
- Valentin, 183, 332
- Van Laer, 451
- Van Tieghem, 24, 29, 185, 186, 188, 190,
199, 203, 220, 221, 222, 227, 235, 269,
282, 355, 356, 364, 365, 374, 390, 431,
437, 454, 477, 481, 524
- Veillon, 230
- Veley, 277
- Vianna, 245, 335
- Vincenzi, 500
- Von Baer, 16, 357
- Von Hibler, 179
- Vuillemin, 62, 77, 118, 119, 180, 203,
211, 225, 227, 228, 229, 250, 251, 258,
268, 279, 295, 297, 315, 331, 336, 343,
357, 360, 363, 365, 367, 369, 376, 377,
378, 379, 386, 387, 390, 396, 404, 410,
413, 415, 417, 426, 427, 435, 456, 457,
460, 463, 473, 474, 475, 478, 483, 484,
486, 491, 493, 494, 495, 501, 513, 526,
531
- Wade, 158, 162, 164, 165, 246, 295, 377,
404
- Waksman, 165, 501
- Wallroth, 163, 231, 406
- Ward, H Marshall, 11
- Warming, 155, 219, 418, 483, 503
- Weichselbaum, 31, 290, 292
- Weinberg, 329, 330, 361, 463, 528
- Weis, 196, 202, 224, 236, 237, 250, 251,
264, 270, 274, 282, 435, 456, 470, 513,
515
- Weiss, 256
- Weldin, J C , 13
- Wenner, 338, 531
- Werkman, Mrs C H , 13
- West, 71, 241, 266, 325, 330, 349, 486
- Wille, 325
- Winogradsky, 19, 36, 46, 64, 174, 175,
176, 189, 235, 236, 256, 265, 268, 269,
342, 343, 398, 401, 402, 407, 408, 441,
442, 502, 503, 505, 506, 508, 509, 511,
512
- Winslow, C-E A , 10, 61, 65, 78, 86, 96,
102, 153, 154, 158, 171, 172, 173, 190,
191, 193, 194, 195, 196, 203, 205, 208,
225, 226, 240, 246, 251, 257, 258, 268,
269, 270, 279, 284, 292, 293, 302, 303,
304, 308, 309, 315, 328, 329, 332, 338,
339, 353, 356, 357, 368, 371, 373, 376,
384, 387, 388, 394, 395, 396, 399, 400,
402, 404, 411, 414, 420, 425, 428, 433,
434, 436, 443, 446, 447, 453, 456, 457,
461, 470, 473, 474, 480, 481, 488, 489,
492, 493, 494, 502, 514, 519, 520, 531
- Winter, 26, 27, 29, 188, 198, 203, 221, 227,
235, 261, 265, 276, 281, 305, 344, 347,
355, 374, 390, 454, 468, 472, 477, 483
- Wisnouch, 510
- Wolf, 167
- Wolf, 64, 161, 417
- Wollmann, 321
- Wright, 161, 164, 167, 403, 404
- Zeidler, 498, 500
- Zettinow, 410, 475, 476
- Zopf, 28, 30, 44, 53, 189, 199, 203, 204,
206, 208, 221, 222, 227, 228, 235, 259,
260, 264, 267, 282, 283, 345, 347, 351,
355, 364, 375, 410, 414, 423, 446, 455,
472, 493, 524
- Zuber, 230
- Zuelzer, 479, 480
- Zukal, 222, 255, 394, 430

Sans Tache



All of these processes, except that of binding into cloth or leather covers, are carried on under our roof.

The motto of the Williams & Wilkins Company is *Sans Tache*. Our ideal is to publish books "*without blemish*"—worthy books, worthily printed, with worthy typography—books to which we shall be proud to attach our imprint, made by craftsmen who are willing to accept open responsibility for their work, and who are entitled to credit for creditable performance.

The printing craftsman of today is quite as much a craftsman as his predecessor. There is quite as much discrimination between poor work and good. We are of the opinion that the individuality of the worker should not be wholly lost. The members of our staff who have contributed their skill of hand and brain to this volume are

Composing Zeddie Breithaupt, Howard Fogle, Harry Harmeyer, Kurt Woernle, William Kidner, Ray Kauffman, Benjamin Hatcher, Thomas Loving, Herbert Leitch, William Fite, Harry LaMotte, Edward Poland Stultz, John Dotterweich, William Koch, George Behr, Ernest Salgado, William Sanders, George Moss, Andrew Rassa, Frank Zeller, Walter Phillips, Richard King, Henry Shea

Press August Hildebrand, Raymond Gallagher, Robert Gallagher, Clarence Ridgeway, Henry Eckert, Henry Augsburg

Keyboard Minnie Goad, Harry Susemihl, Vera Taylor, Anna Kelly, Kathryn Wilson, Anna Thomas, Eleanor Luecke, Catherine Kocent, Hannah Scott

Custer Kenneth Brown, Ernest Wann, Charles Aher, Mahlon Robinson, George Smith, Theodore Nilson, Oscar Hellung, Frank Malanosky

Folder Lawrence Krug

Proof Room Sarah Katzin, Mary Reed, Alice Reuter, Ruth Trieschman, Ethel Strasinger, Lucille Bull, Angeline Eifert, Audrey Tanner, Edna Clark, Lewis Forney, Dorothy Strasinger, Lillian Gilland, Arthur Baker

